



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2021 – 12:17 PM EDT

PDB ID : 1BIS  
Title : HIV-1 INTEGRASE CORE DOMAIN  
Authors : Goldgur, Y.; Dyda, F.; Hickman, A.B.; Jenkins, T.M.; Craigie, R.; Davies, D.R.  
Deposited on : 1998-06-19  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	 5% 66% 20% • 12%
1	B	166	 12% 68% 20% • • 7%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3274 atoms, of which 820 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HIV-1 INTEGRASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	146	1370	713	250	194	208	5	0	0	0
1	B	154	1448	751	266	205	221	5	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	48	SER	GLU	conflict	UNP P12497
A	49	HIS	ALA	conflict	UNP P12497
A	131	GLU	TRP	engineered mutation	UNP P12497
A	185	LYS	PHE	engineered mutation	UNP P12497
B	48	SER	GLU	conflict	UNP P12497
B	49	HIS	ALA	conflict	UNP P12497
B	131	GLU	TRP	engineered mutation	UNP P12497
B	185	LYS	PHE	engineered mutation	UNP P12497

- Molecule 2 is water.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	H	O		
2	A	80	240	160	80	0	0
2	B	72	216	144	72	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HIV-1 INTEGRASE

Chain A: 



- Molecule 1: HIV-1 INTEGRASE

Chain B: 



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.09Å 45.08Å 49.43Å 68.82° 64.74° 62.63°	Depositor
Resolution (Å)	(Not available) – 1.95 21.25 – 1.95	Depositor EDS
% Data completeness (in resolution range)	93.8 ((Not available)-1.95) 94.1 (21.25-1.95)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.09 (at 1.95Å)	Xtrriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.204 , 0.256 0.214 , 0.255	Depositor DCC
$R_{free}$ test set	1012 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtrriage
Anisotropy	0.205	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.017 for h,h-k,h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3274	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/1140	0.67	1/1540 (0.1%)
1	B	0.45	0/1205	0.68	0/1631
All	All	0.46	0/2345	0.68	1/3171 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	127	LYS	CD-CE-NZ	5.40	124.11	111.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1120	250	1134	26	0
1	B	1182	266	1190	30	1
2	A	80	160	0	5	1
2	B	72	144	0	8	0
All	All	2454	820	2324	55	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:GLN:HG3	2:B:669:HOH:O	1.75	0.85
1:B:124:THR:HA	2:B:738:HOH:O	1.87	0.73
1:B:178:MET:HE1	2:B:1005:HOH:O	1.94	0.66
1:B:84:ILE:HD11	1:B:158:LEU:HD13	1.77	0.65
1:B:56:CYS:HG	1:B:143:TYR:HH	1.46	0.63
1:A:127:LYS:HE2	2:A:856:HOH:O	1.99	0.61
1:B:79:VAL:HG22	1:B:151:ILE:HD13	1.84	0.60
1:B:118:GLY:HA3	2:B:931:HOH:O	2.04	0.57
1:B:144:ASN:N	1:B:145:PRO:HD3	2.23	0.54
1:B:103:LYS:HD3	2:B:669:HOH:O	2.08	0.54
1:B:181:PHE:O	1:B:185:LYS:HG2	2.10	0.52
1:A:204:ILE:O	1:A:208:ILE:HG23	2.09	0.52
1:B:143:TYR:CD1	1:B:143:TYR:N	2.77	0.51
1:B:127:LYS:HD2	2:B:738:HOH:O	2.11	0.50
1:A:84:ILE:HD11	1:A:158:LEU:HD13	1.93	0.50
1:A:82:GLY:O	1:A:183:HIS:HE1	1.94	0.50
1:B:208:ILE:O	1:B:208:ILE:HG22	2.12	0.49
1:A:62:GLN:HG3	1:A:79:VAL:HG22	1.95	0.49
1:B:79:VAL:HG11	1:B:147:SER:HB3	1.95	0.48
1:A:156:LYS:HD3	1:A:157:GLU:N	2.29	0.48
1:B:79:VAL:CG1	1:B:147:SER:HB3	2.44	0.47
1:B:121:PHE:HE2	2:B:934:HOH:O	1.96	0.47
1:B:107:ARG:HG3	1:B:108:TRP:CD1	2.49	0.47
1:B:82:GLY:O	1:B:183:HIS:HE1	1.99	0.46
1:A:200:ILE:HG12	1:A:204:ILE:HD12	1.97	0.45
1:B:62:GLN:HG2	1:B:114:HIS:HB2	1.99	0.45
1:A:151:ILE:HG23	1:A:152:GLU:H	1.82	0.44
1:B:125:THR:HG22	2:B:878:HOH:O	2.16	0.44
1:B:146:GLN:NE2	1:B:146:GLN:HA	2.33	0.44
1:A:140:GLY:HA2	2:A:483:HOH:O	2.18	0.44
1:B:74:LEU:HD22	1:B:89:ILE:HD13	2.00	0.44
1:A:165:VAL:HG11	1:A:182:ILE:CD1	2.48	0.43
1:B:202:ASP:O	1:B:206:THR:HG23	2.19	0.43
1:A:72:VAL:HG13	1:A:89:ILE:O	2.19	0.43
1:A:181:PHE:O	1:A:185:LYS:HG2	2.18	0.43
1:B:68:LEU:HD22	1:B:69:GLU:HG3	2.00	0.43
1:B:142:PRO:O	1:B:145:PRO:HD3	2.19	0.43
1:A:151:ILE:HG23	1:A:152:GLU:N	2.33	0.42
1:B:74:LEU:HD21	1:B:97:THR:HA	2.01	0.42
1:A:205:ALA:O	1:A:208:ILE:HG12	2.20	0.42
1:A:178:MET:HE2	1:B:132:TRP:CE2	2.55	0.42
1:A:142:PRO:CD	2:A:537:HOH:O	2.68	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:PRO:HG3	2:A:637:HOH:O	2.20	0.42
1:A:156:LYS:HD3	1:A:157:GLU:H	1.83	0.41
1:A:84:ILE:CD1	1:A:158:LEU:HD13	2.50	0.41
1:B:146:GLN:NE2	1:B:148:GLN:OE1	2.51	0.41
1:B:147:SER:O	1:B:151:ILE:HB	2.20	0.41
1:B:204:ILE:O	1:B:208:ILE:HG13	2.20	0.41
1:A:65:CYS:SG	1:A:74:LEU:HD13	2.61	0.41
1:A:158:LEU:O	1:A:162:ILE:HG13	2.20	0.41
1:B:56:CYS:SG	1:B:143:TYR:OH	2.64	0.41
1:A:63:LEU:HD11	1:A:100:PHE:CE2	2.56	0.41
1:A:142:PRO:HD3	2:A:537:HOH:O	2.20	0.41
1:A:57:SER:HA	1:A:58:PRO:HD3	1.89	0.40
1:A:172:LEU:O	1:A:176:VAL:HG23	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:LYS:HZ2	2:A:815:HOH:H1[1_554]	1.15	0.45

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/166 (86%)	138 (97%)	3 (2%)	1 (1%)	22	11
1	B	152/166 (92%)	147 (97%)	1 (1%)	4 (3%)	5	1
All	All	294/332 (89%)	285 (97%)	4 (1%)	5 (2%)	9	2

All (5) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	B	143	TYR
1	B	145	PRO
1	B	146	GLN
1	A	152	GLU
1	B	142	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/135 (87%)	113 (96%)	5 (4%)	30	17
1	B	125/135 (93%)	117 (94%)	8 (6%)	17	6
All	All	243/270 (90%)	230 (95%)	13 (5%)	22	10

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	111	LYS
1	A	131	GLU
1	A	156	LYS
1	A	209	GLN
1	B	62	GLN
1	B	65	CYS
1	B	68	LEU
1	B	107	ARG
1	B	111	LYS
1	B	146	GLN
1	B	154	MET
1	B	209	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	67	HIS
1	A	183	HIS
1	B	137	GLN
1	B	183	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	146/166 (87%)	0.56	9 (6%) 20 29	13, 25, 54, 84	0
1	B	154/166 (92%)	0.87	20 (12%) 3 5	14, 26, 77, 93	0
All	All	300/332 (90%)	0.72	29 (9%) 7 12	13, 25, 70, 93	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	150	VAL	11.5
1	A	56	CYS	9.9
1	B	144	ASN	9.8
1	A	151	ILE	8.7
1	B	143	TYR	8.4
1	B	148	GLN	6.2
1	B	147	SER	5.9
1	B	146	GLN	5.4
1	B	151	ILE	5.3
1	B	141	ILE	5.3
1	B	149	GLY	5.2
1	A	152	GLU	4.1
1	B	145	PRO	4.1
1	A	208	ILE	3.9
1	B	153	SER	3.8
1	A	153	SER	3.5
1	B	152	GLU	3.4
1	B	156	LYS	3.3
1	A	156	LYS	3.1
1	A	209	GLN	3.0
1	A	141	ILE	2.9
1	B	70	GLY	2.8
1	B	56	CYS	2.8
1	A	140	GLY	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	208	ILE	2.6
1	B	121	PHE	2.5
1	B	209	GLN	2.4
1	B	154	MET	2.4
1	B	158	LEU	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.