



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 02:53 PM EST

PDB ID : 1KOL
Title : Crystal structure of formaldehyde dehydrogenase
Authors : Tanaka, N.; Kusakabe, Y.; Ito, K.; Yoshimoto, T.; Nakamura, K.T.
Deposited on : 2001-12-21
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	398	 90% 8% ..
1	B	398	 89% 10% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called formaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	396	2934	1848	521	549	16	0	0	0
1	B	396	2934	1848	521	549	16	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	VAL	LEU	SEE REMARK 999	UNP P46154
B	139	VAL	LEU	SEE REMARK 999	UNP P46154

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	B	1	44	21	7	14	2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	472	Total	O	0	0
			472	472		
5	B	495	Total	O	0	0
			495	495		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: formaldehyde dehydrogenase



- Molecule 1: formaldehyde dehydrogenase



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	85.69Å 85.69Å 190.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.65	Depositor
% Data completeness (in resolution range)	(Not available) (30.00-1.65)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.171 , 0.206	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6937	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, ZN, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2991	1.10	9/4058 (0.2%)
1	B	0.48	0/2991	1.12	11/4058 (0.3%)
All	All	0.47	0/5982	1.11	20/8116 (0.2%)

There are no bond length outliers.

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	267	ARG	CD-NE-CZ	12.96	141.75	123.60
1	B	146	PHE	CB-CG-CD2	-7.52	115.54	120.80
1	B	101	ARG	NE-CZ-NH2	-7.10	116.75	120.30
1	B	355	ARG	CD-NE-CZ	7.06	133.48	123.60
1	A	162	ARG	NE-CZ-NH1	6.93	123.77	120.30
1	A	291	ARG	NE-CZ-NH2	-6.77	116.91	120.30
1	A	345	ARG	NE-CZ-NH2	-6.47	117.06	120.30
1	B	291	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	267	ARG	NE-CZ-NH1	6.37	123.49	120.30
1	A	374	ARG	NE-CZ-NH2	6.04	123.32	120.30
1	B	206	ARG	NE-CZ-NH1	-5.91	117.35	120.30
1	A	4	ARG	NE-CZ-NH1	-5.59	117.50	120.30
1	A	374	ARG	CD-NE-CZ	5.40	131.16	123.60
1	B	206	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	355	ARG	CD-NE-CZ	5.29	131.00	123.60
1	B	374	ARG	N-CA-CB	5.25	120.04	110.60
1	B	101	ARG	NH1-CZ-NH2	5.15	125.06	119.40
1	B	374	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	B	88	LEU	CA-CB-CG	5.11	127.05	115.30
1	B	206	ARG	NH1-CZ-NH2	5.09	125.00	119.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2934	0	2936	21	0
1	B	2934	0	2936	22	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	44	0	26	1	0
4	B	44	0	26	1	0
5	A	472	0	0	1	1
5	B	495	0	0	5	0
All	All	6937	0	5924	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:393:LYS:HE3	1:A:393:LYS:HA	1.56	0.85
1:A:34:HIS:HD2	1:A:143:TYR:H	1.31	0.77
1:B:34:HIS:HD2	1:B:143:TYR:H	1.37	0.72
1:A:374:ARG:HH21	1:A:374:ARG:HG2	1.55	0.71
1:A:34:HIS:CD2	1:A:143:TYR:H	2.08	0.71
1:B:277:GLU:HG2	5:B:2780:HOH:O	1.92	0.70
1:A:88:LEU:HG	1:A:151:LEU:HD12	1.76	0.65
1:A:393:LYS:HE3	1:A:393:LYS:CA	2.25	0.65
1:A:393:LYS:HE2	1:A:396:SER:OG	1.98	0.64
1:B:34:HIS:CD2	1:B:143:TYR:H	2.16	0.64
1:B:243:HIS:HB3	1:B:288:GLN:NE2	2.14	0.62
1:B:374:ARG:HG3	5:B:2855:HOH:O	2.01	0.61
1:B:243:HIS:HB3	1:B:288:GLN:HE22	1.70	0.56
1:A:26:ASP:HB2	1:A:27:PRO:CD	2.37	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:LYS:HE3	5:B:2808:HOH:O	2.07	0.54
1:A:288:GLN:HE21	1:A:288:GLN:HA	1.77	0.50
1:A:299:PRO:O	4:A:1403:NAD:H2N	2.13	0.49
1:B:299:PRO:O	4:B:2403:NAD:H2N	2.13	0.49
1:A:9:LEU:HD11	1:A:15:GLU:HB3	1.95	0.49
1:B:225:HIS:HE1	1:B:361:VAL:O	1.95	0.48
1:A:46:CYS:HB2	1:A:68:GLU:OE1	2.14	0.48
1:A:371:ASP:O	1:A:374:ARG:HG3	2.13	0.48
1:A:225:HIS:HE1	1:A:361:VAL:O	1.96	0.48
1:B:11:SER:O	1:B:13:LYS:HD2	2.14	0.47
1:B:227:LYS:HE3	5:B:2734:HOH:O	2.14	0.47
1:B:9:LEU:HD11	1:B:15:GLU:HB3	1.98	0.46
1:B:43:THR:HA	1:B:68:GLU:O	2.16	0.46
1:A:374:ARG:HG2	1:A:374:ARG:NH2	2.22	0.45
1:A:193:GLY:O	1:A:198:GLY:HA3	2.16	0.45
1:B:25:GLN:NE2	1:B:29:GLY:HA2	2.32	0.45
1:A:393:LYS:HA	1:A:393:LYS:CE	2.38	0.44
1:B:360:GLU:HG3	5:B:2511:HOH:O	2.18	0.44
1:B:374:ARG:O	1:B:378:GLU:HG3	2.19	0.43
1:B:293:ALA:HA	1:B:330:SER:O	2.19	0.43
1:B:203:ALA:HB2	1:B:361:VAL:HG11	2.00	0.42
1:B:26:ASP:HB2	1:B:27:PRO:CD	2.50	0.42
1:B:193:GLY:O	1:B:198:GLY:HA3	2.19	0.42
1:A:152:PRO:O	1:A:153:ASP:C	2.58	0.42
1:A:366:VAL:HA	1:A:388:VAL:O	2.20	0.41
1:B:46:CYS:HB2	1:B:68:GLU:OE1	2.20	0.41
1:B:78:ARG:HH11	1:B:78:ARG:HD3	1.70	0.41
1:A:277:GLU:HG2	5:A:1832:HOH:O	2.20	0.40
1:A:43:THR:HA	1:A:68:GLU:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1524:HOH:O	5:A:1630:HOH:O[4_555]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	394/398 (99%)	382 (97%)	12 (3%)	0	100	100
1	B	394/398 (99%)	386 (98%)	8 (2%)	0	100	100
All	All	788/796 (99%)	768 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/304 (100%)	291 (96%)	12 (4%)	31	9
1	B	303/304 (100%)	296 (98%)	7 (2%)	50	25
All	All	606/608 (100%)	587 (97%)	19 (3%)	40	14

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLU
1	A	18	LYS
1	A	30	LYS
1	A	58	THR
1	A	153	ASP
1	A	267	ARG
1	A	288	GLN
1	A	316	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	319	LEU
1	A	374	ARG
1	A	391	PRO
1	A	393	LYS
1	B	44	ASN
1	B	60	GLN
1	B	88	LEU
1	B	218	LEU
1	B	319	LEU
1	B	354	ASP
1	B	374	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	HIS
1	A	225	HIS
1	A	288	GLN
1	A	331	HIS
1	B	34	HIS
1	B	60	GLN
1	B	225	HIS
1	B	331	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	1002	-	4,4,4	0.63	0	6,6,6	0.27	0
3	SO4	A	1003	-	4,4,4	0.65	0	6,6,6	0.17	0
4	NAD	A	1403	-	42,48,48	1.26	6 (14%)	50,73,73	2.11	11 (22%)
4	NAD	B	2403	-	42,48,48	1.25	3 (7%)	50,73,73	1.99	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAD	B	2403	-	-	5/26/62/62	0/5/5/5
4	NAD	A	1403	-	-	7/26/62/62	0/5/5/5

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1403	NAD	C3N-C7N	4.23	1.56	1.50
4	B	2403	NAD	C3N-C7N	3.83	1.56	1.50
4	B	2403	NAD	O4D-C1D	3.07	1.45	1.41
4	B	2403	NAD	C6N-N1N	2.61	1.41	1.35
4	A	1403	NAD	C6N-N1N	2.40	1.41	1.35
4	A	1403	NAD	O4D-C1D	2.30	1.44	1.41
4	A	1403	NAD	C2N-C3N	-2.14	1.35	1.39
4	A	1403	NAD	PN-O2N	-2.02	1.45	1.55
4	A	1403	NAD	C2N-N1N	2.00	1.37	1.35

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1403	NAD	C5N-C4N-C3N	-7.80	111.11	120.34
4	B	2403	NAD	C5N-C4N-C3N	-7.31	111.69	120.34
4	A	1403	NAD	C6N-C5N-C4N	5.69	127.70	119.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	2403	NAD	C6N-C5N-C4N	5.54	127.48	119.44
4	B	2403	NAD	C2N-C3N-C4N	5.28	124.24	118.26
4	A	1403	NAD	C2N-C3N-C4N	5.16	124.11	118.26
4	A	1403	NAD	O4D-C1D-C2D	-4.20	100.79	106.93
4	A	1403	NAD	C5N-C6N-N1N	-3.85	114.88	120.40
4	B	2403	NAD	O4D-C1D-C2D	-3.80	101.37	106.93
4	B	2403	NAD	C5A-C6A-N6A	3.54	125.73	120.35
4	B	2403	NAD	C5N-C6N-N1N	-3.24	115.75	120.40
4	B	2403	NAD	C4N-C3N-C7N	-3.18	112.54	121.04
4	A	1403	NAD	C4N-C3N-C7N	-3.13	112.67	121.04
4	A	1403	NAD	C5A-C6A-N6A	3.09	125.04	120.35
4	A	1403	NAD	C2N-N1N-C1D	-2.71	113.10	119.14
4	A	1403	NAD	O4B-C1B-C2B	-2.68	103.01	106.93
4	A	1403	NAD	C1B-N9A-C4A	-2.28	122.64	126.64
4	B	2403	NAD	O4B-C1B-C2B	-2.21	103.70	106.93
4	A	1403	NAD	O7N-C7N-N7N	2.06	125.50	122.58

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1403	NAD	O4D-C1D-N1N-C2N
4	A	1403	NAD	O4D-C1D-N1N-C6N
4	A	1403	NAD	C2D-C1D-N1N-C2N
4	A	1403	NAD	C2D-C1D-N1N-C6N
4	B	2403	NAD	O4D-C1D-N1N-C2N
4	B	2403	NAD	O4D-C1D-N1N-C6N
4	B	2403	NAD	C2D-C1D-N1N-C2N
4	B	2403	NAD	C2D-C1D-N1N-C6N
4	A	1403	NAD	C5B-O5B-PA-O3
4	B	2403	NAD	O4B-C4B-C5B-O5B
4	A	1403	NAD	C5B-O5B-PA-O1A
4	A	1403	NAD	O4B-C4B-C5B-O5B


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1403	NAD	1	0
4	B	2403	NAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.