



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 10:06 AM EDT

PDB ID : 1NFK
Title : STRUCTURE OF THE NUCLEAR FACTOR KAPPA-B (NF-KB) P50 HOMODIMER
Authors : Ghosh, G.; Van Duyne, G.; Ghosh, S.; Sigler, P.B.
Deposited on : 1995-02-28
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	11	45% (yellow) 55% (orange)
1	D	11	64% (yellow) 36% (orange)
2	A	325	52% (green) 35% (yellow) 7% (orange) 7% (red) 7% (grey)
2	B	325	41% (green) 45% (yellow) 9% (orange) 5% (red) 5% (grey)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7477 atoms, of which 1810 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*GP*GP*GP*AP*AP*TP*TP*CP*CP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	C	11	Total	C	H	N	O	P	0	0	0
			246	107	24	40	65	10			
1	D	11	Total	C	H	N	O	P	0	0	0
			246	107	24	40	65	10			

- Molecule 2 is a protein called PROTEIN (NUCLEAR FACTOR KAPPA-B (NF-KB)).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	A	312	Total	C	H	N	O	S	0	0	0
			3017	1554	564	428	459	12			
2	B	312	Total	C	H	N	O	S	0	0	0
			3017	1554	564	428	459	12			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	25	Total	H	O	0	0
			75	50	25		
3	D	26	Total	H	O	0	0
			78	52	26		
3	A	152	Total	H	O	0	0
			456	304	152		
3	B	114	Total	H	O	0	0
			342	228	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*AP*AP*TP*TP*CP*CP*C)-3')

Chain C: 




- Molecule 1: DNA (5'-D(*TP*GP*GP*GP*AP*AP*TP*TP*CP*CP*C)-3')

Chain D: 




- Molecule 2: PROTEIN (NUCLEAR FACTOR KAPPA-B (NF-KB))

Chain A: 



- Molecule 2: PROTEIN (NUCLEAR FACTOR KAPPA-B (NF-KB))

Chain B: 



A108	A185	G262	P342
H109	L186	G263	ILE
S110	T187	E264	LYS
L111	D188	E265	LYS
V112	R189	I266	LYS
G113	E190	I267	GLU
K114	K191	L268	GLU
H115	E192	D276	VAL
C116	I193	I194	GLN
C121	I194	I195	ARG
T122	R195	I196	ARG
V123	Q196	V199	GLN
T124	V199	Q200	LYS
A125	Q200	Q201	LYS
G126	Q201	T202	LYS
V132	T202	K203	LYS
A135	K203	F204	ARG
M136	F204	M205	GLN
L137	M205	D206	LYS
G138	D206	L207	LYS
I139	L207	R211	LYS
L140	R211	L212	LYS
H141	L212	A216	LYS
V142	A216	F217	LYS
T143	F217	L218	LYS
K144	L218	P219	LYS
K145	P219	D220	LYS
K146	D220	S221	LYS
V147	S221	T222	LYS
F148	T222	G223	LYS
E149	G223	F225	LYS
T150	F225	T226	LYS
A153	T226	R227	LYS
R154	R227	R228	LYS
M155	R228	L229	LYS
T156	L229	V232	LYS
E157	V232	V233	LYS
A158	V233	S234	LYS
C159	S234	D235	LYS
I160	D235	A236	LYS
M164	A236	S240	LYS
P165	S240	K241	LYS
G166	K241	L248	LYS
L167	L248	R252	LYS
L168	R252	M253	LYS
D172	M253	D254	LYS
L173	D254	R255	LYS
A174	R255	T256	LYS
Y175	T256	V260	LYS
L176	V260	T261	LYS
Q177	T261		
A178			
R184			

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	84.20Å 132.10Å 80.10Å 90.00° 93.10° 90.00°	Depositor
Resolution (Å)	6.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.230 , 0.340	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7477	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	C	2.99	26/248 (10.5%)	2.09	11/381 (2.9%)
1	D	3.47	38/248 (15.3%)	2.18	14/381 (3.7%)
2	A	0.60	0/2505	0.89	4/3384 (0.1%)
2	B	0.55	0/2505	0.84	3/3384 (0.1%)
All	All	1.12	64/5506 (1.2%)	1.06	32/7530 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

The worst 5 of 64 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	7	DT	C5-C7	12.11	1.57	1.50
1	D	7	DT	C5'-C4'	-11.19	1.39	1.51
1	C	1	DT	N3-C4	-10.50	1.30	1.38
1	D	6	DA	N9-C4	10.27	1.44	1.37
1	C	8	DT	C5'-C4'	-9.51	1.40	1.51

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	8	DT	C4-C5-C7	-9.73	113.16	119.00
1	D	5	DA	O4'-C4'-C3'	-7.38	101.55	104.50
1	D	5	DA	O4'-C1'-N9	7.32	113.12	108.00
1	C	11	DC	O4'-C1'-N1	7.18	113.03	108.00
1	C	7	DT	C4-C5-C7	-6.69	114.98	119.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	175	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	222	24	125	5	0
1	D	222	24	126	7	0
2	A	2453	564	2449	121	0
2	B	2453	564	2451	152	0
3	A	152	304	0	10	0
3	B	114	228	0	4	0
3	C	25	50	0	1	0
3	D	26	52	0	2	0
All	All	5667	1810	5151	275	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 275 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:9:DC:H2'	2:A:59:CYS:SG	1.97	1.05
1:C:1:DT:H2''	1:C:2:DG:H5'	1.42	0.99
2:B:218:LEU:HB2	2:B:227:ARG:HB3	1.41	0.99
2:A:92:LYS:HG2	2:A:124:THR:HG22	1.44	0.98
2:B:147:VAL:HG12	2:B:202:THR:HG22	1.54	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	310/325 (95%)	257 (83%)	35 (11%)	18 (6%)	1	0
2	B	310/325 (95%)	251 (81%)	39 (13%)	20 (6%)	1	0
All	All	620/650 (95%)	508 (82%)	74 (12%)	38 (6%)	1	0

5 of 38 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	87	TYR
2	A	172	ASP
2	A	261	THR
2	A	317	LYS
2	A	322	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	268/281 (95%)	228 (85%)	40 (15%)	3	3
2	B	268/281 (95%)	241 (90%)	27 (10%)	7	9
All	All	536/562 (95%)	469 (88%)	67 (12%)	4	5

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	207	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	248	LEU
2	B	316	TYR
2	A	213	MET
2	A	210	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	43	GLN
2	B	141	HIS
2	B	75	ASN
2	B	201	GLN
2	B	109	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.