



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2021 – 10:42 PM EDT

PDB ID : 1U12
Title : M. loti cyclic nucleotide binding domain mutant
Authors : Clayton, G.M.; Silverman, W.R.; Heginbotham, L.; Morais-Cabral, J.H.
Deposited on : 2004-07-14
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	138	 7% (poor fit), 52% (0 outliers), 33% (1 outlier), 7% (2 outliers), 5% (3+ outliers)
1	B	138	 2% (poor fit), 58% (0 outliers), 28% (1 outlier), 5% (2 outliers), 8% (3+ outliers)

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	IOD	A	132	-	-	X	-
2	IOD	B	211	-	-	X	-
2	IOD	B	911	-	-	X	-
2	IOD	B	913	-	-	X	-
2	IOD	B	918	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1918 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cyclic nucleotide binding domain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	131	942	604	160	173	5	0	0	0
1	B	127	920	587	159	169	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	348	ALA	ARG	engineered mutation	UNP Q98GN8
B	348	ALA	ARG	engineered mutation	UNP Q98GN8

- Molecule 2 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	12	Total	I	0	0
			12	12		
2	B	19	Total	I	0	0
			19	19		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total K 3 3	0	0
4	B	1	Total K 1 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total O 2 2	0	0
5	B	9	Total O 9 9	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cyclic nucleotide binding domain

Chain A: 7% 52% 33% 7% 5%



- Molecule 1: cyclic nucleotide binding domain

Chain B: 2% 58% 28% 5% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	57.13Å 57.13Å 193.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.39 – 2.70 40.39 – 2.60	Depositor EDS
% Data completeness (in resolution range)	96.5 (40.39-2.70) 98.6 (40.39-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 2.61Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.253 , 0.281 0.275 , 0.289	Depositor DCC
R_{free} test set	810 reflections (4.35%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtrriage
Anisotropy	0.559	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	1918	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, K, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/961	1.21	8/1314 (0.6%)
1	B	0.54	0/938	1.05	5/1281 (0.4%)
All	All	0.55	0/1899	1.14	13/2595 (0.5%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	287	PRO	CA-N-CD	-14.32	91.45	111.50
1	B	287	PRO	CA-N-CD	-13.57	92.51	111.50
1	A	285	PRO	CA-N-CD	-13.15	93.09	111.50
1	A	306	PRO	CA-N-CD	-13.13	93.12	111.50
1	B	234	PRO	CA-N-CD	-12.52	93.97	111.50
1	A	268	PRO	CA-N-CD	-11.14	95.90	111.50
1	B	285	PRO	CA-N-CD	-10.56	96.72	111.50
1	A	335	PRO	CA-N-CD	-10.11	97.35	111.50
1	A	257	PRO	CA-N-CD	-9.82	97.75	111.50
1	A	241	PRO	CA-N-CD	-9.23	98.58	111.50
1	A	292	PRO	CA-N-CD	-9.05	98.83	111.50
1	B	306	PRO	CA-N-CD	-7.76	100.64	111.50
1	B	234	PRO	CA-C-N	-7.00	101.81	117.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	942	0	928	40	0
1	B	920	0	923	47	0
2	A	12	0	0	6	0
2	B	19	0	0	15	0
3	A	10	0	0	0	0
4	A	3	0	0	0	0
4	B	1	0	0	0	0
5	A	2	0	0	1	0
5	B	9	0	0	2	0
All	All	1918	0	1851	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:344:THR:HG22	1:B:344:THR:O	1.56	1.06
1:B:279:SER:H	1:B:315:THR:HG22	1.33	0.92
1:B:234:PRO:HG2	2:B:911:IOD:I	2.40	0.92
1:A:303:SER:OG	2:A:192:IOD:I	2.62	0.88
2:B:914:IOD:I	2:B:918:IOD:I	3.32	0.87
1:A:331:CYS:SG	1:A:338:ALA:HB2	2.17	0.85
1:A:264:ARG:O	1:A:267:GLU:HG2	1.78	0.84
1:A:239:LEU:HD21	1:A:340:ILE:HD12	1.60	0.82
1:B:264:ARG:O	1:B:267:GLU:HG2	1.79	0.80
1:A:249:ARG:NH1	2:A:711:IOD:I	2.84	0.80
1:B:249:ARG:HG2	2:B:918:IOD:I	2.53	0.79
1:B:344:THR:O	1:B:344:THR:CG2	2.29	0.78
1:A:245:VAL:HG21	1:B:245:VAL:HG11	1.65	0.78
1:B:242:ALA:HB3	2:B:211:IOD:I	2.57	0.74
1:A:263:CYS:SG	1:A:307:ARG:HD3	2.30	0.72
1:B:306:PRO:HD3	2:B:913:IOD:I	2.60	0.71
1:B:300:ALA:HB2	1:B:307:ARG:NH2	2.06	0.71
1:B:336:GLU:O	1:B:340:ILE:HG12	1.89	0.71
1:B:245:VAL:O	1:B:249:ARG:HG3	1.91	0.71
1:A:331:CYS:O	1:A:335:PRO:HD3	1.95	0.67
1:A:235:LEU:HD11	1:A:298:GLU:HG2	1.77	0.64
1:B:234:PRO:HB2	2:B:911:IOD:I	2.67	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:226:ASN:ND2	1:B:320:LEU:HD11	2.12	0.64
1:A:328:GLN:NE2	2:A:132:IOD:I	3.00	0.64
1:B:234:PRO:CG	2:B:911:IOD:I	3.15	0.64
1:A:315:THR:O	1:A:317:VAL:HG23	1.98	0.63
1:B:343:LYS:O	1:B:345:ALA:N	2.32	0.63
1:B:263:CYS:SG	1:B:307:ARG:HD3	2.41	0.61
1:A:239:LEU:CD2	1:A:340:ILE:HD12	2.33	0.58
1:B:262:ILE:HD13	1:B:319:LEU:HD11	1.85	0.58
1:A:241:PRO:O	1:A:245:VAL:HG23	2.04	0.58
1:A:277:GLU:HA	1:A:292:PRO:HG3	1.85	0.58
1:B:324:SER:O	1:B:328:GLN:HG3	2.03	0.57
1:A:343:LYS:O	1:A:347:GLU:HG3	2.05	0.57
1:A:278:GLY:O	1:A:292:PRO:HD3	2.04	0.56
1:B:299:MET:HE3	1:B:302:ILE:HG12	1.86	0.56
1:B:279:SER:H	1:B:315:THR:CG2	2.13	0.56
1:B:256:VAL:HB	1:B:317:VAL:HG23	1.88	0.56
1:B:300:ALA:HB2	1:B:307:ARG:HH21	1.71	0.55
1:A:272:MET:HE3	1:A:298:GLU:HG3	1.89	0.55
1:B:297:GLY:HA2	2:B:162:IOD:I	2.78	0.54
1:A:302:ILE:HD13	1:A:342:ARG:CB	2.40	0.52
1:B:242:ALA:CB	2:B:211:IOD:I	3.28	0.52
2:B:915:IOD:I	5:B:8:HOH:O	2.90	0.51
1:A:227:TRP:CH2	1:A:244:LEU:HB2	2.45	0.51
1:A:326:ASP:O	1:A:329:MET:HG3	2.11	0.51
1:B:306:PRO:CD	2:B:913:IOD:I	3.29	0.51
1:A:296:PHE:HA	1:A:298:GLU:OE2	2.11	0.51
1:B:324:SER:C	1:B:328:GLN:HG3	2.31	0.50
1:B:226:ASN:HD21	1:B:320:LEU:HD11	1.74	0.50
1:B:276:VAL:HG21	1:B:320:LEU:HG	1.94	0.50
1:B:276:VAL:CG2	1:B:320:LEU:HG	2.41	0.49
1:B:315:THR:HG23	1:B:316:THR:O	2.12	0.49
1:A:225:ARG:HD2	1:A:226:ASN:N	2.29	0.48
1:A:253:ALA:C	1:A:254:ARG:HG2	2.35	0.47
1:A:279:SER:O	1:A:317:VAL:HG21	2.14	0.47
1:B:268:PRO:HG3	2:B:913:IOD:I	2.85	0.47
1:A:225:ARG:HD2	1:A:225:ARG:C	2.34	0.47
1:B:234:PRO:CB	2:B:911:IOD:I	3.31	0.47
1:B:221:GLY:N	2:B:112:IOD:I	3.18	0.47
1:A:287:PRO:HB3	2:A:511:IOD:I	2.86	0.46
1:A:331:CYS:SG	1:A:338:ALA:N	2.89	0.45
1:A:325:ALA:HB2	2:A:132:IOD:I	2.86	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:306:PRO:HG3	2:B:913:IOD:I	2.86	0.45
1:B:256:VAL:CG1	1:B:260:ALA:HB3	2.47	0.45
1:A:222:ASP:HA	1:A:225:ARG:HG3	1.98	0.44
1:B:299:MET:HB3	5:B:7:HOH:O	2.17	0.44
1:A:331:CYS:SG	1:A:337:ILE:HB	2.58	0.43
1:B:281:SER:N	1:B:312:SER:O	2.43	0.43
1:A:301:LEU:HD22	1:A:324:SER:HA	2.00	0.43
1:B:237:GLN:HG3	1:B:238:LYS:N	2.34	0.43
1:A:246:GLU:HB3	1:A:330:LEU:HD22	2.01	0.42
1:A:347:GLU:HA	2:A:812:IOD:I	2.90	0.42
1:A:256:VAL:CG1	1:A:260:ALA:HB3	2.49	0.42
1:B:244:LEU:O	1:B:248:VAL:HG23	2.20	0.42
1:A:269:GLY:O	1:A:300:ALA:HB1	2.20	0.42
1:B:343:LYS:C	1:B:345:ALA:H	2.21	0.41
1:A:220:ARG:O	1:A:224:VAL:HG23	2.19	0.41
1:B:235:LEU:HD12	1:B:344:THR:HG22	2.03	0.41
1:A:331:CYS:SG	1:A:338:ALA:CB	2.98	0.41
1:B:313:ALA:HB1	1:B:317:VAL:HG22	2.01	0.41
1:B:224:VAL:HG23	1:B:225:ARG:N	2.36	0.41
1:A:324:SER:O	1:A:327:PHE:HB3	2.21	0.41
1:B:256:VAL:HA	1:B:257:PRO:HD3	1.92	0.41
1:A:242:ALA:HB3	5:A:9:HOH:O	2.20	0.40
1:B:226:ASN:ND2	1:B:320:LEU:CD1	2.83	0.40
1:B:224:VAL:CG2	1:B:225:ARG:N	2.84	0.40
1:A:279:SER:O	1:A:317:VAL:CG2	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	129/138 (94%)	122 (95%)	6 (5%)	1 (1%)	19 43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	125/138 (91%)	118 (94%)	6 (5%)	1 (1%)	19	43
All	All	254/276 (92%)	240 (94%)	12 (5%)	2 (1%)	19	43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	344	THR
1	A	284	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/106 (90%)	79 (83%)	16 (17%)	2	5
1	B	96/106 (91%)	87 (91%)	9 (9%)	8	20
All	All	191/212 (90%)	166 (87%)	25 (13%)	4	10

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	225	ARG
1	A	229	LEU
1	A	241	PRO
1	A	257	PRO
1	A	268	PRO
1	A	284	THR
1	A	285	PRO
1	A	287	PRO
1	A	292	PRO
1	A	298	GLU
1	A	302	ILE
1	A	306	PRO
1	A	307	ARG
1	A	329	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	335	PRO
1	A	339	GLU
1	B	234	PRO
1	B	247	ILE
1	B	285	PRO
1	B	287	PRO
1	B	302	ILE
1	B	306	PRO
1	B	307	ARG
1	B	316	THR
1	B	317	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	323	HIS
1	B	226	ASN
1	B	286	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 37 ligands modelled in this entry, 35 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	144	4	4,4,4	0.35	0	6,6,6	0.15	0
3	SO4	A	145	-	4,4,4	0.28	0	6,6,6	0.16	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/138 (94%)	0.52	10 (7%) 13 12	14, 31, 52, 60	0
1	B	127/138 (92%)	0.34	3 (2%) 59 60	13, 29, 46, 75	0
All	All	258/276 (93%)	0.43	13 (5%) 28 27	13, 30, 52, 75	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	347	GLU	6.2
1	A	303	SER	4.5
1	A	242	ALA	3.3
1	A	237	GLN	3.1
1	A	331	CYS	2.9
1	A	330	LEU	2.9
1	A	335	PRO	2.6
1	B	345	ALA	2.4
1	A	334	SER	2.2
1	A	243	VAL	2.1
1	A	241	PRO	2.1
1	A	238	LYS	2.1
1	B	264	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IOD	B	918	1/1	0.48	0.19	82,82,82,82	1
4	K	A	814	1/1	0.83	0.12	41,41,41,41	1
2	IOD	B	912	1/1	0.84	0.10	64,64,64,64	1
2	IOD	A	811	1/1	0.85	0.11	58,58,58,58	1
2	IOD	A	411	1/1	0.85	0.14	65,65,65,65	1
2	IOD	A	813	1/1	0.86	0.11	69,69,69,69	1
4	K	A	155	1/1	0.87	0.23	35,35,35,35	1
2	IOD	B	914	1/1	0.88	0.10	61,61,61,61	1
2	IOD	B	172	1/1	0.88	0.09	57,57,57,57	1
2	IOD	B	917	1/1	0.89	0.11	54,54,54,54	1
2	IOD	B	112	1/1	0.89	0.08	67,67,67,67	1
3	SO4	A	144	5/5	0.90	0.22	26,27,27,28	3
2	IOD	B	919	1/1	0.91	0.27	58,58,58,58	1
2	IOD	B	211	1/1	0.93	0.09	60,60,60,60	1
2	IOD	A	132	1/1	0.94	0.12	64,64,64,64	1
2	IOD	A	812	1/1	0.94	0.11	64,64,64,64	1
2	IOD	B	915	1/1	0.94	0.10	53,53,53,53	1
2	IOD	B	916	1/1	0.94	0.10	44,44,44,44	1
2	IOD	A	356	1/1	0.94	0.05	53,53,53,53	1
2	IOD	A	711	1/1	0.95	0.13	59,59,59,59	1
2	IOD	A	511	1/1	0.95	0.14	51,51,51,51	1
2	IOD	B	911	1/1	0.95	0.17	61,61,61,61	1
2	IOD	B	202	1/1	0.96	0.07	61,61,61,61	1
2	IOD	A	192	1/1	0.96	0.09	67,67,67,67	1
2	IOD	B	913	1/1	0.96	0.10	57,57,57,57	1
3	SO4	A	145	5/5	0.97	0.21	22,25,26,27	4
2	IOD	B	122	1/1	0.97	0.11	60,60,60,60	1
2	IOD	A	611	1/1	0.97	0.20	57,57,57,57	1
4	K	A	455	1/1	0.97	0.06	26,26,26,26	1
4	K	B	920	1/1	0.97	0.32	19,19,19,19	1
2	IOD	A	152	1/1	0.98	0.11	50,50,50,50	1
2	IOD	B	111	1/1	0.98	0.11	52,52,52,52	1
2	IOD	B	162	1/1	0.98	0.12	49,49,49,49	1
2	IOD	B	102	1/1	0.98	0.11	48,48,48,48	1
2	IOD	B	182	1/1	0.98	0.08	52,52,52,52	1
2	IOD	A	142	1/1	0.99	0.12	35,35,35,35	1
2	IOD	B	212	1/1	0.99	0.12	48,48,48,48	1

6.5 Other polymers

There are no such residues in this entry.