



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 10:09 AM EDT

PDB ID : 1XBV
Title : Crystal structure of 3-keto-L-gulonate 6-phosphate decarboxylase with bound D-ribulose 5-phosphate
Authors : Wise, E.L.; Yew, W.S.; Akana, J.; Gerlt, J.A.; Rayment, I.
Deposited on : 2004-08-31
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	
1	B	216	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-keto-L-gulonate 6-phosphate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	213	Total 1654	C 1048	N 290	O 308	S 8	0	2	0
1	B	215	Total 1632	C 1034	N 286	O 304	S 8	0	1	0

- Molecule 2 is RIBULOSE-5-PHOSPHATE (three-letter code: 5RP) (formula: C₅H₁₁O₈P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	Total 13	C 5	O 7	P 1	0	0
2	B	1	Total 13	C 5	O 7	P 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	B	1	Total Mg 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	205	Total O 205 205	0	0
4	B	220	Total O 220 220	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3-keto-L-gulonate 6-phosphate decarboxylase



- Molecule 1: 3-keto-L-gulonate 6-phosphate decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	123.81Å 41.47Å 91.18Å 90.00° 96.65° 90.00°	Depositor
Resolution (Å)	91.29 – 1.66 23.64 – 1.66	Depositor EDS
% Data completeness (in resolution range)	99.1 (91.29-1.66) 99.0 (23.64-1.66)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.66Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.162 , 0.194 0.176 , 0.207	Depositor DCC
R_{free} test set	2763 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtrriage
Anisotropy	0.404	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 53.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3739	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 5RP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.77	0/1698	0.95	6/2303 (0.3%)
1	B	0.88	1/1665 (0.1%)	1.00	6/2259 (0.3%)
All	All	0.83	1/3363 (0.0%)	0.97	12/4562 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	2	SER	C-N	7.40	1.51	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	2	SER	O-C-N	-12.50	102.70	122.70
1	A	23	ARG	NE-CZ-NH1	9.56	125.08	120.30
1	A	82	ASP	CB-CG-OD2	6.98	124.58	118.30
1	B	139	ARG	NE-CZ-NH2	-6.43	117.08	120.30
1	B	16	ASP	CB-CG-OD2	6.14	123.83	118.30
1	B	91	ASP	CB-CG-OD2	6.10	123.79	118.30
1	B	30	ASP	CB-CG-OD2	6.03	123.72	118.30
1	A	126	ARG	NE-CZ-NH2	-5.84	117.38	120.30
1	B	45	ARG	NE-CZ-NH1	5.65	123.12	120.30
1	A	152	ASP	CB-CG-OD2	5.40	123.16	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	100	ASP	CB-CG-OD2	5.25	123.02	118.30
1	A	23	ARG	NE-CZ-NH2	-5.05	117.77	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	2	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1654	0	1663	8	1
1	B	1632	0	1627	8	1
2	A	13	0	5	1	0
2	B	13	0	6	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	205	0	0	0	0
4	B	220	0	0	2	1
All	All	3739	0	3301	14	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:GLY:OXT	4:B:821:HOH:O	1.79	0.99
1:B:202:GLU:OE1	4:B:820:HOH:O	2.09	0.70
1:B:60:LEU:HD12	1:B:83:TRP:HE3	1.67	0.58
1:A:122:ALA:HA	1:A:125:TRP:CE3	2.40	0.56
1:A:89:CYS:O	1:B:115:GLY:HA2	2.06	0.55
1:B:122:ALA:HA	1:B:125:TRP:CE3	2.44	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:HIS:CE1	2:A:501:5RP:C2	2.92	0.52
1:A:89:CYS:HB2	1:B:115:GLY:HA3	1.92	0.50
1:A:158:ARG:O	1:A:162:MET:HG3	2.14	0.47
1:A:168:VAL:HG12	1:A:172:LEU:HD11	1.98	0.46
1:B:60:LEU:HD11	1:B:110:GLN:NE2	2.30	0.46
1:A:175:GLU:O	1:A:178:PRO:HD2	2.19	0.42
1:A:177:LEU:N	1:A:178:PRO:CD	2.84	0.41
1:B:87:ILE:HD12	1:B:87:ILE:HA	2.00	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:70:LYS:NZ	1:B:104:GLU:OE1[1_545]	2.17	0.03
4:B:688:HOH:O	4:B:812:HOH:O[4_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/216 (99%)	211 (99%)	3 (1%)	0	100	100
1	B	214/216 (99%)	211 (99%)	3 (1%)	0	100	100
All	All	428/432 (99%)	422 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/169 (101%)	168 (99%)	2 (1%)	71	53
1	B	164/169 (97%)	163 (99%)	1 (1%)	86	76
All	All	334/338 (99%)	331 (99%)	3 (1%)	78	66

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	LEU
1	A	112	GLU
1	B	112	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	80	ASN
1	A	110	GLN
1	A	206	GLN
1	B	7	GLN
1	B	13	GLN
1	B	110	GLN
1	B	206	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5RP	A	501	3	12,12,13	1.10	2 (16%)	14,17,18	1.92	3 (21%)
2	5RP	B	502	3	12,12,13	1.43	2 (16%)	14,17,18	2.19	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5RP	A	501	3	-	3/14/14/16	-
2	5RP	B	502	3	-	7/14/14/16	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	502	5RP	O4-C3	3.55	1.30	1.21
2	A	501	5RP	C5-C3	2.42	1.54	1.52
2	B	502	5RP	C5-C3	2.15	1.54	1.52
2	A	501	5RP	O4-C3	2.06	1.26	1.21

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	5RP	O13-C5-C3	-4.74	103.47	111.31
2	B	502	5RP	O4-C3-C5	4.58	125.39	119.24
2	B	502	5RP	C2-C3-C5	-4.13	113.78	118.13
2	A	501	5RP	O4-C3-C5	3.31	123.69	119.24
2	B	502	5RP	O8-C7-C6	3.21	117.92	109.36
2	B	502	5RP	O14-C6-C5	-2.97	105.37	109.67
2	A	501	5RP	O10-P9-O11	2.00	115.31	107.80

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	5RP	O13-C5-C6-O14
2	A	501	5RP	O13-C5-C6-C7
2	B	502	5RP	C3-C5-C6-O14
2	B	502	5RP	O13-C5-C6-O14
2	B	502	5RP	O13-C5-C6-C7
2	B	502	5RP	C5-C6-C7-O8
2	B	502	5RP	O14-C6-C7-O8
2	B	502	5RP	C3-C5-C6-C7
2	A	501	5RP	C6-C7-O8-P9
2	B	502	5RP	C2-C3-C5-O13

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	5RP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/216 (98%)	0.10	12 (5%) 24 23	13, 22, 40, 48	0
1	B	215/216 (99%)	-0.09	6 (2%) 53 53	11, 20, 35, 45	0
All	All	428/432 (99%)	0.00	18 (4%) 36 35	11, 21, 38, 48	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	GLY	5.6
1	A	147	ALA	4.8
1	A	148	TRP	4.5
1	A	145	GLY	4.4
1	B	145	GLY	4.4
1	A	146	VAL	3.7
1	A	141	ALA	3.5
1	A	144	ALA	3.4
1	A	179	LEU	3.1
1	A	143	ALA	3.0
1	A	116	TYR	2.5
1	B	146	VAL	2.5
1	B	114	THR	2.4
1	B	144	ALA	2.3
1	B	216	GLY	2.3
1	B	161	ASP	2.2
1	A	106	ASN	2.2
1	A	175	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	B	602	1/1	0.90	0.07	19,19,19,19	0
3	MG	A	601	1/1	0.96	0.05	20,20,20,20	0
2	5RP	A	501	13/14	0.96	0.09	18,24,34,40	0
2	5RP	B	502	13/14	0.97	0.09	16,27,35,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.