



# wwPDB X-ray Structure Validation Summary Report i

Feb 5, 2024 – 05:29 AM EST

PDB ID : 1XFY  
Title : Crystal structure of anthrax edema factor (EF) in complex with calmodulin  
Authors : Shen, Y.; Zhukovskaya, N.L.; Guo, Q.; Florian, J.; Tang, W.J.  
Deposited on : 2004-09-15  
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)
RSRZ outliers	127900	1115 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

*Continued from previous page...*



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	C	902	-	-	-	X
4	CA	P	803	-	-	-	X

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 42846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calmodulin-sensitive adenylate cyclase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	B	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	C	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	D	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	E	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			
1	F	735	Total	C	N	O	S	0	0	0
			5992	3828	995	1163	6			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	initiating methionine	UNP P40136
A	25	HIS	-	expression tag	UNP P40136
A	26	HIS	-	expression tag	UNP P40136
A	27	HIS	-	expression tag	UNP P40136
A	28	HIS	-	expression tag	UNP P40136
A	29	HIS	-	expression tag	UNP P40136
A	30	HIS	-	expression tag	UNP P40136
A	31	ALA	-	cloning artifact	UNP P40136
A	32	ALA	-	cloning artifact	UNP P40136
B	24	MET	-	initiating methionine	UNP P40136
B	25	HIS	-	expression tag	UNP P40136
B	26	HIS	-	expression tag	UNP P40136
B	27	HIS	-	expression tag	UNP P40136
B	28	HIS	-	expression tag	UNP P40136
B	29	HIS	-	expression tag	UNP P40136
B	30	HIS	-	expression tag	UNP P40136
B	31	ALA	-	cloning artifact	UNP P40136

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	32	ALA	-	cloning artifact	UNP P40136
C	24	MET	-	initiating methionine	UNP P40136
C	25	HIS	-	expression tag	UNP P40136
C	26	HIS	-	expression tag	UNP P40136
C	27	HIS	-	expression tag	UNP P40136
C	28	HIS	-	expression tag	UNP P40136
C	29	HIS	-	expression tag	UNP P40136
C	30	HIS	-	expression tag	UNP P40136
C	31	ALA	-	cloning artifact	UNP P40136
C	32	ALA	-	cloning artifact	UNP P40136
D	24	MET	-	initiating methionine	UNP P40136
D	25	HIS	-	expression tag	UNP P40136
D	26	HIS	-	expression tag	UNP P40136
D	27	HIS	-	expression tag	UNP P40136
D	28	HIS	-	expression tag	UNP P40136
D	29	HIS	-	expression tag	UNP P40136
D	30	HIS	-	expression tag	UNP P40136
D	31	ALA	-	cloning artifact	UNP P40136
D	32	ALA	-	cloning artifact	UNP P40136
E	24	MET	-	initiating methionine	UNP P40136
E	25	HIS	-	expression tag	UNP P40136
E	26	HIS	-	expression tag	UNP P40136
E	27	HIS	-	expression tag	UNP P40136
E	28	HIS	-	expression tag	UNP P40136
E	29	HIS	-	expression tag	UNP P40136
E	30	HIS	-	expression tag	UNP P40136
E	31	ALA	-	cloning artifact	UNP P40136
E	32	ALA	-	cloning artifact	UNP P40136
F	24	MET	-	initiating methionine	UNP P40136
F	25	HIS	-	expression tag	UNP P40136
F	26	HIS	-	expression tag	UNP P40136
F	27	HIS	-	expression tag	UNP P40136
F	28	HIS	-	expression tag	UNP P40136
F	29	HIS	-	expression tag	UNP P40136
F	30	HIS	-	expression tag	UNP P40136
F	31	ALA	-	cloning artifact	UNP P40136
F	32	ALA	-	cloning artifact	UNP P40136

- Molecule 2 is a protein called Calmodulin 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	O	146	Total	C	N	O	S	0	0	0
			1146	702	186	249	9			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	146	Total	C 1146	N 702	O 186	S 249	9	0	0
2	Q	146	Total	C 1146	N 702	O 186	S 249	9	0	0
2	R	146	Total	C 1146	N 702	O 186	S 249	9	0	0
2	S	146	Total	C 1146	N 702	O 186	S 249	9	0	0
2	T	146	Total	C 1146	N 702	O 186	S 249	9	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg 1	0	0
3	B	1	Total	Mg 1	0	0
3	C	1	Total	Mg 1	0	0
3	D	1	Total	Mg 1	0	0
3	E	1	Total	Mg 1	0	0
3	F	1	Total	Mg 1	0	0


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	O	2	Total	Ca 2	0	0
4	P	2	Total	Ca 2	0	0
4	Q	2	Total	Ca 2	0	0
4	R	2	Total	Ca 2	0	0
4	S	2	Total	Ca 2	0	0
4	T	2	Total	Ca 2	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.








- Molecule 1: Calmodulin-sensitive adenylate cyclase












• Molecule 1: Calmodulin-sensitive adenylate cyclase






- Molecule 1: Calmodulin-sensitive adenylate cyclase






### • Molecule 1: Calmodulin-sensitive adenylate cyclase










- Molecule 2: Calmodulin 2



- Molecule 2: Calmodulin 2





- Molecule 2: Calmodulin 2







- Molecule 2: Calmodulin 2



- Molecule 2: Calmodulin 2



- Molecule 2: Calmodulin 2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	315.62 Å    182.04 Å    141.02 Å 90.00°    89.93°    90.00°	Depositor
Resolution (Å)	10.00 – 3.30 29.82 – 3.26	Depositor EDS
% Data completeness (in resolution range)	93.9 (10.00-3.30) 91.8 (29.82-3.26)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	CNS 1.1	Depositor
$R$ , $R_{free}$	0.269 , 0.289 0.273 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	101.4	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 56.2	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.448 for -1/2*h-3/2*k,-1/2*h+1/2*k,-l 0.448 for -1/2*h+3/2*k,1/2*h+1/2*k,-l 0.448 for 1/2*h-3/2*k,-1/2*h-1/2*k,-l 0.448 for 1/2*h+3/2*k,1/2*h-1/2*k,-l 0.458 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	42846	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.54	0/6104	0.84	15/8208 (0.2%)
1	B	0.54	0/6104	0.85	15/8208 (0.2%)
1	C	0.54	0/6104	0.85	17/8208 (0.2%)
1	D	0.54	1/6104 (0.0%)	0.85	17/8208 (0.2%)
1	E	0.54	0/6104	0.85	16/8208 (0.2%)
1	F	0.54	0/6104	0.85	16/8208 (0.2%)
2	O	0.54	0/1158	0.85	4/1553 (0.3%)
2	P	0.54	0/1158	0.86	4/1553 (0.3%)
2	Q	0.54	0/1158	0.85	4/1553 (0.3%)
2	R	0.55	0/1158	0.85	4/1553 (0.3%)
2	S	0.54	0/1158	0.85	4/1553 (0.3%)
2	T	0.53	0/1158	0.85	4/1553 (0.3%)
All	All	0.54	1/43572 (0.0%)	0.85	120/58566 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	190	PRO	N-CA	-5.69	1.37	1.47

The worst 5 of 120 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	160	ALA	N-CA-C	9.55	136.80	111.00
1	F	160	ALA	N-CA-C	9.55	136.78	111.00
1	E	160	ALA	N-CA-C	9.54	136.76	111.00
1	A	160	ALA	N-CA-C	9.54	136.75	111.00
1	D	160	ALA	N-CA-C	9.53	136.74	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5992	0	6010	866	0
1	B	5992	0	6010	875	0
1	C	5992	0	6010	855	0
1	D	5992	0	6010	863	0
1	E	5992	0	6010	863	0
1	F	5992	0	6010	862	0
2	O	1146	0	1071	195	0
2	P	1146	0	1071	195	0
2	Q	1146	0	1071	189	0
2	R	1146	0	1071	196	0
2	S	1146	0	1071	196	0
2	T	1146	0	1071	192	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
4	O	2	0	0	0	0
4	P	2	0	0	0	0
4	Q	2	0	0	0	0
4	R	2	0	0	0	0
4	S	2	0	0	0	0
4	T	2	0	0	0	0
All	All	42846	0	42486	6152	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 72.

The worst 5 of 6152 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:183:SER:O	1:A:187:SER:CB	1.70	1.39
1:D:183:SER:O	1:D:187:SER:CB	1.70	1.38
1:F:183:SER:O	1:F:187:SER:CB	1.70	1.37

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:183:SER:O	1:E:187:SER:CB	1.70	1.36
1:B:183:SER:O	1:B:187:SER:CB	1.70	1.35

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	733/777 (94%)	490 (67%)	170 (23%)	73 (10%)	0 3
1	B	733/777 (94%)	492 (67%)	171 (23%)	70 (10%)	0 4
1	C	733/777 (94%)	494 (67%)	171 (23%)	68 (9%)	0 4
1	D	733/777 (94%)	493 (67%)	170 (23%)	70 (10%)	0 4
1	E	733/777 (94%)	492 (67%)	171 (23%)	70 (10%)	0 4
1	F	733/777 (94%)	490 (67%)	172 (24%)	71 (10%)	0 3
2	O	144/149 (97%)	92 (64%)	39 (27%)	13 (9%)	1 4
2	P	144/149 (97%)	89 (62%)	39 (27%)	16 (11%)	0 2
2	Q	144/149 (97%)	90 (62%)	37 (26%)	17 (12%)	0 2
2	R	144/149 (97%)	90 (62%)	38 (26%)	16 (11%)	0 2
2	S	144/149 (97%)	90 (62%)	39 (27%)	15 (10%)	0 3
2	T	144/149 (97%)	90 (62%)	40 (28%)	14 (10%)	0 3
All	All	5262/5556 (95%)	3492 (66%)	1257 (24%)	513 (10%)	0 3

5 of 513 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	77	ASP
1	A	111	LEU
1	A	113	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	135	VAL
1	A	162	ASN

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	664/705 (94%)	580 (87%)	84 (13%)	4 19
1	B	664/705 (94%)	576 (87%)	88 (13%)	4 17
1	C	664/705 (94%)	578 (87%)	86 (13%)	4 17
1	D	664/705 (94%)	577 (87%)	87 (13%)	4 17
1	E	664/705 (94%)	578 (87%)	86 (13%)	4 17
1	F	664/705 (94%)	579 (87%)	85 (13%)	4 18
2	O	123/127 (97%)	106 (86%)	17 (14%)	3 16
2	P	123/127 (97%)	106 (86%)	17 (14%)	3 16
2	Q	123/127 (97%)	106 (86%)	17 (14%)	3 16
2	R	123/127 (97%)	106 (86%)	17 (14%)	3 16
2	S	123/127 (97%)	106 (86%)	17 (14%)	3 16
2	T	123/127 (97%)	106 (86%)	17 (14%)	3 16
All	All	4722/4992 (95%)	4104 (87%)	618 (13%)	4 17

5 of 618 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	284	LYS
2	R	42	ASN
1	F	408	LEU
1	F	279	ILE
2	O	28	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 191 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	794	GLN
1	E	789	ASN
1	E	368	GLN
1	E	581	GLN
1	F	387	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 18 ligands modelled in this entry, 18 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	735/777 (94%)	3.03	439 (59%) 0 0	24, 83, 143, 153	0
1	B	735/777 (94%)	2.95	441 (60%) 0 0	25, 83, 143, 153	0
1	C	735/777 (94%)	3.07	483 (65%) 0 0	25, 83, 143, 153	0
1	D	735/777 (94%)	2.93	458 (62%) 0 0	25, 83, 143, 153	0
1	E	735/777 (94%)	2.94	453 (61%) 0 0	24, 83, 142, 152	0
1	F	735/777 (94%)	2.93	447 (60%) 0 0	27, 83, 143, 154	0
2	O	146/149 (97%)	2.80	85 (58%) 0 0	22, 74, 121, 134	0
2	P	146/149 (97%)	2.48	78 (53%) 0 0	20, 75, 121, 135	0
2	Q	146/149 (97%)	2.49	85 (58%) 0 0	21, 75, 121, 134	0
2	R	146/149 (97%)	2.58	75 (51%) 0 0	20, 75, 120, 134	0
2	S	146/149 (97%)	2.67	87 (59%) 0 0	21, 76, 121, 134	0
2	T	146/149 (97%)	2.81	80 (54%) 0 0	21, 75, 120, 134	0
All	All	5286/5556 (95%)	2.92	3211 (60%) 0 0	20, 80, 142, 154	0

The worst 5 of 3211 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	468	LYS	21.6
1	D	204	ASP	21.2
2	P	78	ASP	20.1
1	D	211	SER	19.8
1	E	593	ILE	19.8

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	E	904	1/1	-0.20	0.34	23,23,23,23	0
4	CA	O	802	1/1	0.08	0.13	33,33,33,33	0
4	CA	S	809	1/1	0.29	0.20	29,29,29,29	0
4	CA	R	808	1/1	0.30	0.21	36,36,36,36	0
3	MG	C	902	1/1	0.33	0.57	23,23,23,23	0
4	CA	R	807	1/1	0.35	0.13	30,30,30,30	0
3	MG	D	903	1/1	0.39	0.31	21,21,21,21	0
3	MG	B	901	1/1	0.40	0.29	16,16,16,16	0
4	CA	T	812	1/1	0.40	0.25	35,35,35,35	0
4	CA	Q	806	1/1	0.41	0.16	35,35,35,35	0
4	CA	P	803	1/1	0.49	0.48	32,32,32,32	0
3	MG	A	900	1/1	0.53	0.28	23,23,23,23	0
4	CA	S	810	1/1	0.55	0.28	40,40,40,40	0
4	CA	O	801	1/1	0.60	0.13	33,33,33,33	0
4	CA	Q	805	1/1	0.65	0.21	29,29,29,29	0
4	CA	P	804	1/1	0.67	0.28	34,34,34,34	0
4	CA	T	811	1/1	0.79	0.23	28,28,28,28	0
3	MG	F	905	1/1	0.92	0.29	17,17,17,17	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.