



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 5, 2023 – 10:32 pm GMT

PDB ID : 2CGO
Title : FACTOR INHIBITING HIF-1 ALPHA with fumarate
Authors : McDonough, M.A.; Clifton, I.J.; Schofield, C.J.
Deposited on : 2006-03-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	349	 75% 16% 6% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HYPOXIA-INDUCIBLE FACTOR 1 ALPHA INHIBITOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	339	2777	1779	473	514	11	0	0	0

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		

- Molecule 3 is FUMARIC ACID (three-letter code: FUM) (formula: C₄H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	4	4		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	1
			10	8	2		
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	121	Total	O	0	0
			121	121		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HYPOXIA-INDUCIBLE FACTOR 1 ALPHA INHIBITOR

Chain A:  75% 16% 6%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.66Å 86.66Å 147.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.74 – 2.30 37.48 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (74.74-2.30) 99.5 (37.48-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.237 0.246 , 0.282	Depositor DCC
R_{free} test set	2532 reflections (9.89%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtrriage
Anisotropy	0.325	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2932	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FE, FUM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.28	11/2857 (0.4%)	1.07	13/3877 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	225	GLU	CG-CD	8.28	1.64	1.51
1	A	29	GLU	CG-CD	7.20	1.62	1.51
1	A	225	GLU	CD-OE2	6.84	1.33	1.25
1	A	70	VAL	CB-CG1	6.82	1.67	1.52
1	A	40	ARG	CG-CD	6.29	1.67	1.51
1	A	71	TYR	CZ-OH	6.04	1.48	1.37
1	A	238	ARG	CZ-NH1	5.94	1.40	1.33
1	A	335	GLU	CD-OE1	5.80	1.32	1.25
1	A	59	GLU	CG-CD	-5.22	1.44	1.51
1	A	176	LYS	CE-NZ	5.15	1.61	1.49
1	A	222	ASP	C-O	-5.13	1.13	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	ASP	CB-CG-OD2	11.89	129.00	118.30
1	A	201	ASP	CB-CG-OD1	-10.09	109.22	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177	ARG	NE-CZ-NH2	-9.26	115.67	120.30
1	A	177	ARG	NE-CZ-NH1	7.90	124.25	120.30
1	A	238	ARG	NE-CZ-NH2	-7.84	116.38	120.30
1	A	326	LEU	CB-CG-CD2	-7.07	98.98	111.00
1	A	237	ASP	CB-CG-OD1	6.17	123.86	118.30
1	A	251	ARG	NE-CZ-NH1	6.01	123.31	120.30
1	A	269	ASP	CB-CG-OD2	-5.72	113.16	118.30
1	A	243	ASP	CB-CG-OD2	-5.57	113.29	118.30
1	A	238	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	A	40	ARG	N-CA-CB	-5.44	100.80	110.60
1	A	17	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	304	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2777	0	2660	45	0
2	A	1	0	0	0	0
3	A	8	0	1	0	0
4	A	25	0	0	0	0
5	A	121	0	0	5	0
All	All	2932	0	2661	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:ASN:HA	5:A:2045:HOH:O	1.82	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:LYS:NZ	5:A:2057:HOH:O	2.15	0.77
1:A:334:GLN:HE21	1:A:334:GLN:HA	1.53	0.71
1:A:309:PRO:O	1:A:310:LEU:HB2	1.91	0.70
1:A:85:ILE:CD1	1:A:85:ILE:O	2.43	0.66
1:A:112:GLN:O	1:A:114:PHE:N	2.31	0.64
1:A:9:VAL:HG12	1:A:40:ARG:HD2	1.80	0.63
1:A:133:GLN:HE21	1:A:133:GLN:HA	1.64	0.63
1:A:132:LEU:HD23	1:A:144:LEU:HD12	1.80	0.61
1:A:306:ILE:HD13	1:A:306:ILE:N	2.16	0.60
1:A:322:ILE:HA	1:A:325:MET:HE2	1.86	0.58
1:A:85:ILE:O	1:A:85:ILE:HD13	2.05	0.56
1:A:305:ARG:HB3	1:A:306:ILE:HD13	1.88	0.55
1:A:53:GLU:HG3	1:A:177:ARG:HH22	1.72	0.54
1:A:219:PHE:HB2	1:A:278:TRP:HB2	1.89	0.53
1:A:334:GLN:HE21	1:A:334:GLN:CA	2.20	0.53
1:A:87:ASN:N	1:A:87:ASN:OD1	2.43	0.52
1:A:85:ILE:O	1:A:85:ILE:HD12	2.08	0.52
1:A:223:GLN:NE2	1:A:258:VAL:HG22	2.25	0.51
1:A:346:GLY:H	1:A:349:ASN:HD21	1.59	0.51
1:A:223:GLN:HE21	1:A:258:VAL:HG23	1.77	0.50
1:A:115:LYS:N	5:A:2044:HOH:O	2.38	0.49
1:A:223:GLN:NE2	1:A:258:VAL:CG2	2.76	0.49
1:A:75:LYS:NZ	1:A:161:ASP:OD1	2.46	0.48
1:A:330:LEU:HD13	1:A:335:GLU:HB3	1.96	0.47
1:A:326:LEU:HD11	1:A:339:LEU:HD22	1.97	0.47
1:A:10:ALA:HA	1:A:39:THR:O	2.15	0.47
1:A:115:LYS:HB2	5:A:2044:HOH:O	2.14	0.47
1:A:306:ILE:HD13	1:A:306:ILE:H	1.80	0.47
1:A:134:ASP:OD2	1:A:138:ARG:NH1	2.48	0.46
1:A:135:ILE:HD13	1:A:142:GLU:O	2.16	0.46
1:A:172:LYS:HG2	1:A:176:LYS:NZ	2.29	0.46
1:A:82:GLN:NE2	1:A:126:HIS:HB2	2.31	0.46
1:A:185:ASN:HA	1:A:294:ASN:O	2.16	0.45
1:A:275:MET:HE1	5:A:2027:HOH:O	2.16	0.45
1:A:108:MET:HG2	1:A:114:PHE:CE2	2.52	0.44
1:A:311:LYS:HD3	1:A:313:HIS:CE1	2.53	0.44
1:A:232:VAL:HG11	1:A:324:LYS:HG3	2.00	0.43
1:A:223:GLN:HE21	1:A:258:VAL:CG2	2.32	0.43
1:A:227:LEU:HD13	1:A:278:TRP:CG	2.54	0.42
1:A:103:TYR:CD2	1:A:114:PHE:CZ	3.09	0.41
1:A:223:GLN:HE22	1:A:258:VAL:HG22	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:TYR:CD2	1:A:114:PHE:HZ	2.39	0.41
1:A:222:ASP:OD1	1:A:222:ASP:C	2.60	0.40
1:A:349:ASN:HD22	1:A:349:ASN:C	2.24	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/349 (96%)	311 (93%)	20 (6%)	4 (1%)	13 14

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	ASN
1	A	304	LYS
1	A	310	LEU
1	A	309	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	297/304 (98%)	273 (92%)	24 (8%)	11 15

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLU
1	A	29	GLU
1	A	40	ARG
1	A	47	GLN
1	A	53	GLU
1	A	85	ILE
1	A	87	ASN
1	A	99	LYS
1	A	106	LYS
1	A	133	GLN
1	A	138	ARG
1	A	145	TYR
1	A	149	THR
1	A	156	ARG
1	A	172	LYS
1	A	181	GLN
1	A	205	ASN
1	A	258	VAL
1	A	305	ARG
1	A	306	ILE
1	A	307	GLU
1	A	334	GLN
1	A	339	LEU
1	A	349	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	GLN
1	A	82	GLN
1	A	133	GLN
1	A	137	GLN
1	A	205	ASN
1	A	223	GLN
1	A	313	HIS
1	A	321	ASN
1	A	334	GLN
1	A	349	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	A	1353	-	4,4,4	0.11	0	6,6,6	0.37	0
3	FUM	A	351	2	7,7,7	1.26	1 (14%)	8,8,8	1.00	0
4	SO4	A	1354[B]	-	4,4,4	0.03	0	6,6,6	0.86	0
4	SO4	A	1354[A]	-	4,4,4	0.11	0	6,6,6	1.37	1 (16%)
4	SO4	A	1352	-	4,4,4	0.15	0	6,6,6	0.23	0
4	SO4	A	1355	-	4,4,4	0.25	0	6,6,6	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FUM	A	351	2	-	0/5/5/5	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	351	FUM	O7-C6	2.18	1.28	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	1354[A]	SO4	O4-S-O3	-2.42	98.74	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.