



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2024 – 01:37 AM EST

PDB ID : 2P2B  
Title : Acetyl-CoA Synthetase, V386A mutation  
Authors : Reger, A.S.; Gulick, A.M.  
Deposited on : 2007-03-07  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	 92% 5% •
1	B	652	 % 90% 8% ••

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetyl-coenzyme A synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	633	4899	3114	845	923	17	0	0	0
1	B	639	4906	3122	846	920	18	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	386	ALA	VAL	engineered mutation	UNP Q8ZKF6
B	386	ALA	VAL	engineered mutation	UNP Q8ZKF6

- Molecule 2 is COENZYME A (three-letter code: COA) (formula:  $C_{21}H_{36}N_7O_{16}P_3S$ ).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
2	A	1	48	21	7	16	3	1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	B	1	48	21	7	16	3	1	0	0

- Molecule 3 is ADENOSINE-5'-MONOPHOSPHATE-PROPYL ESTER (three-letter code: PRX) (formula: C<sub>13</sub>H<sub>20</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	26	13	5	7	1	0	0
3	B	1	26	13	5	7	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	180	180	180	0	0
4	B	130	130	130	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetyl-coenzyme A synthetase



- Molecule 1: Acetyl-coenzyme A synthetase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.91Å 143.08Å 71.38Å 90.00° 90.97° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.95 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.9 (30.00-2.20) 96.9 (29.95-2.20)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.43 (at 2.20Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.166 , 0.211 0.167 , 0.214	Depositor DCC
$R_{free}$ test set	2980 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtrriage
Anisotropy	0.340	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.075 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10263	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, PRX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/5031	0.64	0/6864
1	B	0.57	0/5037	0.64	0/6876
All	All	0.59	0/10068	0.64	0/13740

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4899	0	4709	24	0
1	B	4906	0	4717	29	0
2	A	48	0	32	3	0
2	B	48	0	32	1	0
3	A	26	0	19	1	0
3	B	26	0	19	0	0
4	A	180	0	0	2	0
4	B	130	0	0	0	0
All	All	10263	0	9528	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:628:ASP:O	1:B:631:THR:HG22	1.77	0.84
1:B:154:ARG:HE	1:B:248:HIS:HD2	1.29	0.81
1:B:154:ARG:NE	1:B:248:HIS:HD2	1.95	0.65
1:B:552:HIS:HD2	1:B:555:LYS:H	1.46	0.64
1:B:552:HIS:CD2	1:B:555:LYS:H	2.16	0.64
1:B:154:ARG:HE	1:B:248:HIS:CD2	2.15	0.62
1:B:529:THR:HG22	1:B:612:ARG:HH21	1.69	0.58
1:A:552:HIS:HD2	1:A:555:LYS:H	1.52	0.58
1:A:387:GLY:HA2	3:A:999:PRX:H3P3	1.87	0.56
1:B:76:ASN:HD22	1:B:79:ALA:H	1.53	0.55
1:A:76:ASN:HD22	1:A:79:ALA:H	1.55	0.55
1:A:335:ASN:H	1:A:335:ASN:HD22	1.54	0.54
1:A:335:ASN:ND2	2:A:990:COA:O9P	2.40	0.54
1:B:552:HIS:CD2	1:B:554:ILE:H	2.26	0.54
1:A:529:THR:HG21	4:A:1108:HOH:O	2.09	0.53
1:B:353:ILE:HG23	1:B:382:ILE:HB	1.91	0.52
1:A:552:HIS:CD2	1:A:555:LYS:H	2.30	0.50
1:B:552:HIS:HD2	1:B:554:ILE:H	1.57	0.50
1:B:525:HIS:CE1	1:B:590:LEU:HD22	2.46	0.50
1:B:601:SER:HB3	1:B:637:VAL:HB	1.93	0.49
1:B:299:ASP:OD1	1:B:381:ARG:NH1	2.44	0.48
1:A:335:ASN:H	1:A:335:ASN:ND2	2.11	0.47
1:B:335:ASN:H	1:B:335:ASN:HD22	1.62	0.47
1:A:610:ILE:HD11	1:A:615:LEU:CD1	2.44	0.47
1:A:610:ILE:HD11	1:A:615:LEU:HD11	1.97	0.47
1:B:20:ASN:H	1:B:23:GLN:NE2	2.13	0.47
1:A:301:TYR:HE1	1:A:315:TYR:O	1.97	0.47
1:A:638:VAL:O	1:A:642:LEU:HG	2.14	0.46
1:B:521:ASN:O	1:B:557:GLN:HB2	2.16	0.46
1:A:333:VAL:HB	2:A:990:COA:O9P	2.15	0.46
1:A:196:ILE:HG21	2:A:990:COA:H2A	1.97	0.45
1:B:196:ILE:HG21	2:B:991:COA:H2A	1.98	0.45
1:A:358:PRO:HD2	1:A:388:GLU:HG2	1.98	0.45
1:B:134:VAL:HG11	1:B:178:SER:HB2	1.99	0.44
1:A:386:ALA:HB2	1:A:412:THR:OG1	2.18	0.44
1:A:530:ALA:HB3	4:A:1063:HOH:O	2.18	0.43
1:B:206:LEU:HD22	1:B:211:VAL:HG11	2.00	0.43
1:A:315:TYR:CE1	1:A:421:PHE:CE2	3.07	0.43
1:B:397:TRP:NE1	1:B:401:LYS:HG3	2.34	0.43
1:A:610:ILE:HD13	1:A:611:MET:N	2.33	0.42

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:ILE:HA	1:B:23:GLN:HE21	1.84	0.42
1:B:528:GLY:O	1:B:531:GLU:HG2	2.20	0.42
1:B:397:TRP:CE2	1:B:401:LYS:HG3	2.54	0.42
1:A:335:ASN:HD22	1:A:335:ASN:N	2.13	0.42
1:A:312:GLY:O	1:A:316:LEU:HB3	2.20	0.41
1:A:109:SER:OG	1:A:112:GLU:HG3	2.20	0.41
1:A:576:TYR:CE1	1:A:596:LEU:HB2	2.55	0.41
1:B:301:TYR:HE1	1:B:315:TYR:O	2.03	0.41
1:A:335:ASN:ND2	1:A:335:ASN:N	2.68	0.41
1:B:597:HIS:CD2	1:B:597:HIS:C	2.93	0.41
1:B:386:ALA:HB2	1:B:412:THR:OG1	2.21	0.41
1:B:32:ILE:HD12	1:B:445:GLN:HG2	2.02	0.41
1:B:61:ALA:HA	1:B:494:ASN:OD1	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	629/652 (96%)	608 (97%)	20 (3%)	1 (0%)	47 55
1	B	635/652 (97%)	618 (97%)	15 (2%)	2 (0%)	41 46
All	All	1264/1304 (97%)	1226 (97%)	35 (3%)	3 (0%)	47 55

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	438	THR
1	B	438	THR
1	B	63	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	506/537 (94%)	497 (98%)	9 (2%)	59 72
1	B	503/537 (94%)	490 (97%)	13 (3%)	46 58
All	All	1009/1074 (94%)	987 (98%)	22 (2%)	52 65

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ILE
1	A	53	GLN
1	A	83	ASP
1	A	88	GLU
1	A	262	LEU
1	A	335	ASN
1	A	483	ARG
1	A	610	ILE
1	A	614	ILE
1	B	83	ASP
1	B	106	LYS
1	B	127	LEU
1	B	250	PRO
1	B	262	LEU
1	B	316	LEU
1	B	335	ASN
1	B	461	GLU
1	B	493	LYS
1	B	521	ASN
1	B	529	THR
1	B	612	ARG
1	B	614	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	53	GLN
1	A	76	ASN
1	A	104	GLN
1	A	335	ASN
1	A	344	GLN
1	A	445	GLN
1	A	552	HIS
1	A	567	HIS
1	B	23	GLN
1	B	30	GLN
1	B	64	ASN
1	B	76	ASN
1	B	248	HIS
1	B	254	ASN
1	B	335	ASN
1	B	344	GLN
1	B	350	GLN
1	B	521	ASN
1	B	552	HIS
1	B	567	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	COA	A	990	-	41,50,50	1.66	3 (7%)	52,75,75	1.26	4 (7%)
3	PRX	A	999	-	25,28,28	0.87	1 (4%)	27,41,41	1.15	2 (7%)
3	PRX	B	998	-	25,28,28	0.88	1 (4%)	27,41,41	1.17	1 (3%)
2	COA	B	991	-	41,50,50	1.66	2 (4%)	52,75,75	1.22	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	990	-	-	12/44/64/64	0/3/3/3
3	PRX	A	999	-	-	0/11/31/31	0/3/3/3
3	PRX	B	998	-	-	1/11/31/31	0/3/3/3
2	COA	B	991	-	-	17/44/64/64	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	991	COA	O9P-C9P	9.05	1.41	1.23
2	A	990	COA	O9P-C9P	8.72	1.40	1.23
2	A	990	COA	C2A-N3A	4.11	1.38	1.32
2	B	991	COA	C2A-N3A	3.65	1.38	1.32
3	A	999	PRX	O4'-C1'	2.39	1.44	1.41
2	A	990	COA	C2A-N1A	2.19	1.38	1.33
3	B	998	PRX	O4'-C1'	2.16	1.44	1.41

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	990	COA	N3A-C2A-N1A	-5.73	119.72	128.68
2	B	991	COA	N3A-C2A-N1A	-5.70	119.77	128.68
3	A	999	PRX	N3-C2-N1	-4.43	121.76	128.68
3	B	998	PRX	N3-C2-N1	-4.33	121.91	128.68
2	A	990	COA	CEP-CBP-CCP	3.49	113.92	108.23
2	B	991	COA	CDP-CBP-CAP	3.10	114.20	108.82

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	991	COA	P2A-O3A-P1A	-2.32	124.87	132.83
2	A	990	COA	O6A-CCP-CBP	2.24	114.15	110.55
2	A	990	COA	C4A-C5A-N7A	-2.12	107.19	109.40
3	A	999	PRX	C4-C5-N7	-2.03	107.28	109.40

There are no chirality outliers.

All (30) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	990	COA	C3B-O3B-P3B-O7A
2	A	990	COA	CDP-CBP-CCP-O6A
2	A	990	COA	CAP-CBP-CCP-O6A
2	A	990	COA	N8P-C9P-CAP-OAP
2	A	990	COA	CAP-C9P-N8P-C7P
2	A	990	COA	O9P-C9P-N8P-C7P
2	B	991	COA	CCP-O6A-P2A-O4A
2	B	991	COA	OAP-CAP-CBP-CCP
2	B	991	COA	C9P-CAP-CBP-CCP
2	B	991	COA	OAP-CAP-CBP-CDP
2	B	991	COA	C9P-CAP-CBP-CDP
2	B	991	COA	OAP-CAP-CBP-CEP
2	B	991	COA	C9P-CAP-CBP-CEP
2	B	991	COA	CAP-C9P-N8P-C7P
2	B	991	COA	O9P-C9P-N8P-C7P
2	B	991	COA	C6P-C5P-N4P-C3P
2	B	991	COA	O5P-C5P-N4P-C3P
2	A	990	COA	CEP-CBP-CCP-O6A
2	B	991	COA	C5P-C6P-C7P-N8P
2	A	990	COA	O9P-C9P-CAP-OAP
2	A	990	COA	O5P-C5P-C6P-C7P
2	A	990	COA	P1A-O3A-P2A-O6A
2	A	990	COA	N4P-C5P-C6P-C7P
2	B	991	COA	O9P-C9P-CAP-CBP
2	B	991	COA	N8P-C9P-CAP-CBP
2	A	990	COA	CCP-O6A-P2A-O3A
2	B	991	COA	C3B-O3B-P3B-O8A
2	B	991	COA	CCP-O6A-P2A-O3A
2	B	991	COA	P2A-O3A-P1A-O2A
3	B	998	PRX	C5'-O5'-P-O1P



There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	990	COA	3	0
3	A	999	PRX	1	0
2	B	991	COA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	633/652 (97%)	-0.55	1 (0%) 95 94	15, 22, 35, 47	0
1	B	639/652 (98%)	-0.35	8 (1%) 77 75	17, 26, 37, 52	0
All	All	1272/1304 (97%)	-0.45	9 (0%) 87 86	15, 24, 37, 52	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	568	GLY	2.7
1	B	63	GLY	2.5
1	B	11	ALA	2.5
1	B	621	GLY	2.5
1	B	19	ILE	2.5
1	B	646	GLN	2.2
1	B	635	PRO	2.1
1	B	630	SER	2.1
1	B	33	ASN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	COA	B	991	48/48	0.94	0.15	28,45,70,70	0
2	COA	A	990	48/48	0.95	0.13	24,43,69,70	0
3	PRX	A	999	26/26	0.98	0.07	15,18,20,21	0
3	PRX	B	998	26/26	0.98	0.08	18,20,23,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around COA A 990:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around PRX A 999:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.