



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 15, 2024 – 05:58 PM EDT

PDB ID : 2QJH
Title : M. jannaschii ADH synthase covalently bound to dihydroxyacetone phosphate
Authors : Ealick, S.E.; Morar, M.
Deposited on : 2007-07-07
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	 2% 73% 21% . .
1	B	273	 2% 68% 26% . .
1	C	273	 % 73% 22% . .
1	D	273	 71% 24% . .
1	E	273	 73% 21% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	273	 68% 26% . .
1	G	273	 68% 26% . .
1	H	273	 68% 26% . .
1	I	273	 74% 20% . .
1	J	273	 72% 22% . .
1	K	273	 73% 21% . .
1	L	273	 71% 23% . .
1	M	273	 72% 23% . .
1	N	273	 72% 21% . .
1	O	273	 69% 25% . .
1	P	273	 71% 22% . .
1	Q	273	 70% 23% . .
1	R	273	 68% 26% . .
1	S	273	 69% 24% . .
1	T	273	 68% 27% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 40004 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative aldolase MJ0400.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	Total 1960	C 1231	N 347	O 371	S 11	0	0	0
1	B	264	Total 1964	C 1233	N 347	O 373	S 11	0	0	0
1	C	265	Total 1984	C 1246	N 350	O 376	S 12	0	0	0
1	D	266	Total 1995	C 1251	N 353	O 379	S 12	0	0	0
1	E	264	Total 1972	C 1238	N 348	O 375	S 11	0	0	0
1	F	264	Total 1968	C 1235	N 347	O 375	S 11	0	0	0
1	G	264	Total 1960	C 1231	N 347	O 371	S 11	0	0	0
1	H	264	Total 1964	C 1234	N 348	O 371	S 11	0	0	0
1	I	264	Total 1982	C 1244	N 352	O 375	S 11	0	0	0
1	J	264	Total 1968	C 1235	N 347	O 375	S 11	0	0	0
1	K	264	Total 1970	C 1237	N 351	O 371	S 11	0	0	0
1	L	265	Total 1979	C 1242	N 352	O 374	S 11	0	0	0
1	M	264	Total 1968	C 1237	N 349	O 371	S 11	0	0	0
1	N	264	Total 1964	C 1233	N 347	O 373	S 11	0	0	0
1	O	264	Total 1968	C 1236	N 348	O 373	S 11	0	0	0
1	P	264	Total 1966	C 1234	N 350	O 371	S 11	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Q	264	Total	C	N	O	S	0	0	0
			1972	1239	349	373	11			
1	R	266	Total	C	N	O	S	0	0	0
			1986	1246	350	379	11			
1	S	264	Total	C	N	O	S	0	0	0
			1961	1232	346	372	11			
1	T	264	Total	C	N	O	S	0	0	0
			1960	1231	347	371	11			

- Molecule 2 is 1,3-DIHYDROXYACETONEPHOSPHATE (three-letter code: 13P) (formula: C₃H₇O₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			9	3	5	1		
2	B	1	Total	C	O	P	0	0
			9	3	5	1		
2	C	1	Total	C	O	P	0	0
			9	3	5	1		
2	D	1	Total	C	O	P	0	0
			9	3	5	1		
2	E	1	Total	C	O	P	0	0
			9	3	5	1		
2	F	1	Total	C	O	P	0	0
			9	3	5	1		
2	G	1	Total	C	O	P	0	0
			9	3	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	H	1	Total 9	C 3	O 5	P 1	0	0
2	I	1	Total 9	C 3	O 5	P 1	0	0
2	J	1	Total 9	C 3	O 5	P 1	0	0
2	K	1	Total 9	C 3	O 5	P 1	0	0
2	L	1	Total 9	C 3	O 5	P 1	0	0
2	M	1	Total 9	C 3	O 5	P 1	0	0
2	N	1	Total 9	C 3	O 5	P 1	0	0
2	O	1	Total 9	C 3	O 5	P 1	0	0
2	P	1	Total 9	C 3	O 5	P 1	0	0
2	Q	1	Total 9	C 3	O 5	P 1	0	0
2	R	1	Total 9	C 3	O 5	P 1	0	0
2	S	1	Total 9	C 3	O 5	P 1	0	0
2	T	1	Total 9	C 3	O 5	P 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total 13	O 13	0	0
3	B	15	Total 15	O 15	0	0
3	C	20	Total 20	O 20	0	0
3	D	23	Total 23	O 23	0	0
3	E	23	Total 23	O 23	0	0
3	F	25	Total 25	O 25	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	G	17	Total 17	O 17	0	0
3	H	13	Total 13	O 13	0	0
3	I	24	Total 24	O 24	0	0
3	J	19	Total 19	O 19	0	0
3	K	27	Total 27	O 27	0	0
3	L	26	Total 26	O 26	0	0
3	M	23	Total 23	O 23	0	0
3	N	20	Total 20	O 20	0	0
3	O	27	Total 27	O 27	0	0
3	P	17	Total 17	O 17	0	0
3	Q	23	Total 23	O 23	0	0
3	R	24	Total 24	O 24	0	0
3	S	20	Total 20	O 20	0	0
3	T	14	Total 14	O 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative aldolase MJ0400



- Molecule 1: Putative aldolase MJ0400



- Molecule 1: Putative aldolase MJ0400







- Molecule 1: Putative aldolase MJ0400

Chain D: 71% 24%





- Molecule 1: Putative aldolase MJ0400

Chain E: 73% 21%




- Molecule 1: Putative aldolase MJ0400


Chain F: 68% 26%




- Molecule 1: Putative aldolase MJ0400

Chain G: 68% 26%







● Molecule 1: Putative aldolase MJ0400



● Molecule 1: Putative aldolase MJ0400




● Molecule 1: Putative aldolase MJ0400




● Molecule 1: Putative aldolase MJ0400







- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400




- Molecule 1: Putative aldolase MJ0400




• Molecule 1: Putative aldolase MJ0400



• Molecule 1: Putative aldolase MJ0400




• Molecule 1: Putative aldolase MJ0400






• Molecule 1: Putative aldolase MJ0400



• Molecule 1: Putative aldolase MJ0400



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	94.49Å 101.86Å 154.16Å 90.31° 86.70° 82.44°	Depositor
Resolution (Å)	48.28 – 2.60 54.20 – 2.35	Depositor EDS
% Data completeness (in resolution range)	84.2 (48.28-2.60) 84.3 (54.20-2.35)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.00 (at 2.34Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.203 , 0.244 0.198 , 0.236	Depositor DCC
R_{free} test set	16785 reflections (9.94%)	wwPDB-VP
Wilson B-factor (Å ²)	51.0	Xtrriage
Anisotropy	0.382	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	40004	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 13P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1990	0.63	2/2699 (0.1%)
1	B	0.38	0/1994	0.62	0/2704
1	C	0.39	0/2014	0.64	0/2727
1	D	0.40	0/2025	0.64	0/2742
1	E	0.61	6/2002 (0.3%)	0.76	3/2713 (0.1%)
1	F	0.36	0/1998	0.61	0/2709
1	G	0.37	0/1990	0.63	0/2699
1	H	0.38	0/1994	0.61	0/2703
1	I	0.42	0/2012	0.64	0/2724
1	J	0.40	0/1998	0.64	1/2709 (0.0%)
1	K	0.40	0/2000	0.64	2/2710 (0.1%)
1	L	0.40	0/2009	1.10	6/2722 (0.2%)
1	M	0.36	0/1998	0.64	2/2707 (0.1%)
1	N	0.37	0/1994	0.63	2/2704 (0.1%)
1	O	0.39	0/1998	0.62	0/2708
1	P	0.40	0/1996	0.64	2/2706 (0.1%)
1	Q	0.40	1/2002 (0.0%)	0.66	2/2712 (0.1%)
1	R	0.38	0/2016	0.65	2/2732 (0.1%)
1	S	0.38	0/1991	0.67	2/2700 (0.1%)
1	T	0.36	0/1990	0.62	1/2699 (0.0%)
All	All	0.40	7/40011 (0.0%)	0.67	27/54229 (0.0%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	251	ARG	CZ-NH1	-12.47	1.16	1.33
1	E	251	ARG	CZ-NH2	-8.12	1.22	1.33
1	E	251	ARG	N-CA	-8.02	1.30	1.46
1	E	251	ARG	NE-CZ	-6.82	1.24	1.33
1	Q	141	CYS	CB-SG	-5.84	1.72	1.81

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	173	ARG	NE-CZ-NH1	-31.62	104.49	120.30
1	L	173	ARG	NE-CZ-NH2	29.11	134.85	120.30
1	L	173	ARG	CD-NE-CZ	15.64	145.49	123.60
1	E	251	ARG	NE-CZ-NH1	-13.58	113.51	120.30
1	E	251	ARG	NE-CZ-NH2	13.06	126.83	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1960	0	1945	56	0
1	B	1964	0	1948	56	0
1	C	1984	0	1986	55	0
1	D	1995	0	1992	55	0
1	E	1972	0	1963	56	0
1	F	1968	0	1952	68	0
1	G	1960	0	1944	64	0
1	H	1964	0	1955	66	0
1	I	1982	0	1985	50	0
1	J	1968	0	1953	60	0
1	K	1970	0	1967	58	0
1	L	1979	0	1973	61	0
1	M	1968	0	1967	55	0
1	N	1964	0	1949	59	0
1	O	1968	0	1959	65	0
1	P	1966	0	1956	64	0
1	Q	1972	0	1971	59	0
1	R	1986	0	1975	80	0
1	S	1961	0	1944	66	0
1	T	1960	0	1944	60	0
2	A	9	0	5	2	0
2	B	9	0	5	2	0
2	C	9	0	5	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	9	0	5	2	0
2	E	9	0	5	2	0
2	F	9	0	5	2	0
2	G	9	0	5	2	0
2	H	9	0	5	2	0
2	I	9	0	5	2	0
2	J	9	0	5	2	0
2	K	9	0	5	2	0
2	L	9	0	5	2	0
2	M	9	0	5	2	0
2	N	9	0	5	2	0
2	O	9	0	5	2	0
2	P	9	0	5	2	0
2	Q	9	0	5	2	0
2	R	9	0	5	2	0
2	S	9	0	5	2	0
2	T	9	0	5	2	0
3	A	13	0	0	3	0
3	B	15	0	0	1	0
3	C	20	0	0	1	0
3	D	23	0	0	3	0
3	E	23	0	0	7	0
3	F	25	0	0	5	0
3	G	17	0	0	8	0
3	H	13	0	0	2	0
3	I	24	0	0	2	0
3	J	19	0	0	4	0
3	K	27	0	0	4	0
3	L	26	0	0	6	0
3	M	23	0	0	3	0
3	N	20	0	0	6	0
3	O	27	0	0	4	0
3	P	17	0	0	4	0
3	Q	23	0	0	2	0
3	R	24	0	0	7	0
3	S	20	0	0	4	0
3	T	14	0	0	0	0
All	All	40004	0	39328	1149	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1149 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Q:136:MET:HE3	1:Q:136:MET:HA	1.44	0.98
1:L:136:MET:HE3	1:L:136:MET:HA	1.45	0.98
1:D:136:MET:HE3	1:D:136:MET:HA	1.46	0.98
1:E:136:MET:HE3	1:E:136:MET:HA	1.47	0.97
1:H:136:MET:HA	1:H:136:MET:HE3	1.48	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/273 (95%)	247 (95%)	13 (5%)	0	100	100
1	B	260/273 (95%)	249 (96%)	10 (4%)	1 (0%)	34	57
1	C	261/273 (96%)	247 (95%)	14 (5%)	0	100	100
1	D	262/273 (96%)	252 (96%)	10 (4%)	0	100	100
1	E	260/273 (95%)	249 (96%)	11 (4%)	0	100	100
1	F	260/273 (95%)	249 (96%)	8 (3%)	3 (1%)	13	27
1	G	260/273 (95%)	244 (94%)	16 (6%)	0	100	100
1	H	260/273 (95%)	247 (95%)	12 (5%)	1 (0%)	34	57
1	I	260/273 (95%)	247 (95%)	11 (4%)	2 (1%)	19	39
1	J	260/273 (95%)	247 (95%)	12 (5%)	1 (0%)	34	57
1	K	260/273 (95%)	245 (94%)	15 (6%)	0	100	100
1	L	261/273 (96%)	249 (95%)	12 (5%)	0	100	100
1	M	260/273 (95%)	243 (94%)	16 (6%)	1 (0%)	34	57
1	N	260/273 (95%)	243 (94%)	16 (6%)	1 (0%)	34	57
1	O	260/273 (95%)	250 (96%)	10 (4%)	0	100	100
1	P	260/273 (95%)	248 (95%)	12 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	260/273 (95%)	245 (94%)	14 (5%)	1 (0%)	34	57
1	R	262/273 (96%)	250 (95%)	12 (5%)	0	100	100
1	S	260/273 (95%)	248 (95%)	12 (5%)	0	100	100
1	T	260/273 (95%)	247 (95%)	13 (5%)	0	100	100
All	All	5206/5460 (95%)	4946 (95%)	249 (5%)	11 (0%)	47	71

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	267	LEU
1	J	268	LYS
1	F	270	ILE
1	I	263	VAL
1	M	3	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/220 (92%)	192 (95%)	11 (5%)	22	44
1	B	204/220 (93%)	191 (94%)	13 (6%)	17	35
1	C	208/220 (94%)	197 (95%)	11 (5%)	22	45
1	D	209/220 (95%)	196 (94%)	13 (6%)	18	37
1	E	206/220 (94%)	195 (95%)	11 (5%)	22	45
1	F	205/220 (93%)	194 (95%)	11 (5%)	22	44
1	G	203/220 (92%)	191 (94%)	12 (6%)	19	39
1	H	204/220 (93%)	194 (95%)	10 (5%)	25	48
1	I	208/220 (94%)	196 (94%)	12 (6%)	20	40
1	J	205/220 (93%)	194 (95%)	11 (5%)	22	44
1	K	205/220 (93%)	193 (94%)	12 (6%)	19	39
1	L	206/220 (94%)	195 (95%)	11 (5%)	22	45

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	M	205/220 (93%)	195 (95%)	10 (5%)	25	48
1	N	204/220 (93%)	190 (93%)	14 (7%)	15	31
1	O	205/220 (93%)	194 (95%)	11 (5%)	22	44
1	P	204/220 (93%)	193 (95%)	11 (5%)	22	44
1	Q	206/220 (94%)	194 (94%)	12 (6%)	20	40
1	R	207/220 (94%)	193 (93%)	14 (7%)	16	32
1	S	203/220 (92%)	191 (94%)	12 (6%)	19	39
1	T	203/220 (92%)	191 (94%)	12 (6%)	19	39
All	All	4103/4400 (93%)	3869 (94%)	234 (6%)	20	41

5 of 234 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	4	PHE
1	S	222	MET
1	M	136	MET
1	S	173	ARG
1	R	70	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 122 such sidechains are listed below:

Mol	Chain	Res	Type
1	J	160	GLN
1	R	214	ASN
1	L	160	GLN
1	R	170	HIS
1	T	95	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	13P	N	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	C	501	1	8,8,9	1.55	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	L	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	S	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	M	501	1	8,8,9	1.55	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	I	501	1	8,8,9	1.51	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	D	501	1	8,8,9	1.53	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	A	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	J	501	1	8,8,9	1.53	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	P	501	1	8,8,9	1.53	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	H	501	1	8,8,9	1.53	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	T	501	1	8,8,9	1.55	1 (12%)	10,10,12	1.33	2 (20%)
2	13P	O	501	1	8,8,9	1.53	1 (12%)	10,10,12	1.36	2 (20%)
2	13P	K	501	1	8,8,9	1.51	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	R	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	B	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	E	501	1	8,8,9	1.54	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	G	501	1	8,8,9	1.55	1 (12%)	10,10,12	1.35	2 (20%)
2	13P	Q	501	1	8,8,9	1.52	1 (12%)	10,10,12	1.34	2 (20%)
2	13P	F	501	1	8,8,9	1.55	1 (12%)	10,10,12	1.34	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	13P	N	501	1	-	4/6/6/8	-
2	13P	C	501	1	-	4/6/6/8	-
2	13P	L	501	1	-	4/6/6/8	-
2	13P	S	501	1	-	4/6/6/8	-
2	13P	M	501	1	-	4/6/6/8	-
2	13P	I	501	1	-	4/6/6/8	-
2	13P	D	501	1	-	4/6/6/8	-
2	13P	A	501	1	-	4/6/6/8	-
2	13P	J	501	1	-	4/6/6/8	-
2	13P	P	501	1	-	4/6/6/8	-
2	13P	H	501	1	-	4/6/6/8	-
2	13P	T	501	1	-	4/6/6/8	-
2	13P	O	501	1	-	4/6/6/8	-
2	13P	K	501	1	-	4/6/6/8	-
2	13P	R	501	1	-	4/6/6/8	-
2	13P	B	501	1	-	4/6/6/8	-
2	13P	E	501	1	-	4/6/6/8	-
2	13P	G	501	1	-	4/6/6/8	-
2	13P	Q	501	1	-	4/6/6/8	-
2	13P	F	501	1	-	4/6/6/8	-

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	M	501	13P	P-O1P	3.46	1.61	1.50
2	G	501	13P	P-O1P	3.45	1.61	1.50
2	C	501	13P	P-O1P	3.45	1.61	1.50
2	S	501	13P	P-O1P	3.45	1.61	1.50
2	F	501	13P	P-O1P	3.45	1.61	1.50

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	501	13P	C1-C2-C3	3.13	119.73	113.55
2	A	501	13P	C1-C2-C3	3.12	119.70	113.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	O	501	13P	C1-C2-C3	3.12	119.69	113.55
2	J	501	13P	C1-C2-C3	3.10	119.67	113.55
2	M	501	13P	C1-C2-C3	3.10	119.67	113.55

There are no chirality outliers.

5 of 80 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	13P	C1-O1-P-O1P
2	A	501	13P	C1-O1-P-O2P
2	A	501	13P	C1-O1-P-O3P
2	A	501	13P	C1-C2-C3-O3
2	B	501	13P	C1-O1-P-O1P

There are no ring outliers.

20 monomers are involved in 40 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	N	501	13P	2	0
2	C	501	13P	2	0
2	L	501	13P	2	0
2	S	501	13P	2	0
2	M	501	13P	2	0
2	I	501	13P	2	0
2	D	501	13P	2	0
2	A	501	13P	2	0
2	J	501	13P	2	0
2	P	501	13P	2	0
2	H	501	13P	2	0
2	T	501	13P	2	0
2	O	501	13P	2	0
2	K	501	13P	2	0
2	R	501	13P	2	0
2	B	501	13P	2	0
2	E	501	13P	2	0
2	G	501	13P	2	0
2	Q	501	13P	2	0
2	F	501	13P	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/273 (96%)	-0.05	5 (1%) 66 62	33, 58, 98, 119	0
1	B	264/273 (96%)	-0.05	5 (1%) 66 62	34, 57, 85, 115	0
1	C	265/273 (97%)	-0.02	3 (1%) 80 78	34, 55, 86, 122	0
1	D	266/273 (97%)	-0.40	0 100 100	32, 54, 86, 107	0
1	E	264/273 (96%)	-0.30	0 100 100	36, 52, 85, 130	0
1	F	264/273 (96%)	-0.12	2 (0%) 86 84	37, 61, 95, 115	0
1	G	264/273 (96%)	0.06	4 (1%) 73 70	34, 62, 90, 110	0
1	H	264/273 (96%)	0.04	4 (1%) 73 70	36, 60, 91, 124	0
1	I	264/273 (96%)	-0.28	1 (0%) 92 91	31, 51, 78, 119	0
1	J	264/273 (96%)	-0.24	1 (0%) 92 91	30, 52, 83, 122	0
1	K	264/273 (96%)	-0.42	0 100 100	35, 54, 82, 118	0
1	L	265/273 (97%)	-0.25	4 (1%) 73 70	36, 57, 88, 115	0
1	M	264/273 (96%)	-0.02	7 (2%) 54 48	38, 59, 94, 129	0
1	N	264/273 (96%)	-0.08	6 (2%) 60 54	35, 58, 88, 130	0
1	O	264/273 (96%)	-0.30	2 (0%) 86 84	34, 54, 84, 163	0
1	P	264/273 (96%)	-0.36	1 (0%) 92 91	36, 56, 83, 115	0
1	Q	264/273 (96%)	-0.32	0 100 100	32, 53, 82, 109	0
1	R	266/273 (97%)	0.03	13 (4%) 29 23	39, 60, 98, 141	0
1	S	264/273 (96%)	0.10	10 (3%) 40 33	35, 66, 98, 122	0
1	T	264/273 (96%)	-0.19	2 (0%) 86 84	41, 61, 92, 107	0
All	All	5286/5460 (96%)	-0.16	70 (1%) 77 73	30, 57, 90, 163	0

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	S	246	VAL	5.4
1	S	47	ILE	5.1
1	T	268	LYS	4.1
1	C	1	MET	4.0
1	R	211	PRO	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	13P	O	501	9/10	0.92	0.15	62,64,65,65	0
2	13P	D	501	9/10	0.93	0.19	61,63,65,65	0
2	13P	S	501	9/10	0.93	0.14	67,69,70,71	0
2	13P	A	501	9/10	0.94	0.16	66,66,68,69	0
2	13P	F	501	9/10	0.94	0.17	65,67,72,72	0
2	13P	M	501	9/10	0.95	0.13	66,68,68,70	0
2	13P	G	501	9/10	0.95	0.17	66,66,67,68	0
2	13P	H	501	9/10	0.95	0.15	64,65,66,67	0
2	13P	C	501	9/10	0.96	0.13	66,67,69,69	0
2	13P	N	501	9/10	0.96	0.14	64,65,67,67	0
2	13P	I	501	9/10	0.96	0.18	58,61,64,64	0
2	13P	P	501	9/10	0.96	0.17	64,65,66,66	0
2	13P	Q	501	9/10	0.96	0.18	65,66,68,68	0
2	13P	R	501	9/10	0.96	0.13	68,69,71,72	0
2	13P	L	501	9/10	0.96	0.14	64,65,67,67	0
2	13P	T	501	9/10	0.96	0.14	64,66,67,67	0
2	13P	B	501	9/10	0.97	0.17	62,63,66,66	0
2	13P	J	501	9/10	0.97	0.14	62,64,67,67	0
2	13P	K	501	9/10	0.97	0.17	64,66,67,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	13P	E	501	9/10	0.98	0.18	63,65,67,69	0

6.5 Other polymers [i](#)

There are no such residues in this entry.