



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 06:26 AM EDT

PDB ID : 2R1Z  
Title : Crystal Structure of the BARD1 BRCT Repeat  
Authors : Lee, M.S.; Edwards, R.A.; Williams, R.S.; Glover, M.J.N.  
Deposited on : 2007-08-23  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	 23% 81% 14%
1	B	209	 25% 84% 14%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BRCA1-associated RING domain protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	Total 1630	C 1046	N 279	O 294	S 11	0	1	0
1	B	208	Total 1682	C 1081	N 288	O 302	S 11	0	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0
2	B	1	Total 6	C 3	O 3	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	57	Total 57	O 57	0	0
3	B	58	Total 58	O 58	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BRCA1-associated RING domain protein 1



- Molecule 1: BRCA1-associated RING domain protein 1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.82Å 75.55Å 117.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.49 – 2.10 27.76 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.0 (29.49-2.10) 95.0 (27.76-2.10)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.80 (at 2.10Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.222 , 0.262 0.220 , 0.260	Depositor DCC
$R_{free}$ test set	1445 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	43.4	Xtrriage
Anisotropy	0.258	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 58.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3439	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	3/1673 (0.2%)	0.60	0/2259
1	B	0.43	0/1724	0.59	0/2333
All	All	0.48	3/3397 (0.1%)	0.60	0/4592

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	587	GLU	CD-OE2	6.76	1.33	1.25
1	A	587	GLU	CD-OE1	6.38	1.32	1.25
1	A	587	GLU	CG-CD	5.90	1.60	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1630	0	1650	22	0
1	B	1682	0	1696	19	0
2	A	6	0	8	0	0
2	B	6	0	8	2	0
3	A	57	0	0	0	0
3	B	58	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3439	0	3362	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:621:MET:HG2	1:B:768:MET:HE3	1.56	0.87
1:B:621:MET:HG2	1:B:768:MET:CE	2.04	0.86
1:A:730:GLN:HE22	1:A:774:PRO:HA	1.54	0.73
1:A:714:THR:HG23	1:A:735:GLN:OE1	1.97	0.65
1:B:578:SER:H	1:B:581:GLN:HE21	1.46	0.64
1:A:578:SER:H	1:A:581:GLN:HE21	1.46	0.64
1:B:730:GLN:HE22	1:B:774:PRO:HA	1.63	0.64
1:A:621:MET:HG2	1:A:768:MET:CE	2.29	0.62
1:B:575:SER:O	1:B:610:PRO:HD3	1.98	0.62
1:B:670:LYS:HE3	2:B:802:GOL:H11	1.81	0.61
1:B:621:MET:HG2	1:B:768:MET:HE1	1.82	0.61
1:A:724:ALA:HB3	1:A:731:ARG:HD2	1.83	0.59
1:B:711:SER:HB3	1:B:714:THR:HG23	1.85	0.59
1:A:621:MET:HG2	1:A:768:MET:HE1	1.90	0.53
1:B:574:GLY:HA3	1:B:577:LEU:HD22	1.92	0.51
1:A:621:MET:HG2	1:A:768:MET:HE3	1.91	0.51
1:B:670:LYS:HE3	2:B:802:GOL:C1	2.41	0.51
1:B:730:GLN:NE2	1:B:774:PRO:HA	2.27	0.50
1:B:572:LEU:HD23	1:B:606:HIS:HB2	1.94	0.50
1:A:652:GLU:OE2	1:A:661:ARG:HD3	2.13	0.49
1:A:629:TRP:CZ3	1:A:661:ARG:HD2	2.48	0.48
1:A:574:GLY:HA3	1:A:577:LEU:HD22	1.94	0.48
1:A:646:GLU:HG3	1:A:651:TYR:CE1	2.49	0.47
1:B:578:SER:H	1:B:581:GLN:NE2	2.12	0.47
1:B:631:LEU:HD21	1:B:648:GLU:HB2	1.97	0.47
1:A:646:GLU:CG	1:A:651:TYR:CE1	2.98	0.46
1:B:609:VAL:HG13	3:B:43:HOH:O	2.16	0.45
1:A:646:GLU:HG2	1:A:651:TYR:CZ	2.51	0.45
1:A:717:ILE:HD13	1:A:717:ILE:HA	1.81	0.45
1:A:646:GLU:HG3	1:A:651:TYR:HE1	1.82	0.45
1:A:730:GLN:NE2	1:A:774:PRO:HA	2.25	0.45
1:A:581:GLN:HA	1:A:584:MET:HE3	1.98	0.44
1:B:768:MET:HA	1:B:768:MET:HE2	1.99	0.44

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:629:TRP:CD2	1:A:648:GLU:HG3	2.54	0.43
1:A:631:LEU:HD21	1:A:648:GLU:HB2	2.01	0.43
1:B:725:ARG:HA	1:B:726:PRO:HD3	1.88	0.43
1:A:573:ILE:HD12	1:A:619:LYS:HD3	2.02	0.41
1:B:614:VAL:HG21	1:B:624:ILE:HD11	2.02	0.41
1:A:629:TRP:CE2	1:A:648:GLU:HG3	2.56	0.41
1:A:577:LEU:HG	1:A:581:GLN:HB3	2.03	0.40
1:B:652:GLU:OE2	1:B:661:ARG:HD3	2.22	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/209 (95%)	195 (98%)	4 (2%)	0	100	100
1	B	206/209 (99%)	198 (96%)	8 (4%)	0	100	100
All	All	405/418 (97%)	393 (97%)	12 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/189 (97%)	171 (93%)	12 (7%)	16	14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	188/189 (100%)	176 (94%)	12 (6%)	17	14
All	All	371/378 (98%)	347 (94%)	24 (6%)	17	14

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	573	ILE
1	A	577	LEU
1	A	609	VAL
1	A	642	ARG
1	A	661	ARG
1	A	664	ARG
1	A	671	LEU
1	A	705	ARG
1	A	718	ASN
1	A	741	ASP
1	A	749	ARG
1	A	755	VAL
1	B	577	LEU
1	B	580	GLU
1	B	609	VAL
1	B	626	ASN
1	B	641	ARG
1	B	649	GLU
1	B	671	LEU
1	B	714	THR
1	B	727	ASP
1	B	744	ASN
1	B	748	GLU
1	B	755	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	581	GLN
1	A	626	ASN
1	A	663	ASN
1	A	690	ASN
1	A	718	ASN
1	A	730	GLN
1	A	752	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	581	GLN
1	B	690	ASN
1	B	730	GLN
1	B	744	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	B	802	-	5,5,5	0.31	0	5,5,5	0.25	0
2	GOL	A	801	-	5,5,5	0.33	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	802	-	-	0/4/4/4	-
2	GOL	A	801	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	802	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	202/209 (96%)	1.31	49 (24%) 0 0	37, 46, 61, 70	0
1	B	208/209 (99%)	1.37	52 (25%) 0 0	37, 46, 56, 65	0
All	All	410/418 (98%)	1.34	101 (24%) 0 0	37, 46, 60, 70	0

All (101) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	777	SER	8.9
1	B	712	ASP	7.9
1	B	710	ASP	6.1
1	B	667	LEU	4.9
1	B	709	PRO	4.9
1	B	607	VAL	4.7
1	B	624	ILE	4.6
1	B	644	VAL	4.6
1	A	685	HIS	4.5
1	A	727	ASP	4.5
1	B	625	LEU	4.4
1	A	712	ASP	4.3
1	A	591	ILE	4.3
1	A	716	THR	4.3
1	A	630	ILE	4.1
1	A	644	VAL	4.1
1	B	713	VAL	4.0
1	B	569	PRO	4.0
1	B	630	ILE	3.8
1	B	570	LEU	3.8
1	B	579	SER	3.6
1	A	741	ASP	3.6
1	A	578	SER	3.5
1	B	716	THR	3.4

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	598	THR	3.4
1	A	579	SER	3.4
1	A	569	PRO	3.4
1	B	614	VAL	3.4
1	B	767	VAL	3.4
1	B	665	GLU	3.4
1	B	711	SER	3.4
1	A	726	PRO	3.4
1	B	597	TYR	3.3
1	B	705	ARG	3.3
1	A	733	CYS	3.3
1	B	645	CYS	3.3
1	B	722	TYR	3.3
1	B	666	GLN	3.2
1	B	620	CYS	3.2
1	A	590	VAL	3.2
1	B	776	ASP	3.1
1	A	586	SER	3.1
1	A	717	ILE	3.1
1	A	602	SER	3.0
1	B	580	GLU	3.0
1	A	645	CYS	3.0
1	A	749	ARG	2.9
1	B	643	LYS	2.9
1	A	594	ALA	2.9
1	B	599	GLU	2.9
1	A	684	LYS	2.9
1	A	710	ASP	2.9
1	B	733	CYS	2.8
1	A	732	PHE	2.8
1	A	597	TYR	2.8
1	A	736	TYR	2.8
1	B	608	VAL	2.7
1	A	776	ASP	2.7
1	A	755	VAL	2.7
1	B	662	LEU	2.7
1	B	708	LYS	2.7
1	B	668	LEU	2.7
1	A	653	ILE	2.6
1	B	715	GLN	2.6
1	B	743	CYS	2.6
1	A	713	VAL	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	631	LEU	2.5
1	A	657	PRO	2.4
1	A	603	THR	2.4
1	A	711	SER	2.4
1	A	592	LEU	2.4
1	B	621	MET	2.4
1	B	749	ARG	2.3
1	A	593	LYS	2.3
1	B	631	LEU	2.3
1	B	742	LEU	2.3
1	B	584	MET	2.3
1	A	725	ARG	2.3
1	B	689	ASP	2.3
1	B	732	PHE	2.3
1	B	658	ARG	2.3
1	A	754	LYS	2.3
1	A	753	GLY	2.3
1	A	620[A]	CYS	2.2
1	A	767	VAL	2.2
1	A	756	TRP	2.2
1	B	726	PRO	2.2
1	A	580	GLU	2.2
1	B	613	ALA	2.2
1	B	727	ASP	2.2
1	B	714	THR	2.1
1	A	608	VAL	2.1
1	A	589	ALA	2.1
1	A	624	ILE	2.1
1	A	686	HIS	2.1
1	B	641	ARG	2.1
1	B	775	LEU	2.1
1	A	714	THR	2.0
1	B	671	LEU	2.0
1	A	675	CYS	2.0
1	B	766	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	802	6/6	0.69	0.34	46,47,48,48	6
2	GOL	A	801	6/6	0.96	0.28	48,48,49,51	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.