



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 23, 2023 – 02:54 PM EDT

PDB ID : 3C1N
Title : Crystal Structure of Allosteric Inhibition Threonine-sensitive Aspartokinase from *Methanococcus jannaschii* with L-threonine
Authors : Liu, X.; Pavlovshy, A.G.; Viola, R.E.
Deposited on : 2008-01-23
Resolution : 2.72 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3359 (2.74-2.70)
Clashscore	141614	3686 (2.74-2.70)
Ramachandran outliers	138981	3622 (2.74-2.70)
Sidechain outliers	138945	3623 (2.74-2.70)
RSRZ outliers	127900	3276 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	
1	B	473	
1	C	473	
1	D	473	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	THR	A	471	-	-	X	-
2	THR	A	473	-	-	X	-
2	THR	B	470	-	X	X	-
2	THR	B	474	-	-	X	-
2	THR	C	475	-	-	-	X
2	THR	D	472	-	-	X	X

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 13893 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable aspartokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	458	Total 3456	C 2193	N 573	O 680	S 10	0	0	0
1	B	459	Total 3459	C 2194	N 574	O 682	S 9	0	0	0
1	C	456	Total 3441	C 2186	N 571	O 674	S 10	0	0	0
1	D	452	Total 3410	C 2167	N 567	O 667	S 9	0	0	0

- Molecule 2 is THREONINE (three-letter code: THR) (formula: C₄H₉NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 8	C 4	N 1	O 3	0	0
2	A	1	Total 8	C 4	N 1	O 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			8	4	1	3		
2	B	1	Total	C	N	O	0	0
			8	4	1	3		
2	C	1	Total	C	N	O	0	0
			8	4	1	3		
2	D	1	Total	C	N	O	0	0
			8	4	1	3		
2	D	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	O	0	0
			18	18		
3	B	20	Total	O	0	0
			20	20		
3	C	19	Total	O	0	0
			19	19		
3	D	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable aspartokinase




- Molecule 1: Probable aspartokinase



- Molecule 1: Probable aspartokinase



● Molecule 1: Probable aspartokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	110.04Å 144.32Å 155.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.72 14.98 – 2.72	Depositor EDS
% Data completeness (in resolution range)	84.4 (50.00-2.72) 84.4 (14.98-2.72)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 2.73Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.236 , 0.290 0.226 , 0.279	Depositor DCC
R_{free} test set	2872 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å ²)	75.3	Xtrriage
Anisotropy	0.213	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 73.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13893	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	1/3492 (0.0%)	0.59	1/4708 (0.0%)
1	B	0.42	0/3496	0.57	1/4716 (0.0%)
1	C	0.36	0/3478	0.53	0/4691
1	D	0.44	2/3445 (0.1%)	0.53	0/4642
All	All	0.41	3/13911 (0.0%)	0.56	2/18757 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	82	GLU	CD-OE2	10.28	1.36	1.25
1	D	82	GLU	CD-OE1	10.13	1.36	1.25
1	A	86	SER	CA-CB	5.58	1.61	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	LEU	CA-CB-CG	7.29	132.07	115.30
1	B	390	LEU	CA-CB-CG	5.48	127.91	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3456	0	3549	63	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3459	0	3548	68	0
1	C	3441	0	3541	41	0
1	D	3410	0	3511	61	0
2	A	16	0	12	12	0
2	B	16	0	12	10	0
2	C	8	0	6	1	0
2	D	16	0	12	7	0
3	A	18	0	0	1	0
3	B	20	0	0	0	0
3	C	19	0	0	0	0
3	D	14	0	0	0	0
All	All	13893	0	14191	228	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 228 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:440:GLN:HE22	2:A:471:THR:HG21	1.15	1.08
1:B:8:GLY:HA3	2:B:474:THR:HA	1.30	1.07
1:B:11:SER:HG	2:B:474:THR:N	1.56	1.03
1:D:351:LEU:HB2	1:D:364:VAL:HG13	1.47	0.95
1:A:328:MET:HG2	1:A:334:THR:HG21	1.49	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	450/473 (95%)	421 (94%)	24 (5%)	5 (1%)	14 32

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	453/473 (96%)	423 (93%)	26 (6%)	4 (1%)	17	38
1	C	450/473 (95%)	412 (92%)	31 (7%)	7 (2%)	9	23
1	D	442/473 (93%)	409 (92%)	28 (6%)	5 (1%)	14	32
All	All	1795/1892 (95%)	1665 (93%)	109 (6%)	21 (1%)	13	30

5 of 21 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	307	SER
1	C	161	ASP
1	C	300	ASN
1	A	161	ASP
1	A	328	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/397 (94%)	341 (91%)	34 (9%)	9	21
1	B	375/397 (94%)	344 (92%)	31 (8%)	11	25
1	C	373/397 (94%)	344 (92%)	29 (8%)	12	28
1	D	369/397 (93%)	337 (91%)	32 (9%)	10	23
All	All	1492/1588 (94%)	1366 (92%)	126 (8%)	10	24

5 of 126 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	363	LEU
1	D	191	THR
1	C	104	LEU
1	D	184	GLU
1	D	370	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	391	ASN
1	B	440	GLN
1	D	391	ASN
1	C	54	GLN
1	A	300	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	THR	A	473	-	6,7,7	1.10	1 (16%)	7,9,9	1.00	0
2	THR	B	474	-	6,7,7	1.13	1 (16%)	7,9,9	0.91	0
2	THR	C	475	-	6,7,7	0.88	0	7,9,9	1.40	2 (28%)
2	THR	D	476	-	6,7,7	0.96	1 (16%)	7,9,9	1.32	2 (28%)
2	THR	B	470	-	6,7,7	1.12	1 (16%)	7,9,9	1.16	1 (14%)
2	THR	D	472	-	6,7,7	1.01	1 (16%)	7,9,9	1.15	1 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	THR	A	471	-	6,7,7	0.93	0	7,9,9	1.22	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	THR	A	473	-	-	2/8/8/8	-
2	THR	B	474	-	-	0/8/8/8	-
2	THR	C	475	-	-	1/8/8/8	-
2	THR	D	476	-	-	1/8/8/8	-
2	THR	B	470	-	-	8/8/8/8	-
2	THR	D	472	-	-	5/8/8/8	-
2	THR	A	471	-	-	4/8/8/8	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	474	THR	OXT-C	-2.55	1.22	1.30
2	B	470	THR	OXT-C	-2.37	1.22	1.30
2	A	473	THR	OXT-C	-2.34	1.22	1.30
2	D	472	THR	OXT-C	-2.20	1.23	1.30
2	D	476	THR	OXT-C	-2.17	1.23	1.30

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	475	THR	OXT-C-O	-2.95	117.40	124.09
2	A	471	THR	OXT-C-O	-2.64	118.09	124.09
2	D	472	THR	OXT-C-O	-2.62	118.14	124.09
2	D	476	THR	OXT-C-O	-2.57	118.25	124.09
2	B	470	THR	OXT-C-O	-2.51	118.39	124.09

There are no chirality outliers.

5 of 21 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	471	THR	C-CA-CB-OG1
2	A	471	THR	C-CA-CB-CG2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	473	THR	OXT-C-CA-N
2	B	470	THR	N-CA-CB-OG1
2	B	470	THR	C-CA-CB-OG1

There are no ring outliers.

7 monomers are involved in 30 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	473	THR	6	0
2	B	474	THR	6	0
2	C	475	THR	1	0
2	D	476	THR	2	0
2	B	470	THR	4	0
2	D	472	THR	5	0
2	A	471	THR	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/473 (96%)	0.21	23 (5%) 28 28	50, 61, 72, 76	0
1	B	459/473 (97%)	0.22	16 (3%) 44 44	49, 61, 72, 79	0
1	C	456/473 (96%)	0.38	35 (7%) 13 12	49, 63, 73, 78	0
1	D	452/473 (95%)	0.50	60 (13%) 3 2	61, 69, 76, 80	0
All	All	1825/1892 (96%)	0.33	134 (7%) 15 13	49, 64, 74, 80	0

The worst 5 of 134 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	382	GLY	8.6
1	B	331	VAL	7.4
1	C	331	VAL	7.2
1	A	302	MET	5.1
1	C	469	GLU	4.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	THR	C	475	8/8	0.34	0.51	51,53,53,53	8
2	THR	D	472	8/8	0.71	0.67	71,71,71,71	8
2	THR	D	476	8/8	0.78	0.35	62,62,62,62	8
2	THR	B	470	8/8	0.87	0.25	68,69,70,70	0
2	THR	A	473	8/8	0.87	0.27	66,67,67,67	0
2	THR	B	474	8/8	0.90	0.21	60,60,60,61	0
2	THR	A	471	8/8	0.93	0.18	57,58,58,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.