



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 23, 2023 – 08:10 AM EDT

PDB ID : 3EB0  
Title : Crystal Structure of cgd4\_240 from cryptosporidium Parvum in complex with indirubin E804  
Authors : Wernimont, A.K.; Fedorov, O.; Lam, A.; Ali, A.; Zhao, Y.; Lew, J.; Wasney, G.; Vedadi, M.; Kozieradzki, I.; Schapira, M.; Bochkarev, A.; Wilstrom, M.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.M.; Hui, R.; Lin, Y.H.; Structural Genomics Consortium (SGC)  
Deposited on : 2008-08-26  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	
1	C	383	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	386	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	312	Total	C	N	O	P	S	0	2	0
			2503	1633	414	442	1	13			
1	C	311	Total	C	N	O	P	S	0	1	0
			2490	1621	415	440	1	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP A3FQN0
C	1	GLY	-	expression tag	UNP A3FQN0

- Molecule 2 is 3-({[(3S)-3,4-dihydroxybutyl]oxy}amino)-1H,2'H-2,3'-biindol-2'-one (three-letter code: DRK) (formula: C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total	C	N	O	0	0
			27	20	3	4		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			27	20	3	4		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	86	Total	O	0	0
			86	86		
4	C	64	Total	O	0	0
			64	64		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein



- Molecule 1: Putative uncharacterized protein



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.10Å 138.37Å 43.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.65 34.59 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.5 (35.00-2.65) 99.5 (34.59-2.65)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 2.65Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.238 , 0.295 0.236 , 0.297	Depositor DCC
$R_{free}$ test set	1234 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.3	Xtrriage
Anisotropy	0.387	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5221	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 58.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.9191e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DRK, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.54	0/2543	0.64	2/3442 (0.1%)
1	C	0.55	0/2530	0.64	0/3427
All	All	0.54	0/5073	0.64	2/6869 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128[A]	LEU	CA-CB-CG	5.05	126.91	115.30
1	A	128[B]	LEU	CA-CB-CG	5.05	126.91	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2503	0	2528	33	1
1	C	2490	0	2500	25	1
2	A	27	0	19	3	0
2	C	27	0	19	3	0
3	A	12	0	16	0	0
3	C	12	0	16	2	0
4	A	86	0	0	5	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	64	0	0	4	0
All	All	5221	0	5098	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:GLY:HA2	1:A:25:GLU:O	1.69	0.93
2:C:384:DRK:HNAO	2:C:384:DRK:HAK	1.40	0.86
2:A:384:DRK:HAK	2:A:384:DRK:HNAO	1.39	0.85
1:C:268:ARG:NH2	4:C:440:HOH:O	2.22	0.72
1:A:12:GLY:CA	1:A:25:GLU:O	2.42	0.67
1:A:66:VAL:HB	1:A:116:ILE:HG22	1.77	0.65
1:C:287:ARG:HG2	1:C:299:ILE:HD13	1.79	0.64
2:C:384:DRK:HAK	2:C:384:DRK:NAO	2.11	0.64
1:A:154:VAL:HG21	1:A:228:ILE:HD11	1.79	0.63
3:C:386:GOL:H31	4:C:531:HOH:O	1.97	0.62
1:A:204:SER:H	1:A:208:ARG:HD2	1.64	0.62
1:A:71:THR:HB	1:A:114:ASN:HD21	1.66	0.61
1:A:192:LEU:O	1:A:194:PRO:HD3	2.01	0.61
1:C:307:ARG:HD2	4:C:417:HOH:O	2.01	0.60
1:C:315:ASN:HD21	3:C:385:GOL:H32	1.66	0.59
1:C:64:LYS:HG3	4:C:389:HOH:O	2.02	0.58
1:A:7:LYS:HD3	4:A:533:HOH:O	2.04	0.57
1:A:152:ARG:HA	1:A:320:MET:HE1	1.87	0.56
1:C:151:PHE:HB3	1:C:320:MET:HE2	1.87	0.56
1:A:151:PHE:HB3	1:A:320:MET:HE2	1.86	0.56
1:C:290:LEU:HB3	1:C:291:PRO:HD2	1.87	0.55
1:A:290:LEU:HB3	1:A:291:PRO:HD2	1.86	0.55
1:A:170:GLN:HB3	2:A:384:DRK:HAL	1.89	0.54
1:A:142:LEU:HA	1:A:145:ILE:HD12	1.90	0.52
1:A:110:ASN:N	4:A:452:HOH:O	2.42	0.52
1:A:165:ARG:HG3	1:A:223:ILE:HD11	1.92	0.52
1:A:233:GLY:HA3	1:A:241:LEU:HD11	1.93	0.51
1:A:164:HIS:O	1:A:165:ARG:HB2	2.10	0.51
1:A:138:ILE:HD12	1:A:235:LEU:HD22	1.93	0.50
1:A:371:ARG:HD3	4:A:447:HOH:O	2.10	0.49
1:C:71:THR:HG22	1:C:112:TYR:HB2	1.94	0.49
1:C:63:ILE:HD13	1:C:173:LEU:HD12	1.95	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:152:ARG:HA	1:C:320:MET:HE1	1.94	0.48
2:A:384:DRK:HAK	2:A:384:DRK:NAO	2.20	0.47
1:C:204:SER:H	1:C:208:ARG:HD2	1.80	0.46
1:A:201:PTR:HD2	1:A:201:PTR:C	2.45	0.46
1:A:276:VAL:HG12	1:A:276:VAL:O	2.15	0.46
1:A:71:THR:HB	1:A:114:ASN:ND2	2.30	0.46
1:C:142:LEU:HD11	1:C:146:TYR:CZ	2.51	0.45
1:A:173:LEU:O	1:A:181:LEU:HA	2.17	0.45
2:C:384:DRK:HNAO	2:C:384:DRK:CAK	2.20	0.45
1:C:201:PTR:C	1:C:201:PTR:HD2	2.47	0.45
1:A:6:SER:N	4:A:449:HOH:O	2.49	0.45
1:C:206:PHE:HB2	1:C:207:TYR:CE2	2.51	0.45
1:C:220:THR:HB	1:C:221:PRO:CD	2.47	0.45
1:C:217:THR:HG22	1:C:218:GLU:HG3	1.99	0.44
1:C:278:PHE:HB3	1:C:279:PRO:HD2	1.98	0.44
1:A:300:ASP:OD1	1:A:304:GLN:NE2	2.51	0.44
1:C:204:SER:O	1:C:208:ARG:HD2	2.18	0.43
1:C:309:GLU:HB2	1:C:312:LEU:HB2	2.02	0.42
1:A:138:ILE:CD1	1:A:235:LEU:HD22	2.50	0.41
1:A:208:ARG:NH1	1:A:250:GLN:OE1	2.53	0.41
1:A:59:HIS:HB3	1:A:62:ILE:HD12	2.01	0.41
1:A:166:ASP:O	1:A:171:ASN:ND2	2.53	0.41
1:C:53:ILE:O	1:C:54:MET:C	2.58	0.41
1:A:64:LYS:HG3	4:A:393:HOH:O	2.20	0.41
1:C:234:GLU:HG3	1:C:239:LYS:C	2.42	0.41
1:C:322:HIS:CG	1:C:323:PRO:HD2	2.56	0.41
1:A:248:ILE:H	1:A:248:ILE:HG13	1.51	0.41
1:A:128[A]:LEU:HD12	1:A:235:LEU:HD23	2.04	0.40
1:A:234:GLU:HG3	1:A:239:LYS:C	2.42	0.40
1:C:166:ASP:O	1:C:171:ASN:ND2	2.54	0.40
1:C:249:ASP:O	1:C:253:ARG:HG2	2.21	0.40
1:C:206:PHE:HB2	1:C:207:TYR:CD2	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:321:ALA:O	1:C:287:ARG:NH1[4_556]	2.09	0.11

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/383 (79%)	286 (94%)	17 (6%)	0	100	100
1	C	301/383 (79%)	287 (95%)	12 (4%)	2 (1%)	22	33
All	All	604/766 (79%)	573 (95%)	29 (5%)	2 (0%)	41	56

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	55	LYS
1	C	185	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/349 (78%)	263 (96%)	11 (4%)	31	47
1	C	271/349 (78%)	262 (97%)	9 (3%)	38	54
All	All	545/698 (78%)	525 (96%)	20 (4%)	34	50

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LYS
1	A	26	VAL
1	A	40	VAL
1	A	63	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	71	THR
1	A	72	THR
1	A	137	SER
1	A	188	SER
1	A	217	THR
1	A	248	ILE
1	A	281	LEU
1	C	20	PHE
1	C	71	THR
1	C	188	SER
1	C	217	THR
1	C	245	GLU
1	C	263	LYS
1	C	312	LEU
1	C	327	HIS
1	C	352	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	114	ASN
1	C	352	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	PTR	A	201	1	15,16,17	2.01	2 (13%)	19,22,24	0.79	0
1	PTR	C	201	1	15,16,17	1.98	1 (6%)	19,22,24	0.79	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	201	1	-	2/10/11/13	0/1/1/1
1	PTR	C	201	1	-	2/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	201	PTR	OH-CZ	-7.21	1.24	1.40
1	A	201	PTR	OH-CZ	-7.17	1.24	1.40
1	A	201	PTR	P-OH	2.51	1.63	1.59

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	201	PTR	CA-CB-CG-CD1
1	A	201	PTR	CA-CB-CG-CD1
1	A	201	PTR	CA-CB-CG-CD2
1	C	201	PTR	CA-CB-CG-CD2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	201	PTR	1	0
1	C	201	PTR	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	C	386	-	5,5,5	0.38	0	5,5,5	0.46	0
3	GOL	A	386	-	5,5,5	0.38	0	5,5,5	0.39	0
3	GOL	A	385	-	5,5,5	0.45	0	5,5,5	0.25	0
3	GOL	C	385	-	5,5,5	0.41	0	5,5,5	0.14	0
2	DRK	C	384	-	25,30,30	1.15	1 (4%)	26,42,42	1.25	3 (11%)
2	DRK	A	384	-	25,30,30	1.13	1 (4%)	26,42,42	1.44	4 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	386	-	-	0/4/4/4	-
3	GOL	A	386	-	-	2/4/4/4	-
3	GOL	A	385	-	-	4/4/4/4	-
3	GOL	C	385	-	-	0/4/4/4	-
2	DRK	C	384	-	-	7/7/25/25	0/4/4/4
2	DRK	A	384	-	-	5/7/25/25	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	384	DRK	CAS-NAP	-2.75	1.32	1.37
2	C	384	DRK	CAS-NAP	-2.40	1.33	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	384	DRK	CAT-NAQ-CAX	4.04	112.21	103.90

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	384	DRK	CAZ-CAV-CAS	-3.53	104.90	107.90
2	C	384	DRK	CAT-NAQ-CAX	3.51	111.12	103.90
2	C	384	DRK	CAE-CAH-CAW	2.48	120.45	117.35
2	A	384	DRK	CAJ-CAY-CAU	-2.39	134.19	135.57
2	A	384	DRK	CAE-CAH-CAW	2.16	120.05	117.35
2	C	384	DRK	CAZ-CAV-CAS	-2.06	106.15	107.90

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	384	DRK	OAR-CAM-CAN-CBA
2	C	384	DRK	OAR-CAM-CAN-CBA
2	C	384	DRK	CAM-CAN-CBA-OAC
2	C	384	DRK	CAM-CAN-CBA-CAL
3	A	385	GOL	O1-C1-C2-C3
2	A	384	DRK	OAB-CAL-CBA-CAN
2	C	384	DRK	OAB-CAL-CBA-CAN
3	A	385	GOL	C1-C2-C3-O3
3	A	386	GOL	C1-C2-C3-O3
2	A	384	DRK	OAB-CAL-CBA-OAC
2	C	384	DRK	OAB-CAL-CBA-OAC
3	A	385	GOL	O1-C1-C2-O2
3	A	385	GOL	O2-C2-C3-O3
3	A	386	GOL	O2-C2-C3-O3
2	C	384	DRK	CAN-CAM-OAR-NAO
2	A	384	DRK	CAU-NAO-OAR-CAM
2	A	384	DRK	CAN-CAM-OAR-NAO
2	C	384	DRK	CAU-NAO-OAR-CAM


There are no ring outliers.

4 monomers are involved in 8 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	386	GOL	1	0
3	C	385	GOL	1	0
2	C	384	DRK	3	0
2	A	384	DRK	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight  $> 250$  and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/383 (81%)	0.24	18 (5%) 23 19	14, 20, 26, 34	14 (4%)
1	C	310/383 (80%)	0.30	13 (4%) 36 33	15, 20, 26, 30	6 (1%)
All	All	621/766 (81%)	0.27	31 (4%) 28 25	14, 20, 26, 34	20 (3%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	GLY	4.7
1	A	12	GLY	4.0
1	C	370	ASN	3.7
1	C	349	ASN	3.6
1	C	365	PRO	3.6
1	A	130	SER	3.5
1	C	369	LEU	3.5
1	A	264	GLU	3.4
1	A	21	GLY	3.3
1	A	287	ARG	3.2
1	C	110	ASN	3.1
1	A	330	ASN	3.1
1	A	350	ILE	3.1
1	A	70	TYR	3.0
1	A	136	ARG	2.8
1	C	358	PRO	2.7
1	A	46	TYR	2.7
1	C	350	ILE	2.7
1	C	136	ARG	2.6
1	C	21	GLY	2.6
1	A	20	PHE	2.4
1	A	267	ILE	2.4
1	C	331	SER	2.3
1	A	253	ARG	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	11	LEU	2.3
1	A	7	LYS	2.2
1	A	43	ASP	2.2
1	C	46	TYR	2.1
1	C	203	CYS	2.1
1	A	29	ILE	2.1
1	C	167	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PTR	A	201	16/17	0.95	0.14	18,19,20,21	0
1	PTR	C	201	16/17	0.96	0.15	14,16,19,20	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	C	385	6/6	0.57	0.28	58,59,60,60	0
3	GOL	A	386	6/6	0.72	0.42	62,62,63,63	0
3	GOL	C	386	6/6	0.73	0.20	42,44,45,45	0
3	GOL	A	385	6/6	0.80	0.17	49,49,50,50	0
2	DRK	A	384	27/27	0.87	0.26	27,29,44,45	1
2	DRK	C	384	27/27	0.88	0.27	33,36,48,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.