



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2023 – 11:18 PM EDT

PDB ID : 3F3W
Title : Drug resistant cSrc kinase domain in complex with inhibitor RL45 (Type II)
Authors : Grutter, C.; Kluter, S.; Getlik, M.; Rauh, D.
Deposited on : 2008-10-31
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	 10% 66% 22% • 9%
1	B	286	 6% 67% 20% • 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	261	2070	1334	337	380	19	0	0	0
1	B	260	2089	1343	347	380	19	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
A	338	MET	THR	engineered mutation	UNP P00523
A	345	CYS	SER	engineered mutation	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523
B	338	MET	THR	engineered mutation	UNP P00523
B	345	CYS	SER	engineered mutation	UNP P00523

- Molecule 2 is 1-{4-[(6-aminoquinazolin-4-yl)amino]phenyl}-3-[3-tert-butyl-1-(3-methylphenyl)-1H-pyrazol-5-yl]urea (three-letter code: 1BU) (formula: C₂₉H₃₀N₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	N	O	0	0
			38	29	8	1		
2	B	1	Total	C	N	O	0	0
			38	29	8	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	37	Total	O	0	0
			37	37		
3	B	36	Total	O	0	0
			36	36		


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.06Å 63.69Å 75.12Å 78.70° 87.02° 89.96°	Depositor
Resolution (Å)	40.00 – 2.60 43.58 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (40.00-2.60) 98.2 (43.58-2.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.47 (at 2.61Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.221 , 0.277 0.232 , 0.283	Depositor DCC
R_{free} test set	923 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.4	Xtrriage
Anisotropy	0.415	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	4308	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
1BU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	2/2122 (0.1%)	0.88	1/2878 (0.0%)
1	B	0.92	3/2141 (0.1%)	0.89	1/2899 (0.0%)
All	All	0.91	5/4263 (0.1%)	0.89	2/5777 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	345	CYS	CB-SG	-8.70	1.67	1.82
1	B	345	CYS	CB-SG	-6.49	1.71	1.82
1	B	504	GLU	CG-CD	5.21	1.59	1.51
1	A	504	GLU	CG-CD	5.18	1.59	1.51
1	B	470	GLU	CG-CD	5.12	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	388	ARG	NE-CZ-NH2	-5.07	117.76	120.30
1	B	386	ASP	CB-CG-OD1	5.04	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2070	0	2023	43	0
1	B	2089	0	2067	39	0
2	A	38	0	30	2	0
2	B	38	0	30	3	0
3	A	37	0	0	1	0
3	B	36	0	0	4	0
All	All	4308	0	4150	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:355:GLY:HA3	3:B:47:HOH:O	1.40	1.19
1:A:427:LYS:HE3	3:B:34:HOH:O	1.69	0.93
1:A:384:HIS:O	1:A:385:ARG:HB2	1.70	0.92
1:B:424:PHE:HD1	1:B:426:ILE:HG22	1.37	0.89
1:A:280:GLU:H	1:A:296:THR:HB	1.43	0.82
1:A:388:ARG:HB3	1:A:428:TRP:CD1	2.14	0.82
1:B:520:PHE:O	1:B:525:PRO:HA	1.83	0.79
1:A:323:VAL:HG21	1:A:403:ALA:HB2	1.66	0.76
1:A:314:MET:HE2	1:A:325:LEU:HB2	1.67	0.76
1:A:426:ILE:HD13	1:A:472:LEU:HG	1.71	0.72
1:B:343:LYS:O	3:B:62:HOH:O	2.07	0.71
1:A:485:PRO:O	3:A:54:HOH:O	2.08	0.70
1:B:383:VAL:HG12	1:B:385:ARG:HG3	1.73	0.69
1:A:485:PRO:O	1:A:486:GLU:HB2	1.94	0.68
1:A:314:MET:CE	1:A:325:LEU:HB2	2.25	0.67
1:B:526:GLN:H	1:B:526:GLN:CD	1.99	0.66
1:A:270:GLU:HG3	1:A:284:GLY:HA2	1.80	0.63
1:A:347:LEU:O	1:A:351:LYS:HG2	2.00	0.61
1:B:500:ARG:HD3	1:B:505:GLU:HB3	1.84	0.59
1:A:370:ILE:HG12	1:A:392:ILE:HD12	1.84	0.58
1:A:426:ILE:CD1	1:A:472:LEU:HG	2.33	0.58
1:A:322:LEU:HD22	1:A:402:VAL:HB	1.85	0.58
1:A:349:PHE:CE1	1:A:354:MET:HG3	2.39	0.57
1:B:324:GLN:O	1:B:338:MET:HB2	2.04	0.56
1:B:493:ASP:O	1:B:497:GLN:HG3	2.07	0.55
1:A:388:ARG:CB	1:A:428:TRP:CD1	2.89	0.55
1:B:384:HIS:O	1:B:385:ARG:HB2	2.07	0.55
1:B:424:PHE:CD1	1:B:426:ILE:HG22	2.29	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:526:GLN:NE2	1:B:526:GLN:C	2.59	0.55
1:A:460:ARG:NE	1:B:514:ALA:HB1	2.22	0.54
1:B:495:MET:O	1:B:498:CYS:HB2	2.07	0.54
1:B:526:GLN:C	1:B:526:GLN:HE21	2.10	0.54
2:B:1:1BU:HAM	2:B:1:1BU:N1	2.21	0.54
1:A:472:LEU:O	1:A:476:GLU:HB2	2.07	0.54
1:A:343:LYS:HE2	1:A:394:VAL:HG12	1.91	0.53
1:A:283:MET:HG3	1:A:284:GLY:N	2.23	0.52
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.44	0.52
1:A:362:GLN:O	1:A:366:MET:HG3	2.09	0.52
1:A:275:GLN:HG2	1:A:280:GLU:HG2	1.91	0.52
1:A:341:MET:HG3	1:A:393:LEU:HD13	1.92	0.52
1:A:324:GLN:O	1:A:338:MET:HB2	2.11	0.51
1:A:324:GLN:O	1:A:338:MET:CB	2.58	0.51
1:A:323:VAL:CG2	1:A:403:ALA:HB2	2.39	0.50
1:A:311:ALA:HB2	1:A:336:ILE:HD13	1.92	0.50
1:A:332:GLU:HA	1:A:333:PRO:C	2.32	0.50
1:B:515:PHE:HD2	3:B:40:HOH:O	1.93	0.49
1:A:287:ASN:O	1:A:289:THR:N	2.44	0.49
1:A:494:LEU:HD13	1:A:515:PHE:CE1	2.47	0.49
1:A:322:LEU:HD13	2:A:1:1BU:HAB	1.94	0.49
1:A:334:ILE:C	1:A:335:TYR:HD1	2.15	0.49
1:B:269:LEU:HG	1:B:294:ILE:HD13	1.94	0.48
1:B:526:GLN:CD	1:B:526:GLN:N	2.66	0.48
1:B:304:PRO:HG3	1:B:332:GLU:OE1	2.14	0.47
1:B:305:GLU:O	1:B:309:GLN:HG3	2.14	0.47
1:B:484:PRO:O	1:B:485:PRO:C	2.51	0.47
1:B:424:PHE:HB3	1:B:426:ILE:HG23	1.95	0.47
2:A:1:1BU:HAM	2:A:1:1BU:N1	2.28	0.47
1:B:476:GLU:OE2	1:B:476:GLU:HA	2.14	0.47
1:A:285:THR:CG2	1:A:288:GLY:HA2	2.45	0.46
1:B:261:GLU:OE2	1:B:330:SER:OG	2.25	0.46
1:B:378:GLU:HG3	1:B:441:ILE:HG12	1.98	0.46
1:A:384:HIS:O	1:A:385:ARG:CB	2.49	0.46
1:A:346:LEU:N	1:A:392:ILE:O	2.48	0.45
1:A:431:PRO:O	1:A:432:GLU:C	2.54	0.45
1:B:390:ALA:O	1:B:405:PHE:CE2	2.70	0.45
1:B:426:ILE:HD13	1:B:472:LEU:HG	1.99	0.45
1:A:302:MET:HG2	1:A:303:SER:N	2.32	0.44
1:B:466:MET:HA	1:B:470:GLU:OE1	2.17	0.44
1:A:378:GLU:HG3	1:A:441:ILE:HG12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:324:GLN:O	1:B:338:MET:CB	2.65	0.44
1:B:339:GLU:OE1	1:B:401:LYS:NZ	2.44	0.44
1:A:280:GLU:N	1:A:296:THR:HB	2.23	0.44
1:B:311:ALA:O	1:B:315:LYS:HG3	2.17	0.44
1:B:324:GLN:HG2	1:B:339:GLU:OE1	2.18	0.43
1:A:466:MET:HE2	1:A:471:VAL:HG22	1.98	0.43
2:B:1:1BU:N1	2:B:1:1BU:CAM	2.81	0.43
1:A:311:ALA:O	1:A:315:LYS:HG3	2.19	0.43
1:B:390:ALA:O	1:B:405:PHE:HE2	2.02	0.43
1:B:487:CYS:SG	1:B:491:LEU:HB3	2.59	0.42
1:B:339:GLU:O	2:B:1:1BU:H2	2.20	0.42
1:B:364:VAL:O	1:B:367:ALA:HB3	2.20	0.42
1:B:362:GLN:O	1:B:366:MET:HG3	2.20	0.41
1:B:268:ARG:O	1:B:284:GLY:HA3	2.21	0.41
1:A:398:LEU:HA	1:A:398:LEU:HD23	1.88	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/286 (90%)	241 (94%)	14 (5%)	2 (1%)	19	39
1	B	256/286 (90%)	227 (89%)	27 (10%)	2 (1%)	19	39
All	All	513/572 (90%)	468 (91%)	41 (8%)	4 (1%)	19	39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	288	GLY
1	B	385	ARG
1	A	385	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	259	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	219/245 (89%)	205 (94%)	14 (6%)	17	35
1	B	224/245 (91%)	204 (91%)	20 (9%)	9	19
All	All	443/490 (90%)	409 (92%)	34 (8%)	13	25

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	291	ARG
1	A	316	LYS
1	A	338	MET
1	A	339	GLU
1	A	351	LYS
1	A	353	GLU
1	A	388	ARG
1	A	426	ILE
1	A	443	SER
1	A	451	LEU
1	A	470	GLU
1	A	476	GLU
1	A	526	GLN
1	A	528	GLN
1	B	268	ARG
1	B	283	MET
1	B	292	VAL
1	B	296	THR
1	B	303	SER
1	B	318	ARG
1	B	323	VAL
1	B	338	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	343	LYS
1	B	345	CYS
1	B	356	LYS
1	B	424	PHE
1	B	426	ILE
1	B	438	ARG
1	B	458	LYS
1	B	469	ARG
1	B	487	CYS
1	B	523	THR
1	B	526	GLN
1	B	528	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	GLN
1	A	397	ASN
1	B	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1BU	B	1	-	39,42,42	1.44	6 (15%)	51,61,61	1.98	11 (21%)
2	1BU	A	1	-	39,42,42	1.41	6 (15%)	51,61,61	2.25	14 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1BU	B	1	-	-	2/16/22/22	0/5/5/5
2	1BU	A	1	-	-	2/16/22/22	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	1BU	CAR-CBF	-3.73	1.34	1.39
2	A	1	1BU	CBG-NAX	-3.30	1.33	1.39
2	B	1	1BU	CBG-NAX	-2.81	1.34	1.39
2	B	1	1BU	CBC-NAW	-2.76	1.36	1.41
2	A	1	1BU	CAR-CBF	-2.75	1.35	1.39
2	B	1	1BU	C6-C5	-2.72	1.41	1.44
2	A	1	1BU	CBC-NAW	-2.46	1.36	1.41
2	A	1	1BU	C6-C5	-2.45	1.41	1.44
2	B	1	1BU	C6-N1	2.43	1.37	1.34
2	B	1	1BU	CBL-CBF	2.42	1.56	1.52
2	A	1	1BU	C6-N1	2.34	1.37	1.34
2	A	1	1BU	CAJ-CBE	-2.15	1.39	1.41

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	1BU	C2-N1-C6	8.25	123.67	116.59
2	B	1	1BU	C2-N1-C6	6.42	122.10	116.59
2	A	1	1BU	C5-C6-N1	-6.03	116.38	121.35
2	B	1	1BU	C5-C6-N1	-5.19	117.08	121.35
2	A	1	1BU	N3-C2-N1	-4.93	120.98	128.68

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	1BU	N3-C2-N1	-4.79	121.19	128.68
2	A	1	1BU	C6-C5-C4	4.44	118.67	115.88
2	B	1	1BU	C6-C5-C4	4.42	118.66	115.88
2	B	1	1BU	C2-N3-C4	3.74	120.56	115.40
2	A	1	1BU	C2-N3-C4	3.44	120.14	115.40
2	A	1	1BU	CAS-C5-C6	-3.38	121.83	124.88
2	B	1	1BU	CAS-C5-C6	-3.29	121.91	124.88
2	A	1	1BU	CBL-CBF-NAV	2.88	123.69	120.59
2	A	1	1BU	CBB-CAS-C5	-2.88	118.40	120.90
2	B	1	1BU	CBL-CBF-NAV	2.75	123.54	120.59
2	A	1	1BU	C5-C6-NAY	2.58	122.57	119.73
2	A	1	1BU	C5-C4-N3	-2.53	120.14	122.83
2	A	1	1BU	NAW-CAZ-NAX	2.40	116.68	112.49
2	A	1	1BU	CBF-CAR-CBG	-2.39	104.12	106.59
2	B	1	1BU	CAR-CBF-NAV	-2.34	108.02	111.41
2	A	1	1BU	CAR-CBF-CBL	-2.29	126.09	129.07
2	B	1	1BU	C5-C4-N3	-2.28	120.40	122.83
2	B	1	1BU	CBB-CAS-C5	-2.07	119.10	120.90
2	A	1	1BU	CAP-C4-C5	2.06	121.49	119.13
2	B	1	1BU	NAY-C6-N1	2.05	121.47	118.72

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	1BU	C5-C6-NAY-CBD
2	A	1	1BU	N1-C6-NAY-CBD
2	B	1	1BU	N1-C6-NAY-CBD
2	B	1	1BU	C5-C6-NAY-CBD


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	1BU	3	0
2	A	1	1BU	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/286 (91%)	0.39	28 (10%) 6 3	2, 14, 49, 61	0
1	B	260/286 (90%)	0.28	18 (6%) 16 12	3, 13, 37, 48	0
All	All	521/572 (91%)	0.33	46 (8%) 10 7	2, 14, 46, 61	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	267	LEU	5.9
1	A	300	GLY	4.7
1	A	328	VAL	4.6
1	A	299	PRO	4.6
1	A	279	GLY	4.3
1	A	332	GLU	3.7
1	B	277	CYS	3.5
1	A	335	TYR	3.5
1	B	300	GLY	3.5
1	A	301	THR	3.5
1	B	258	ASP	3.5
1	A	269	LEU	3.3
1	A	334	ILE	3.2
1	B	473	ASP	3.1
1	A	306	ALA	3.1
1	A	333	PRO	3.0
1	A	327	ALA	3.0
1	B	335	TYR	3.0
1	B	259	ALA	2.9
1	B	305	GLU	2.9
1	A	330	SER	2.9
1	A	302	MET	2.8
1	A	331	GLU	2.8
1	B	301	THR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	268	ARG	2.7
1	A	278	PHE	2.6
1	A	280	GLU	2.6
1	B	290	THR	2.5
1	A	282	TRP	2.5
1	A	285	THR	2.5
1	A	296	THR	2.4
1	B	299	PRO	2.4
1	B	304	PRO	2.3
1	A	277	CYS	2.3
1	B	267	LEU	2.3
1	B	327	ALA	2.3
1	A	305	GLU	2.3
1	B	265	GLU	2.2
1	B	286	TRP	2.2
1	B	333	PRO	2.2
1	B	269	LEU	2.2
1	A	329	VAL	2.1
1	A	271	VAL	2.1
1	A	308	LEU	2.1
1	B	278	PHE	2.1
1	A	304	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	1BU	A	1	38/38	0.95	0.18	13,21,28,28	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	1BU	B	1	38/38	0.96	0.17	17,23,25,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.