



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 07:17 AM EDT

PDB ID : 3H7F
Title : Crystal structure of serine hydroxymethyltransferase from Mycobacterium tuberculosis
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2009-04-24
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2936 (1.50-1.50)
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	447	
1	B	447	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 7350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	414	3117	1970	547	590	1	9	0	13	0
1	B	428	3270	2062	581	614	1	12	0	16	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP O53441
A	-19	ALA	-	expression tag	UNP O53441
A	-18	HIS	-	expression tag	UNP O53441
A	-17	HIS	-	expression tag	UNP O53441
A	-16	HIS	-	expression tag	UNP O53441
A	-15	HIS	-	expression tag	UNP O53441
A	-14	HIS	-	expression tag	UNP O53441
A	-13	HIS	-	expression tag	UNP O53441
A	-12	MET	-	expression tag	UNP O53441
A	-11	GLY	-	expression tag	UNP O53441
A	-10	THR	-	expression tag	UNP O53441
A	-9	LEU	-	expression tag	UNP O53441
A	-8	GLU	-	expression tag	UNP O53441
A	-7	ALA	-	expression tag	UNP O53441
A	-6	GLN	-	expression tag	UNP O53441
A	-5	THR	-	expression tag	UNP O53441
A	-4	GLN	-	expression tag	UNP O53441
A	-3	GLY	-	expression tag	UNP O53441
A	-2	PRO	-	expression tag	UNP O53441
A	-1	GLY	-	expression tag	UNP O53441
A	0	SER	-	expression tag	UNP O53441
B	-20	MET	-	expression tag	UNP O53441
B	-19	ALA	-	expression tag	UNP O53441
B	-18	HIS	-	expression tag	UNP O53441
B	-17	HIS	-	expression tag	UNP O53441

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP O53441
B	-15	HIS	-	expression tag	UNP O53441
B	-14	HIS	-	expression tag	UNP O53441
B	-13	HIS	-	expression tag	UNP O53441
B	-12	MET	-	expression tag	UNP O53441
B	-11	GLY	-	expression tag	UNP O53441
B	-10	THR	-	expression tag	UNP O53441
B	-9	LEU	-	expression tag	UNP O53441
B	-8	GLU	-	expression tag	UNP O53441
B	-7	ALA	-	expression tag	UNP O53441
B	-6	GLN	-	expression tag	UNP O53441
B	-5	THR	-	expression tag	UNP O53441
B	-4	GLN	-	expression tag	UNP O53441
B	-3	GLY	-	expression tag	UNP O53441
B	-2	PRO	-	expression tag	UNP O53441
B	-1	GLY	-	expression tag	UNP O53441
B	0	SER	-	expression tag	UNP O53441

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	418	Total O 418 418	0	0
3	B	544	Total O 544 544	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase 1

Chain A: 



- Molecule 1: Serine hydroxymethyltransferase 1

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.73Å 102.01Å 132.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.50 19.73 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (20.00-1.50) 99.4 (19.73-1.50)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.5.0088	Depositor
R, R_{free}	0.165 , 0.196 0.175 , 0.206	Depositor DCC
R_{free} test set	6606 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtrriage
Anisotropy	0.066	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7350	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	0/3188	0.78	1/4331 (0.0%)
1	B	1.08	0/3351	0.80	2/4550 (0.0%)
All	All	1.04	0/6539	0.79	3/8881 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	372	ARG	NE-CZ-NH2	-5.68	117.46	120.30
1	B	372	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	A	117	ASP	N-CA-CB	-5.39	100.89	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3117	0	3113	7	1
1	B	3270	0	3305	6	0
2	B	1	0	0	0	0
3	A	418	0	0	0	1
3	B	544	0	0	2	0
All	All	7350	0	6418	13	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:406[B]:THR:HG21	3:B:569:HOH:O	1.93	0.67
1:A:379:PHE:HA	1:A:382:VAL:HG22	1.79	0.64
1:A:106:LEU:O	1:A:166:LYS:HE2	1.97	0.64
1:A:111:GLU:OE1	1:A:166:LYS:HD3	2.00	0.60
1:B:406[B]:THR:OG1	1:B:410:ARG:NH1	2.34	0.60
1:A:120[A]:ASN:HD22	1:A:144:VAL:HG22	1.73	0.52
1:B:402:LYS:O	1:B:406[B]:THR:HG22	2.12	0.49
1:B:117[A]:ASP:OD1	3:B:951:HOH:O	2.18	0.49
1:B:227:LLP:NZ	1:B:227:LLP:O3	2.46	0.48
1:A:185:ARG:HD2	1:A:217:HIS:O	2.14	0.47
1:A:111:GLU:OE1	1:A:166:LYS:NZ	2.38	0.46
1:A:227:LLP:O3	1:A:227:LLP:NZ	2.48	0.45
1:B:379:PHE:HA	1:B:382[B]:VAL:HG22	2.01	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:360:SER:OG	3:A:731:HOH:O[3_555]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	422/447 (94%)	411 (97%)	11 (3%)	0	100	100
1	B	441/447 (99%)	429 (97%)	12 (3%)	0	100	100
All	All	863/894 (96%)	840 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	315/338 (93%)	314 (100%)	1 (0%)	92	85
1	B	337/338 (100%)	336 (100%)	1 (0%)	92	85
All	All	652/676 (96%)	650 (100%)	2 (0%)	92	85

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	52	TYR
1	B	52	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	GLN
1	A	99	ASN
1	A	242	GLN
1	B	44	GLN
1	B	347	ASN
1	B	351	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	A	227	1	23,24,25	1.85	5 (21%)	25,32,34	2.00	6 (24%)
1	LLP	B	227	1	23,24,25	1.73	5 (21%)	25,32,34	1.91	6 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	227	1	-	2/16/17/19	0/1/1/1
1	LLP	B	227	1	-	2/16/17/19	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	227	LLP	O3-C3	-4.53	1.26	1.37
1	A	227	LLP	CE-NZ	3.74	1.55	1.46
1	B	227	LLP	C4'-NZ	3.38	1.38	1.27
1	B	227	LLP	C4-C4'	3.36	1.53	1.46
1	A	227	LLP	C4'-NZ	3.26	1.38	1.27
1	B	227	LLP	C6-N1	3.17	1.41	1.34
1	A	227	LLP	C4-C4'	2.93	1.52	1.46
1	B	227	LLP	O3-C3	-2.90	1.30	1.37
1	A	227	LLP	C6-N1	2.24	1.39	1.34
1	B	227	LLP	C3-C2	-2.02	1.38	1.40

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	227	LLP	OP4-C5'-C5	6.39	121.54	109.35
1	A	227	LLP	OP4-C5'-C5	5.98	120.74	109.35
1	A	227	LLP	C5'-C5-C6	-3.84	113.05	119.37
1	A	227	LLP	C2'-C2-C3	3.46	125.16	120.89
1	B	227	LLP	CD-CE-NZ	3.13	118.60	110.93
1	A	227	LLP	CD-CE-NZ	2.69	117.51	110.93

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	227	LLP	C4-C4'-NZ	-2.66	112.12	124.31
1	B	227	LLP	C4-C4'-NZ	-2.62	112.27	124.31
1	B	227	LLP	OP3-P-OP2	2.34	116.57	107.64
1	B	227	LLP	OP2-P-OP4	-2.18	100.92	106.73
1	A	227	LLP	OP2-P-OP1	2.11	118.93	110.68
1	B	227	LLP	O3-C3-C2	2.07	122.01	117.49

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	227	LLP	C4-C4'-NZ-CE
1	A	227	LLP	O-C-CA-CB
1	B	227	LLP	C4-C4'-NZ-CE
1	B	227	LLP	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	227	LLP	1	0
1	B	227	LLP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	413/447 (92%)	0.04	16 (3%) 39 44	12, 21, 34, 48	0
1	B	427/447 (95%)	-0.25	6 (1%) 75 79	11, 18, 27, 41	0
All	All	840/894 (93%)	-0.11	22 (2%) 56 61	11, 20, 31, 48	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	395	SER	5.9
1	B	-1	GLY	5.1
1	A	394	SER	5.0
1	A	361	GLY	4.6
1	A	360	SER	4.1
1	A	-1	GLY	4.0
1	A	393	GLY	4.0
1	A	346	ARG	3.8
1	B	130	LEU	3.3
1	A	300	PRO	3.0
1	A	299	ALA	3.0
1	B	351	ASN	2.7
1	A	390	LEU	2.4
1	B	61	TYR	2.4
1	B	426	ARG	2.4
1	B	425	GLY	2.4
1	A	392	THR	2.4
1	A	425	GLY	2.3
1	A	410	ARG	2.3
1	A	403	ASP	2.2
1	A	396	VAL	2.1
1	A	377	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	A	227	24/25	0.96	0.06	14,17,21,26	0
1	LLP	B	227	24/25	0.96	0.07	12,14,18,22	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	B	427	1/1	0.97	0.37	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.