



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 10:38 PM EDT

PDB ID : 3MKC  
Title : Crystal structure of a putative racemase  
Authors : Eswaramoorthy, S.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2010-04-14  
Resolution : 1.77 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	9185 (1.80-1.76)
Clashscore	141614	10184 (1.80-1.76)
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)
RSRZ outliers	127900	9032 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	394	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3115 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	380	2963	1898	494	558	4	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	387	GLU	-	expression tag	UNP B6R2Z6
A	388	GLY	-	expression tag	UNP B6R2Z6
A	389	HIS	-	expression tag	UNP B6R2Z6
A	390	HIS	-	expression tag	UNP B6R2Z6
A	391	HIS	-	expression tag	UNP B6R2Z6
A	392	HIS	-	expression tag	UNP B6R2Z6
A	393	HIS	-	expression tag	UNP B6R2Z6
A	394	HIS	-	expression tag	UNP B6R2Z6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	152	Total	O	0	0
			152	152		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: racemase



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.84Å 134.84Å 94.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.77 34.76 – 1.77	Depositor EDS
% Data completeness (in resolution range)	95.0 (50.00-1.77) 94.9 (34.76-1.77)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	7.71 (at 1.77Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.216 , 0.241 0.216 , 0.241	Depositor DCC
$R_{free}$ test set	1612 reflections (3.80%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtrriage
Anisotropy	0.260	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3115	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/3022	0.59	0/4098

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2963	0	2900	33	0
2	A	152	0	0	0	0
All	All	3115	0	2900	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:PRO:HG2	1:A:128:GLN:HG3	1.69	0.73
1:A:341:VAL:HG13	1:A:351:ARG:HE	1.61	0.66
1:A:136:ASP:O	1:A:137:LYS:HG2	2.01	0.59
1:A:287:ASP:OD2	1:A:314:HIS:HD2	1.85	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:LEU:CD2	1:A:274:GLU:HG2	2.33	0.58
1:A:81:ILE:HD13	1:A:122:LEU:HD11	1.88	0.54
1:A:186:TRP:HA	1:A:190:GLU:OE1	2.09	0.53
1:A:209:MSE:HG3	1:A:235:TYR:HB3	1.93	0.50
1:A:169:LYS:HD3	1:A:202:LEU:O	2.13	0.49
1:A:196:ARG:HH11	1:A:196:ARG:HG2	1.78	0.48
1:A:16:GLU:HG2	1:A:18:ILE:HG13	1.95	0.48
1:A:288:TYR:CE2	1:A:313:PRO:HB2	2.49	0.48
1:A:361:LEU:HD13	1:A:366:PHE:CZ	2.49	0.48
1:A:22:TYR:CZ	1:A:349:GLU:HG2	2.50	0.47
1:A:287:ASP:HB3	1:A:290:ARG:HB2	1.97	0.47
1:A:95:GLN:HA	1:A:99:MSE:HG3	1.97	0.47
1:A:341:VAL:HG13	1:A:341:VAL:O	2.14	0.47
1:A:317:LYS:HB3	1:A:318:THR:H	1.53	0.47
1:A:196:ARG:HG2	1:A:196:ARG:NH1	2.30	0.46
1:A:288:TYR:CZ	1:A:313:PRO:HB2	2.51	0.45
1:A:222:VAL:HB	1:A:252:LEU:HD21	1.97	0.45
1:A:141:TYR:CD2	1:A:209:MSE:HE2	2.52	0.45
1:A:330:PHE:CE1	1:A:368:LYS:HG2	2.51	0.45
1:A:246:LEU:HD23	1:A:274:GLU:HG2	2.00	0.44
1:A:262:GLY:O	1:A:263:ALA:HB3	2.18	0.43
1:A:19:PRO:HG2	1:A:56:LEU:HD22	2.00	0.43
1:A:53:ASP:OD1	1:A:53:ASP:N	2.52	0.43
1:A:287:ASP:OD2	1:A:314:HIS:CD2	2.69	0.43
1:A:236:PHE:C	1:A:236:PHE:CD1	2.91	0.42
1:A:82:GLU:O	1:A:86:ILE:HG13	2.21	0.41
1:A:189:LYS:HD3	1:A:189:LYS:HA	1.84	0.41
1:A:236:PHE:C	1:A:236:PHE:HD1	2.25	0.40
1:A:341:VAL:HG13	1:A:351:ARG:NE	2.31	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/394 (95%)	360 (96%)	14 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/313 (98%)	302 (99%)	4 (1%)	69	59

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	153	SER
1	A	155	ASP
1	A	209	MSE
1	A	236	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	GLN
1	A	84	ASN
1	A	314	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	371/394 (94%)	0.50	26 (7%) <b>16</b> <b>16</b>	9, 19, 31, 37	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	PRO	11.1
1	A	147	ALA	6.7
1	A	152	ALA	5.8
1	A	22	TYR	5.8
1	A	148	ILE	5.5
1	A	384	ALA	5.5
1	A	386	LEU	4.8
1	A	185	ASP	4.8
1	A	154	LEU	4.8
1	A	156	VAL	4.0
1	A	145	TYR	3.9
1	A	28	SER	3.8
1	A	316	TRP	3.5
1	A	155	ASP	3.4
1	A	157	ALA	3.1
1	A	136	ASP	2.9
1	A	21	ASN	2.8
1	A	183	LYS	2.8
1	A	23	GLN	2.8
1	A	193	TYR	2.6
1	A	182	ILE	2.5
1	A	380	GLN	2.4
1	A	151	ASP	2.3
1	A	346	CYS	2.2
1	A	353	TYR	2.1
1	A	186	TRP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.