



# Full wwPDB X-ray Structure Validation Report i

Aug 28, 2023 – 11:46 PM EDT

PDB ID : 3PSC  
Title : Bovine GRK2 in complex with Gbetagamma subunits  
Authors : Thal, D.M.; Tesmer, J.J.  
Deposited on : 2010-12-01  
Resolution : 2.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3863 (2.70-2.66)
Clashscore	141614	4210 (2.70-2.66)
Ramachandran outliers	138981	4141 (2.70-2.66)
Sidechain outliers	138945	4141 (2.70-2.66)
RSRZ outliers	127900	3780 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-adrenergic receptor kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	614	Total	C 5037	N 3216	O 876	S 909	36	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	670	ALA	SER	engineered mutation	UNP P21146
A	690	HIS	-	expression tag	UNP P21146
A	691	HIS	-	expression tag	UNP P21146
A	692	HIS	-	expression tag	UNP P21146
A	693	HIS	-	expression tag	UNP P21146
A	694	HIS	-	expression tag	UNP P21146
A	695	HIS	-	expression tag	UNP P21146

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	339	Total	C 2607	N 1607	O 468	S 511	21	0	0

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	61	Total	C 481	N 305	O 83	S 89	4	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-5	HIS	-	expression tag	UNP P63212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	-4	HIS	-	expression tag	UNP P63212
G	-3	HIS	-	expression tag	UNP P63212
G	-2	HIS	-	expression tag	UNP P63212
G	-1	HIS	-	expression tag	UNP P63212
G	0	HIS	-	expression tag	UNP P63212


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	B	3	Total O 3 3	0	0
4	G	1	Total O 1 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-adrenergic receptor kinase 1



- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	187.93Å    73.15Å    122.78Å 90.00°    115.42°    90.00°	Depositor
Resolution (Å)	30.00 – 2.67 29.56 – 2.67	Depositor EDS
% Data completeness (in resolution range)	99.5 (30.00-2.67) 99.5 (29.56-2.67)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	3.12 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
$R$ , $R_{free}$	0.223 , 0.257 0.220 , 0.246	Depositor DCC
$R_{free}$ test set	2165 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	78.0	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 49.9	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8133	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/5150	0.59	0/6918
2	B	0.50	0/2654	0.63	0/3597
3	G	0.49	0/481	0.58	0/646
All	All	0.47	0/8285	0.61	0/11161

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	G	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	G	67	PHE	Mainchain

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5037	0	5042	56	0
2	B	2607	0	2510	41	0
3	G	481	0	493	7	0
4	A	4	0	0	1	0
4	B	3	0	0	0	0
4	G	1	0	0	0	0
All	All	8133	0	8045	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:MET:CE	1:A:381:LEU:HD13	1.72	1.18
1:A:294:MET:HE2	1:A:381:LEU:HD13	1.49	0.95
2:B:33:ILE:HD12	3:G:34:ALA:HB3	1.44	0.94
2:B:86:THR:O	2:B:87:THR:HB	1.69	0.92
2:B:340:ASN:HD21	3:G:59:ASN:HD21	1.26	0.83
1:A:294:MET:HE3	1:A:381:LEU:HD13	1.60	0.80
1:A:63:LEU:HD22	1:A:526:PHE:CE1	2.26	0.70
1:A:609:VAL:HG22	1:A:622:LEU:HD22	1.76	0.66
1:A:338:LEU:HD23	1:A:351:VAL:HG13	1.76	0.66
2:B:128:THR:HG22	2:B:130:GLU:H	1.62	0.65
2:B:61:MET:CE	2:B:70:LEU:HD13	2.27	0.64
1:A:301:ILE:HD11	1:A:323:ILE:HD13	1.82	0.61
1:A:62:LYS:NZ	1:A:519:GLN:OE1	2.34	0.61
1:A:105:SER:HB2	1:A:136:LEU:HD21	1.83	0.60
2:B:33:ILE:HD11	3:G:31:SER:HA	1.85	0.59
1:A:300:GLU:OE1	1:A:331:VAL:HG22	2.03	0.58
1:A:246:VAL:HG11	1:A:254:ILE:HG21	1.85	0.58
2:B:71:VAL:HG23	2:B:105:TYR:CD2	2.39	0.57
2:B:233:CYS:HB2	2:B:276:VAL:HG23	1.87	0.56
2:B:71:VAL:CG2	2:B:105:TYR:CD2	2.87	0.56
1:A:40:ARG:NH2	1:A:166:ILE:O	2.36	0.56
1:A:412:GLU:C	1:A:413:LEU:HD23	2.26	0.56
2:B:33:ILE:CD1	3:G:34:ALA:HB3	2.29	0.55
1:A:314:VAL:HG22	1:A:370:SER:HA	1.89	0.55
2:B:58:ILE:HD13	2:B:336:LEU:CD1	2.37	0.55
2:B:61:MET:HE3	2:B:70:LEU:HD13	1.89	0.54
1:A:380:MET:CE	1:A:381:LEU:HD23	2.36	0.54
2:B:232:ILE:HG13	2:B:243:THR:HG22	1.88	0.54

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:413:LEU:HD12	1:A:422:ARG:HG3	1.89	0.54
2:B:273:ILE:N	2:B:273:ILE:HD12	2.24	0.53
1:A:427:GLY:HA3	1:A:437:LEU:HD12	1.91	0.53
1:A:334:SER:O	1:A:336:LEU:HD12	2.09	0.53
1:A:391:PHE:CD1	1:A:406:THR:HA	2.44	0.53
1:A:529:ILE:O	1:A:533:THR:HG23	2.09	0.52
1:A:314:VAL:HG23	1:A:342:PHE:CD2	2.46	0.50
1:A:601:LEU:HD21	1:A:624:ILE:HD11	1.93	0.50
2:B:33:ILE:HD12	3:G:34:ALA:CB	2.30	0.50
2:B:205:ASP:O	2:B:206:ALA:HB3	2.11	0.50
2:B:301:LYS:O	2:B:302:ALA:HB3	2.12	0.50
1:A:173:ARG:HA	1:A:176:GLN:OE1	2.12	0.49
1:A:273:LEU:HD12	1:A:274:MET:H	1.78	0.48
1:A:294:MET:HE2	1:A:381:LEU:CD1	2.35	0.48
2:B:163:ASP:C	2:B:164:THR:HG23	2.34	0.48
2:B:340:ASN:HD21	3:G:59:ASN:ND2	2.04	0.48
1:A:405:MET:HB3	1:A:409:MET:HE2	1.96	0.47
2:B:247:ASP:O	2:B:248:ALA:HB3	2.14	0.47
1:A:380:MET:HE2	1:A:381:LEU:HD23	1.95	0.47
1:A:111:THR:O	1:A:115:LYS:HB3	2.14	0.47
2:B:262:MET:SD	2:B:302:ALA:HB2	2.54	0.47
1:A:155:GLN:OE1	1:A:158:ARG:NH2	2.46	0.46
1:A:390:PRO:CB	1:A:411:VAL:HG21	2.45	0.46
1:A:301:ILE:CD1	1:A:323:ILE:HD13	2.44	0.46
2:B:128:THR:HG22	2:B:130:GLU:N	2.29	0.46
1:A:533:THR:HA	1:A:536:LEU:HD12	1.96	0.46
2:B:63:TRP:CD2	2:B:321:THR:HG22	2.51	0.46
1:A:274:MET:HE1	1:A:332:ARG:HB2	1.97	0.46
1:A:257:MET:HB3	1:A:257:MET:HE2	1.71	0.46
1:A:197:ILE:O	1:A:197:ILE:HG22	2.16	0.46
2:B:179:THR:HG22	2:B:181:THR:HG23	1.97	0.46
1:A:113:ILE:O	1:A:117:LEU:HB2	2.16	0.45
2:B:51:LEU:HD13	2:B:82:TRP:CD2	2.52	0.45
1:A:565:MET:HE3	1:A:634:CYS:SG	2.56	0.45
2:B:79:LEU:HD11	2:B:114:CYS:HB3	1.98	0.45
1:A:246:VAL:CG1	1:A:254:ILE:HG21	2.47	0.45
2:B:163:ASP:O	2:B:164:THR:HG23	2.16	0.45
1:A:397:LYS:O	1:A:399:LYS:N	2.50	0.44
2:B:58:ILE:HD13	2:B:336:LEU:HD11	1.99	0.44
1:A:380:MET:HE1	1:A:381:LEU:CD2	2.48	0.44
1:A:319:LYS:HA	1:A:380:MET:HG3	1.99	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:45:MET:CE	2:B:308:LEU:HD21	2.48	0.43
2:B:63:TRP:CZ2	2:B:328:ALA:HB2	2.53	0.43
1:A:390:PRO:HB3	1:A:411:VAL:HG21	2.00	0.43
2:B:70:LEU:HD11	2:B:336:LEU:HD22	2.00	0.43
1:A:472:PRO:HA	1:A:473:PRO:HD3	1.84	0.43
1:A:509:PHE:N	1:A:510:PRO:CD	2.82	0.43
2:B:210:LEU:HD22	2:B:255:LEU:HD22	2.00	0.42
2:B:262:MET:HG2	2:B:264:TYR:CZ	2.54	0.42
1:A:467:PRO:HA	1:A:468:PRO:HD3	1.93	0.42
1:A:380:MET:HE1	1:A:381:LEU:HD23	1.99	0.42
2:B:45:MET:HE3	2:B:308:LEU:HD21	2.01	0.42
1:A:565:MET:CE	1:A:634:CYS:SG	3.08	0.42
2:B:249:THR:HG22	2:B:265:SER:HB3	2.01	0.42
1:A:464:GLN:NE2	4:A:692:HOH:O	2.53	0.41
2:B:58:ILE:HD13	2:B:336:LEU:HD12	2.02	0.41
1:A:294:MET:CE	1:A:298:ALA:HB2	2.51	0.41
2:B:165:THR:HG22	2:B:181:THR:HG22	2.03	0.41
1:A:294:MET:HE3	1:A:381:LEU:CD1	2.41	0.41
2:B:58:ILE:CD1	2:B:336:LEU:HD12	2.51	0.41
1:A:544:ASN:HB3	1:A:549:HIS:CB	2.50	0.40
2:B:231:ALA:CB	2:B:275:SER:HA	2.51	0.40
1:A:391:PHE:CE1	1:A:406:THR:HA	2.55	0.40
1:A:274:MET:SD	1:A:332:ARG:HD2	2.61	0.40
1:A:314:VAL:HG23	1:A:342:PHE:CE2	2.57	0.40
2:B:73:ALA:HB2	2:B:103:CYS:HB3	2.02	0.40
2:B:170:ASP:HB3	2:B:173:THR:HB	2.04	0.40
3:G:28:ILE:HG22	3:G:29:LYS:N	2.35	0.40
1:A:474:ARG:HA	1:A:474:ARG:CZ	2.51	0.40
1:A:600:LEU:HD12	1:A:600:LEU:C	2.42	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	608/695 (88%)	573 (94%)	33 (5%)	2 (0%)	41 64
2	B	337/340 (99%)	312 (93%)	24 (7%)	1 (0%)	41 64
3	G	59/74 (80%)	56 (95%)	3 (5%)	0	100 100
All	All	1004/1109 (90%)	941 (94%)	60 (6%)	3 (0%)	41 64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	398	ASP
2	B	53	GLY
1	A	508	ASN

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	553/617 (90%)	536 (97%)	17 (3%)	40 67
2	B	282/283 (100%)	270 (96%)	12 (4%)	29 54
3	G	50/61 (82%)	50 (100%)	0	100 100
All	All	885/961 (92%)	856 (97%)	29 (3%)	38 64

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	ASP
1	A	192	SER
1	A	240	ARG
1	A	262	HIS
1	A	263	THR
1	A	265	ASP
1	A	284	SER
1	A	290	SER
1	A	294	MET
1	A	396	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	398	ASP
1	A	409	MET
1	A	432	ASP
1	A	543	LYS
1	A	618	LYS
1	A	651	TYR
1	A	666	ASN
2	B	40	VAL
2	B	52	ARG
2	B	70	LEU
2	B	87	THR
2	B	105	TYR
2	B	122	SER
2	B	199	PHE
2	B	234	PHE
2	B	292	PHE
2	B	304	ARG
2	B	318	LEU
2	B	340	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	310	ASN
1	A	434	ASN
1	A	459	GLN
1	A	503	GLN
2	B	6	GLN
2	B	132	ASN
2	B	176	GLN
2	B	239	ASN
3	G	59	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CMT	G	68	3	7,7,7	2.14	1 (14%)	6,8,8	2.35	4 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CMT	G	68	3	-	4/8/8/8	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	G	68	CMT	OXT-C	5.51	1.46	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	68	CMT	OXT-C-CA	3.76	121.15	111.52
3	G	68	CMT	C1-OXT-C	2.83	122.35	115.94
3	G	68	CMT	CA-CB-SG	-2.38	109.31	114.44
3	G	68	CMT	OXT-C-O	-2.22	119.51	123.84

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	68	CMT	O-C-CA-N
3	G	68	CMT	OXT-C-CA-N
3	G	68	CMT	O-C-OXT-C1
3	G	68	CMT	CA-C-OXT-C1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	614/695 (88%)	0.47	57 (9%) 8 6	61, 94, 148, 200	0
2	B	339/340 (99%)	0.27	20 (5%) 22 20	50, 76, 126, 231	0
3	G	60/74 (81%)	0.03	1 (1%) 70 71	61, 85, 178, 194	0
All	All	1013/1109 (91%)	0.37	78 (7%) 13 11	50, 88, 148, 231	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	129	ARG	7.5
1	A	475	GLY	7.3
1	A	202	PHE	6.3
2	B	4	LEU	5.6
2	B	130	GLU	5.2
1	A	29	SER	4.8
1	A	497	LYS	4.7
1	A	201	GLY	4.7
1	A	364	LYS	4.6
1	A	229	MET	4.5
1	A	506	TYR	4.5
1	A	550	GLU	4.4
1	A	95	LEU	4.3
1	A	231	GLN	4.2
3	G	9	ILE	4.1
1	A	232	GLY	4.0
1	A	474	ARG	3.9
1	A	365	GLY	3.8
1	A	498	LEU	3.6
1	A	222	LEU	3.6
1	A	235	LEU	3.6
2	B	148	CYS	3.5
2	B	103	CYS	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	2	SER	3.4
1	A	396	THR	3.4
1	A	549	HIS	3.3
1	A	412	GLU	3.3
2	B	8	ARG	3.2
1	A	440	LEU	3.2
2	B	102	THR	3.2
1	A	393	GLN	3.2
1	A	618	LYS	3.2
2	B	5	ASP	3.1
2	B	60	ALA	3.1
1	A	366	VAL	3.0
1	A	376	SER	3.0
2	B	128	THR	3.0
1	A	395	LYS	2.9
1	A	422	ARG	2.9
1	A	502	ASP	2.9
1	A	616	GLU	2.8
1	A	199	ARG	2.7
1	A	499	LEU	2.7
1	A	415	ASP	2.7
1	A	334	SER	2.7
1	A	413	LEU	2.7
2	B	150	ARG	2.6
1	A	230	LYS	2.5
2	B	276	VAL	2.5
1	A	397	LYS	2.5
2	B	42	ARG	2.5
1	A	379	CYS	2.5
1	A	388	HIS	2.4
1	A	345	LYS	2.4
1	A	96	GLU	2.4
1	A	30	LYS	2.4
2	B	149	CYS	2.3
1	A	394	HIS	2.3
1	A	551	GLU	2.3
1	A	615	LYS	2.3
1	A	137	VAL	2.3
1	A	420	GLU	2.2
1	A	298	ALA	2.2
1	A	380	MET	2.2
1	A	382	PHE	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	301	ILE	2.2
1	A	504	GLU	2.2
1	A	652	ARG	2.2
1	A	297	TYR	2.2
2	B	15	LYS	2.2
1	A	356	TYR	2.1
2	B	232	ILE	2.1
1	A	547	LEU	2.1
2	B	189	SER	2.1
2	B	61	MET	2.1
1	A	106	ARG	2.1
2	B	317	CYS	2.0
1	A	294	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CMT	G	68	8/8	0.84	0.16	90,92,98,102	0

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.