



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2024 – 06:43 PM EDT

PDB ID : 3QZW
Title : Plasticity of human CD8 binding to peptide-HLA-A*2402
Authors : Shi, Y.; Qi, J.; Gao, G.F.
Deposited on : 2011-03-07
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 75%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 23%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">75% 23% .</p>
1	D	274	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 70%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 28%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">2% 70% 28% .</p>
2	B	100	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 78%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 5%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">2% 78% 17% 5%</p>
2	E	100	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 74%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 2%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">% 74% 22% . .</p>
3	C	10	<div style="display: flex; align-items: center;"> <div style="width: 50%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 40%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> </div> <p style="text-align: center;">50% 40% 10%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	10	 40% 50% 10%
4	G	114	 % 78% 20% .
4	H	114	 5% 75% 21% . .
4	I	114	 5% 66% 32% .
4	J	114	 18% 61% 36% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, A-24 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2221	1382	403	426	10	0	0	0
1	D	274	2221	1382	403	426	10	0	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	836	533	141	158	4	0	0	0
2	E	100	836	533	141	158	4	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	EXPRESSION TAG	UNP P61769
E	0	MET	-	EXPRESSION TAG	UNP P61769

- Molecule 3 is a protein called 10-mer peptide from Protein Nef.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	10	92	64	14	13	1	0	0	0
3	F	10	92	64	14	13	1	0	0	0

- Molecule 4 is a protein called T-cell surface glycoprotein CD8 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	G	114	906	583	153	166	4	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	114	Total	C	N	O	S	0	0	0
			906	583	153	166	4			
4	I	114	Total	C	N	O	S	0	0	0
			906	583	153	166	4			
4	J	114	Total	C	N	O	S	0	0	0
			906	583	153	166	4			


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	44	Total	O	0	0
			44	44		
5	B	16	Total	O	0	0
			16	16		
5	C	1	Total	O	0	0
			1	1		
5	D	28	Total	O	0	0
			28	28		
5	E	16	Total	O	0	0
			16	16		
5	F	1	Total	O	0	0
			1	1		
5	G	33	Total	O	0	0
			33	33		
5	H	10	Total	O	0	0
			10	10		
5	I	14	Total	O	0	0
			14	14		
5	J	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain



- Molecule 1: HLA class I histocompatibility antigen, A-24 alpha chain



- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin







- Molecule 3: 10-mer peptide from Protein Nef





- Molecule 3: 10-mer peptide from Protein Nef



- Molecule 4: T-cell surface glycoprotein CD8 alpha chain




- Molecule 4: T-cell surface glycoprotein CD8 alpha chain




- Molecule 4: T-cell surface glycoprotein CD8 alpha chain



- Molecule 4: T-cell surface glycoprotein CD8 alpha chain





4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	66.24Å 67.02Å 101.14Å 73.71° 71.84° 81.43°	Depositor
Resolution (Å)	31.61 – 2.80 31.61 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.4 (31.61-2.80) 98.4 (31.61-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.56 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, R_{free}	0.202 , 0.260 0.196 , 0.251	Depositor DCC
R_{free} test set	1928 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtrriage
Anisotropy	0.606	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.010 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10086	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	0/2281	0.46	0/3092
1	D	0.28	0/2281	0.47	1/3092 (0.0%)
2	B	0.30	0/859	0.58	1/1162 (0.1%)
2	E	0.30	0/859	0.56	2/1162 (0.2%)
3	C	0.33	0/97	0.47	0/130
3	F	0.30	0/97	0.47	0/130
4	G	0.32	0/931	0.50	0/1263
4	H	0.32	0/931	0.65	4/1263 (0.3%)
4	I	0.31	0/931	0.47	0/1263
4	J	0.38	1/931 (0.1%)	0.64	3/1263 (0.2%)
All	All	0.31	1/10198 (0.0%)	0.52	11/13820 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	7	PRO	N-CD	8.20	1.59	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	98	ASP	CB-CA-C	9.73	129.85	110.40
4	J	6	SER	CB-CA-C	7.91	125.13	110.10
2	E	2	GLN	N-CA-CB	-7.52	97.06	110.60
4	J	29	PRO	N-CA-C	7.15	130.69	112.10
2	E	2	GLN	N-CA-C	7.13	130.26	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2221	0	2082	52	0
1	D	2221	0	2082	66	0
2	B	836	0	803	28	0
2	E	836	0	803	22	0
3	C	92	0	85	4	0
3	F	92	0	85	8	0
4	G	906	0	885	22	0
4	H	906	0	885	27	0
4	I	906	0	885	35	0
4	J	906	0	885	31	0
5	A	44	0	0	1	0
5	B	16	0	0	1	0
5	C	1	0	0	0	0
5	D	28	0	0	1	0
5	E	16	0	0	0	0
5	F	1	0	0	0	0
5	G	33	0	0	0	0
5	H	10	0	0	2	0
5	I	14	0	0	0	0
5	J	1	0	0	0	0
All	All	10086	0	9480	252	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 252 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:98:ASP:O	2:B:99:MET:HG2	1.29	1.23
4:H:29:PRO:O	4:H:30:THR:HG22	1.03	1.16
2:B:98:ASP:O	2:B:99:MET:CG	1.94	1.16
4:H:29:PRO:O	4:H:30:THR:CG2	1.97	1.11
2:E:1:ILE:HG23	2:E:1:ILE:O	1.56	1.05

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/274 (99%)	256 (94%)	16 (6%)	0	100	100
1	D	272/274 (99%)	250 (92%)	22 (8%)	0	100	100
2	B	98/100 (98%)	91 (93%)	7 (7%)	0	100	100
2	E	98/100 (98%)	93 (95%)	4 (4%)	1 (1%)	15	44
3	C	8/10 (80%)	8 (100%)	0	0	100	100
3	F	8/10 (80%)	8 (100%)	0	0	100	100
4	G	112/114 (98%)	102 (91%)	10 (9%)	0	100	100
4	H	112/114 (98%)	97 (87%)	13 (12%)	2 (2%)	8	28
4	I	112/114 (98%)	100 (89%)	11 (10%)	1 (1%)	17	46
4	J	112/114 (98%)	98 (88%)	13 (12%)	1 (1%)	17	46
All	All	1204/1224 (98%)	1103 (92%)	96 (8%)	5 (0%)	34	66

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	H	29	PRO
4	H	30	THR
4	I	43	ALA
2	E	1	ILE
4	J	29	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/230 (100%)	223 (97%)	7 (3%)	41	75
1	D	230/230 (100%)	222 (96%)	8 (4%)	36	70
2	B	95/95 (100%)	91 (96%)	4 (4%)	30	63
2	E	95/95 (100%)	91 (96%)	4 (4%)	30	63
3	C	9/9 (100%)	7 (78%)	2 (22%)	1	2
3	F	9/9 (100%)	8 (89%)	1 (11%)	6	19
4	G	100/100 (100%)	96 (96%)	4 (4%)	31	65
4	H	100/100 (100%)	98 (98%)	2 (2%)	55	84
4	I	100/100 (100%)	98 (98%)	2 (2%)	55	84
4	J	100/100 (100%)	98 (98%)	2 (2%)	55	84
All	All	1068/1068 (100%)	1032 (97%)	36 (3%)	37	71

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	G	45	SER
4	J	107	PHE
4	G	47	THR
4	I	8	LEU
3	C	4	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	93	HIS
4	H	2	GLN
1	D	174	ASN
4	J	55	ASN
2	E	51	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/274 (100%)	-0.27	2 (0%) 87 84	30, 44, 66, 82	0
1	D	274/274 (100%)	0.03	6 (2%) 62 52	38, 55, 79, 91	0
2	B	100/100 (100%)	-0.16	2 (2%) 65 56	30, 44, 76, 83	0
2	E	100/100 (100%)	-0.32	1 (1%) 82 77	33, 46, 70, 86	0
3	C	10/10 (100%)	-0.23	0 100 100	37, 40, 48, 52	0
3	F	10/10 (100%)	-0.00	0 100 100	44, 47, 57, 61	0
4	G	114/114 (100%)	-0.37	1 (0%) 84 80	30, 43, 67, 80	0
4	H	114/114 (100%)	0.14	6 (5%) 26 17	38, 59, 84, 105	0
4	I	114/114 (100%)	0.19	6 (5%) 26 17	49, 63, 87, 116	0
4	J	114/114 (100%)	0.91	21 (18%) 1 1	61, 84, 110, 117	0
All	All	1224/1224 (100%)	-0.01	45 (3%) 41 31	30, 52, 87, 117	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	30	THR	10.8
4	J	114	ALA	6.6
4	H	114	ALA	5.2
4	J	42	ALA	4.8
2	E	0	MET	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.