

wwPDB X-ray Structure Validation Summary Report (i)

Jun 18, 2024 – 03:01 PM EDT

PDB ID	:	3X28
Title	:	Crystal structure of Nitrile Hydratase mutant bR56K
Authors	:	Yamanaka, Y.; Hashimoto, K.; Noguchi, K.; Yohda, M.; Odaka, M.
Deposited on		
Resolution	:	1.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	207	86%	8% 5%
2	В	212	92%	8%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 3459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Nitrile hydratase subunit alpha.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	197	Total 1547	C 985	N 260	O 296	S 6	0	0	0

• Molecule 2 is a protein called Nitrile hydratase subunit beta.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	В	212	Total 1668	C 1061	N 281	O 316	S 10	0	3	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	56	LYS	ARG	ENGINEERED MUTATION	UNP P13449

• Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total Fe 1 1	0	0

• Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total Mg 1 1	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0



• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	102	Total O 102 102	0	0
6	В	139	Total O 139 139	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Nitrile hydratase subunit alpha





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	114.79Å 60.58Å 82.00Å	Depositor
a, b, c, α , β , γ	90.00° 124.97° 90.00°	Depositor
Resolution (Å)	33.17 - 1.65	Depositor
Resolution (A)	33.17 - 1.65	EDS
% Data completeness	90.4 (33.17-1.65)	Depositor
(in resolution range)	90.4 (33.17-1.65)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$3.94 (at 1.65 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
D D.	0.180 , 0.218	Depositor
R, R_{free}	0.191 , 0.227	DCC
R_{free} test set	2560 reflections $(5.10%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	15.1	Xtriage
Anisotropy	0.046	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.36 , 45.6	EDS
L-test for twinning ²	$ \langle L \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3459	wwPDB-VP
Average B, all atoms $(Å^2)$	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.64% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, CSO, MG, CL, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
	RMSZ	# Z > 5	RMSZ	# Z > 5		
1	А	1.09	4/1570~(0.3%)	1.10	11/2144~(0.5%)	
2	В	1.13	2/1725~(0.1%)	1.12	6/2345~(0.3%)	
All	All	1.11	6/3295~(0.2%)	1.11	17/4489~(0.4%)	

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	А	150	GLU	CD-OE1	-8.95	1.15	1.25
2	В	187	GLU	CD-OE2	7.18	1.33	1.25
2	В	209	GLU	CD-OE2	-6.45	1.18	1.25
1	А	113	SER	CB-OG	5.75	1.49	1.42
1	А	10	ASN	CG-ND2	5.59	1.46	1.32

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	107	ARG	NE-CZ-NH1	11.57	126.09	120.30
2	В	65	ARG	NE-CZ-NH2	-10.20	115.20	120.30
1	А	147	MET	CG-SD-CE	-10.14	83.98	100.20
2	В	130	ARG	NE-CZ-NH2	9.80	125.20	120.30
1	А	133	ARG	NE-CZ-NH2	-8.82	115.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1547	0	1516	4	0
2	В	1668	0	1597	7	0
3	А	1	0	0	0	0
4	В	1	0	0	0	0
5	В	1	0	0	0	0
6	А	102	0	0	1	0
6	В	139	0	0	1	0
All	All	3459	0	3113	9	0

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 9 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:34:HIS:HE1	6:B:483:HOH:O	1.94	0.50
1:A:93:TYR:OH	2:B:155:HIS:HE1	1.98	0.46
1:A:139:GLU:HA	2:B:195:ASP:OD1	2.16	0.45
2:B:166:ALA:HA	2:B:171:ARG:HD3	1.99	0.44
2:B:161:TRP:HB3	2:B:162:PRO:HD2	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	193/207~(93%)	188 (97%)	4 (2%)	1 (0%)	29	11
2	В	213/212~(100%)	208~(98%)	5(2%)	0	100	100
All	All	406/419~(97%)	396~(98%)	9(2%)	1 (0%)	47	28



All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	113	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outlier		Percentiles		
1	А	161/171~(94%)	160 (99%)	1 (1%)	86 76		
2	В	176/173~(102%)	174 (99%)	2(1%)	73 57		
All	All	337/344~(98%)	334 (99%)	3 (1%)	78 66		

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	205	THR
2	В	103	PHE
2	В	165	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	83	GLN
1	А	173	GLN
2	В	21	ASN
2	В	34	HIS
2	В	155	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Chain	Dec	Dec	Dec	Dec	Dec	Dec	Tinle	В	Bond lengths			Bond angles		
IVIOI	туре	Chain	\mathbf{Res}	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2					
1	CSD	А	112	1,3	4,7,8	2.01	1 (25%)	1,8,10	1.95	0					
1	CSO	А	114	1,3	3,6,7	1.97	2 (66%)	1,6,8	0.34	0					

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	А	112	1,3	-	1/2/6/8	-
1	CSO	А	114	1,3	-	0/1/5/7	-

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
1	А	112	CSD	OD1-SG	3.71	1.51	1.47
1	А	114	CSO	CA-N	-2.09	1.42	1.48
1	А	114	CSO	CB-CA	-2.04	1.48	1.53

All (3) bond length outliers are listed below:

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	А	112	CSD	CA-CB-SG-OD1

There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	А	114	CSO	1	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	195/207~(94%)	0.01	8 (4%) 37 37	9, 16, 33, 56	0
2	В	212/212 (100%)	-0.30	1 (0%) 91 92	9, 14, 26, 51	0
All	All	407/419~(97%)	-0.15	9 (2%) 62 63	9, 15, 30, 56	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	205	THR	6.9
1	А	11	ALA	6.1
2	В	212	ALA	4.7
1	А	153	SER	3.9
1	А	13	PRO	3.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q < 0.9
1	CSO	А	114	7/8	0.98	0.14	$11,\!11,\!15,\!17$	0
1	CSD	А	112	8/9	0.99	0.10	9,10,10,12	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
4	MG	В	301	1/1	0.96	0.10	9,9,9,9	1
5	CL	В	302	1/1	0.98	0.05	21,21,21,21	0
3	\mathbf{FE}	А	300	1/1	1.00	0.08	9,9,9,9	0

6.5 Other polymers (i)

There are no such residues in this entry.

