



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 03:38 PM EDT

PDB ID : 4C8U
Title : Xenopus ZNRF3 ectodomain crystal form II
Authors : Zebisch, M.; Jones, E.Y.
Deposited on : 2013-10-01
Resolution : 3.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2399 (3.04-3.00)
Clashscore	141614	2734 (3.04-3.00)
Ramachandran outliers	138981	2640 (3.04-3.00)
Sidechain outliers	138945	2643 (3.04-3.00)
RSRZ outliers	127900	2287 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	182	 56% 21% 20%
1	B	182	 60% 17% 21%

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 2207 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 UBIQUITIN-PROTEIN LIGASE ZNR3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	146	1111	699	193	212	7	0	0	0
1	B	144	1095	689	191	208	7	0	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLU	-	expression tag	UNP Q08D68
A	23	THR	-	expression tag	UNP Q08D68
A	192	GLY	-	expression tag	UNP Q08D68
A	193	THR	-	expression tag	UNP Q08D68
A	194	HIS	-	expression tag	UNP Q08D68
A	195	HIS	-	expression tag	UNP Q08D68
A	196	HIS	-	expression tag	UNP Q08D68
A	197	HIS	-	expression tag	UNP Q08D68
A	198	HIS	-	expression tag	UNP Q08D68
A	199	HIS	-	expression tag	UNP Q08D68
A	200	HIS	-	expression tag	UNP Q08D68
A	201	HIS	-	expression tag	UNP Q08D68
A	202	HIS	-	expression tag	UNP Q08D68
A	203	HIS	-	expression tag	UNP Q08D68
B	22	GLU	-	expression tag	UNP Q08D68
B	23	THR	-	expression tag	UNP Q08D68
B	192	GLY	-	expression tag	UNP Q08D68
B	193	THR	-	expression tag	UNP Q08D68
B	194	HIS	-	expression tag	UNP Q08D68
B	195	HIS	-	expression tag	UNP Q08D68
B	196	HIS	-	expression tag	UNP Q08D68
B	197	HIS	-	expression tag	UNP Q08D68
B	198	HIS	-	expression tag	UNP Q08D68
B	199	HIS	-	expression tag	UNP Q08D68
B	200	HIS	-	expression tag	UNP Q08D68

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	201	HIS	-	expression tag	UNP Q08D68
B	202	HIS	-	expression tag	UNP Q08D68
B	203	HIS	-	expression tag	UNP Q08D68

- Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Br 1 1	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.39Å 67.41Å 75.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.34 – 3.01 46.23 – 3.01	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.34-3.01) 99.4 (46.23-3.01)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.99 (at 3.01Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.281 , 0.351 0.277 , 0.358	Depositor DCC
R_{free} test set	291 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	48.2	Xtrriage
Anisotropy	0.717	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 69.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	2207	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 52.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.8690e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/1130	0.72	0/1531
1	B	0.60	0/1114	0.77	0/1509
All	All	0.57	0/2244	0.74	0/3040

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1111	0	1089	33	0
1	B	1095	0	1074	25	0
2	A	1	0	0	0	0
All	All	2207	0	2163	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:164:LEU:O	1:A:168:VAL:HG23	1.55	1.06
1:A:164:LEU:C	1:A:168:VAL:HG23	2.14	0.67
1:B:79:LEU:O	1:B:80:CYS:SG	2.53	0.67
1:A:71:ILE:HD11	1:A:176:ALA:HB2	1.78	0.65
1:A:101:GLN:HG3	1:A:103:GLU:OE1	1.97	0.64
1:B:110:LEU:HD22	1:B:114:GLY:HA3	1.82	0.62
1:B:68:GLU:HA	1:B:176:ALA:O	2.00	0.61
1:A:125:THR:OG1	1:B:177:ARG:NH2	2.24	0.60
1:A:92:GLY:HA3	1:B:175:ARG:NE	2.16	0.60
1:A:68:GLU:HA	1:A:176:ALA:O	2.02	0.59
1:A:116:ALA:O	1:A:120:VAL:HG23	2.01	0.59
1:B:92:GLY:HA2	1:B:125:THR:HB	1.86	0.57
1:B:52:THR:HG21	1:B:167:ILE:HD11	1.86	0.56
1:B:125:THR:O	1:B:151:ARG:NE	2.38	0.56
1:B:96:VAL:HA	1:B:128:ILE:O	2.06	0.55
1:A:42:SER:HB3	1:A:43:PRO:HD2	1.89	0.54
1:A:44:ASN:HD22	1:A:44:ASN:H	1.55	0.53
1:B:57:ARG:HB2	1:B:155:TYR:CZ	2.44	0.53
1:A:92:GLY:O	1:B:175:ARG:NH2	2.42	0.53
1:A:117:LYS:HG3	1:A:149:LEU:HD21	1.90	0.52
1:A:126:ALA:O	1:A:127:VAL:HG23	2.09	0.52
1:A:92:GLY:HA2	1:A:125:THR:HB	1.93	0.51
1:B:128:ILE:HD11	1:B:178:ILE:HD11	1.92	0.51
1:A:126:ALA:O	1:A:127:VAL:CG2	2.59	0.50
1:A:68:GLU:O	1:B:68:GLU:OE2	2.28	0.50
1:B:112:VAL:HG22	1:B:141:LEU:HD21	1.93	0.50
1:A:143:GLN:HA	1:A:143:GLN:OE1	2.13	0.49
1:A:94:VAL:HG22	1:A:126:ALA:HB3	1.95	0.48
1:B:68:GLU:OE1	1:B:177:ARG:HG2	2.13	0.48
1:A:93:TRP:CD1	1:A:93:TRP:N	2.81	0.47
1:B:131:VAL:HG22	1:B:156:MET:O	2.14	0.47
1:A:57:ARG:HB2	1:A:155:TYR:CZ	2.50	0.47
1:A:47:TYR:OH	1:B:89:TYR:O	2.32	0.46
1:A:164:LEU:O	1:A:168:VAL:CG2	2.45	0.46
1:B:34:VAL:HG21	1:B:156:MET:CE	2.46	0.46
1:A:64:THR:O	1:A:64:THR:HG22	2.16	0.45
1:A:92:GLY:HA3	1:B:175:ARG:HE	1.81	0.45
1:A:98:LYS:HA	1:A:130:ASP:HB3	1.98	0.45
1:A:177:ARG:CZ	1:B:151:ARG:NH1	2.81	0.44
1:B:57:ARG:HB2	1:B:155:TYR:CE1	2.53	0.43
1:A:44:ASN:H	1:A:44:ASN:ND2	2.16	0.43
1:B:91:TYR:CD1	1:B:123:GLY:HA2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:THR:HG21	1:B:68:GLU:OE2	2.19	0.43
1:A:126:ALA:C	1:A:127:VAL:HG23	2.40	0.42
1:B:125:THR:O	1:B:151:ARG:NH2	2.50	0.42
1:A:179:GLN:HG2	1:A:180:HIS:N	2.35	0.42
1:A:163:LYS:HA	1:A:166:ASN:HB2	2.02	0.41
1:B:96:VAL:HG22	1:B:128:ILE:HB	2.03	0.41
1:A:75:HIS:O	1:A:78:GLY:N	2.47	0.40
1:A:62:GLY:HA3	1:A:153:VAL:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/182 (78%)	127 (89%)	14 (10%)	1 (1%)	22	59
1	B	140/182 (77%)	121 (86%)	19 (14%)	0	100	100
All	All	282/364 (78%)	248 (88%)	33 (12%)	1 (0%)	34	71

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	127	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/155 (76%)	105 (90%)	12 (10%)	7	26
1	B	115/155 (74%)	104 (90%)	11 (10%)	8	30
All	All	232/310 (75%)	209 (90%)	23 (10%)	8	28

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	52	THR
1	A	57	ARG
1	A	80	CYS
1	A	88	LEU
1	A	113	LEU
1	A	133	ASP
1	A	139	GLU
1	A	140	GLN
1	A	165	MET
1	A	166	ASN
1	A	180	HIS
1	B	65	ILE
1	B	68	GLU
1	B	70	GLU
1	B	74	MET
1	B	90	GLU
1	B	109	CYS
1	B	113	LEU
1	B	125	THR
1	B	136	ASP
1	B	166	ASN
1	B	175	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	140	GLN
1	A	166	ASN
1	B	44	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	146/182 (80%)	-0.17	1 (0%) 87 68	28, 65, 107, 127	0
1	B	144/182 (79%)	-0.16	0 100 100	16, 63, 102, 116	0
All	All	290/364 (79%)	-0.16	1 (0%) 94 83	16, 64, 104, 127	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	145	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BR	A	1182	1/1	0.96	0.08	53,53,53,53	1

6.5 Other polymers [i](#)

There are no such residues in this entry.