



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 19, 2024 – 04:46 AM EDT

PDB ID : 4H2Y
Title : Crystal structure of engineered Bradyrhizobium japonicum glycine:[carrier protein] ligase complexed with carrier protein from Agrobacterium tumefaciens and ATP
Authors : Luic, M.; Weygand-Durasevic, I.; Ivic, N.; Mocibob, M.
Deposited on : 2012-09-13
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	 82% 15%
1	B	346	 78% 8% 14%
2	C	103	 45% 59% 16% 25%
2	D	103	 60% 55% 6% 39%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 6086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amino acid--[acyl-carrier-protein] ligase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	295	Total	C	N	O	S	0	7	0
			2300	1451	418	413	18			
1	B	296	Total	C	N	O	S	0	10	0
			2349	1483	423	425	18			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	EXPRESSION TAG	UNP Q89VT8
A	-18	GLY	-	EXPRESSION TAG	UNP Q89VT8
A	-17	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-16	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-15	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-14	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-13	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-12	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-11	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-10	HIS	-	EXPRESSION TAG	UNP Q89VT8
A	-9	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-8	SER	-	EXPRESSION TAG	UNP Q89VT8
A	-7	GLY	-	EXPRESSION TAG	UNP Q89VT8
A	-6	LEU	-	EXPRESSION TAG	UNP Q89VT8
A	-5	VAL	-	EXPRESSION TAG	UNP Q89VT8
A	-4	PRO	-	EXPRESSION TAG	UNP Q89VT8
A	-3	ARG	-	EXPRESSION TAG	UNP Q89VT8
A	-2	GLY	-	EXPRESSION TAG	UNP Q89VT8
A	-1	SER	-	EXPRESSION TAG	UNP Q89VT8
A	0	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-19	MET	-	EXPRESSION TAG	UNP Q89VT8
B	-18	GLY	-	EXPRESSION TAG	UNP Q89VT8
B	-17	SER	-	EXPRESSION TAG	UNP Q89VT8
B	-16	SER	-	EXPRESSION TAG	UNP Q89VT8
B	-15	HIS	-	EXPRESSION TAG	UNP Q89VT8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-13	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-12	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-11	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-10	HIS	-	EXPRESSION TAG	UNP Q89VT8
B	-9	SER	-	EXPRESSION TAG	UNP Q89VT8
B	-8	SER	-	EXPRESSION TAG	UNP Q89VT8
B	-7	GLY	-	EXPRESSION TAG	UNP Q89VT8
B	-6	LEU	-	EXPRESSION TAG	UNP Q89VT8
B	-5	VAL	-	EXPRESSION TAG	UNP Q89VT8
B	-4	PRO	-	EXPRESSION TAG	UNP Q89VT8
B	-3	ARG	-	EXPRESSION TAG	UNP Q89VT8
B	-2	GLY	-	EXPRESSION TAG	UNP Q89VT8
B	-1	SER	-	EXPRESSION TAG	UNP Q89VT8
B	0	HIS	-	EXPRESSION TAG	UNP Q89VT8

- Molecule 2 is a protein called Aminoacyl carrier protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	77	Total	C	N	O	S	0	0	0
			546	349	90	105	2			
2	D	63	Total	C	N	O	S	0	0	0
			413	264	70	78	1			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-19	MET	-	EXPRESSION TAG	UNP A9CHM9
C	-18	GLY	-	EXPRESSION TAG	UNP A9CHM9
C	-17	SER	-	EXPRESSION TAG	UNP A9CHM9
C	-16	SER	-	EXPRESSION TAG	UNP A9CHM9
C	-15	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-14	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-13	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-12	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-11	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-10	HIS	-	EXPRESSION TAG	UNP A9CHM9
C	-9	SER	-	EXPRESSION TAG	UNP A9CHM9
C	-8	SER	-	EXPRESSION TAG	UNP A9CHM9
C	-7	GLY	-	EXPRESSION TAG	UNP A9CHM9
C	-6	LEU	-	EXPRESSION TAG	UNP A9CHM9
C	-5	VAL	-	EXPRESSION TAG	UNP A9CHM9

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-4	PRO	-	EXPRESSION TAG	UNP A9CHM9
C	-3	ARG	-	EXPRESSION TAG	UNP A9CHM9
C	-2	GLY	-	EXPRESSION TAG	UNP A9CHM9
C	-1	SER	-	EXPRESSION TAG	UNP A9CHM9
C	0	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-19	MET	-	EXPRESSION TAG	UNP A9CHM9
D	-18	GLY	-	EXPRESSION TAG	UNP A9CHM9
D	-17	SER	-	EXPRESSION TAG	UNP A9CHM9
D	-16	SER	-	EXPRESSION TAG	UNP A9CHM9
D	-15	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-14	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-13	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-12	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-11	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-10	HIS	-	EXPRESSION TAG	UNP A9CHM9
D	-9	SER	-	EXPRESSION TAG	UNP A9CHM9
D	-8	SER	-	EXPRESSION TAG	UNP A9CHM9
D	-7	GLY	-	EXPRESSION TAG	UNP A9CHM9
D	-6	LEU	-	EXPRESSION TAG	UNP A9CHM9
D	-5	VAL	-	EXPRESSION TAG	UNP A9CHM9
D	-4	PRO	-	EXPRESSION TAG	UNP A9CHM9
D	-3	ARG	-	EXPRESSION TAG	UNP A9CHM9
D	-2	GLY	-	EXPRESSION TAG	UNP A9CHM9
D	-1	SER	-	EXPRESSION TAG	UNP A9CHM9
D	0	HIS	-	EXPRESSION TAG	UNP A9CHM9

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	B	1	Total Zn 1 1	0	0

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
4	A	1	31	10	5	13	3	0	0
4	B	1	31	10	5	13	3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
5	A	1	1	1	0	0

- Molecule 6 is 4'-PHOSPHOPANTETHEINE (three-letter code: PNS) (formula: C₁₁H₂₃N₂O₇PS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
6	C	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		
6	D	1	Total	C	N	O	P	S	0	0
			21	11	2	6	1	1		



- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	190	Total	O	0	0
			190	190		
7	B	180	Total	O	0	0
			180	180		
7	D	1	Total	O	0	0
			1	1		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amino acid--[acyl-carrier-protein] ligase 1



- Molecule 1: Amino acid--[acyl-carrier-protein] ligase 1




- Molecule 2: Aminoacyl carrier protein



- Molecule 2: Aminoacyl carrier protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.86Å 101.59Å 103.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.08 – 2.10 46.08 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.08-2.10) 99.8 (46.08-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.10Å)	Xtriage
Refinement program	PHENIX dev_1116	Depositor
R, R_{free}	0.176 , 0.210 0.179 , 0.212	Depositor DCC
R_{free} test set	3103 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h,l,k 0.013 for -l,-k,-h 0.019 for k,h,-l 0.005 for k,l,h 0.005 for l,h,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6086	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, PNS, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2377	0.53	0/3222
1	B	0.39	0/2437	0.53	0/3302
2	C	0.26	0/554	0.46	0/756
2	D	0.24	0/419	0.41	0/571
All	All	0.38	0/5787	0.52	0/7851

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2300	0	2242	11	0
1	B	2349	0	2290	17	0
2	C	546	0	514	13	0
2	D	413	0	336	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	31	0	12	6	0
4	B	31	0	12	4	0
5	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	21	0	21	0	0
6	D	21	0	21	1	0
7	A	190	0	0	3	0
7	B	180	0	0	1	0
7	D	1	0	0	0	0
All	All	6086	0	5448	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:MET:HB2	2:C:58:LEU:HD22	1.71	0.72
1:A:160[B]:ARG:NH1	7:A:585:HOH:O	2.26	0.69
1:B:45:SER:O	1:B:49[B]:ARG:HG2	1.97	0.65
4:A:402:ATP:H5'1	4:A:402:ATP:H8	1.63	0.64
2:C:5:ILE:HA	2:C:8:ILE:HD12	1.85	0.58
1:A:126[A]:LEU:HD11	1:B:126:LEU:HD11	1.88	0.55
1:A:18:ASP:OD1	1:A:165:HIS:ND1	2.39	0.54
1:A:174[B]:MET:SD	4:A:402:ATP:O4'	2.68	0.52
2:C:8:ILE:HA	2:C:11:LYS:HB2	1.91	0.52
2:D:28:LEU:N	2:D:64:ALA:O	2.43	0.52
1:A:168:ARG:HH22	4:A:402:ATP:PG	2.34	0.49
2:C:42:MET:CE	2:C:58:LEU:HD21	2.43	0.49
1:B:273[A]:GLU:OE1	7:B:630:HOH:O	2.20	0.48
1:A:235:LYS:NZ	4:A:402:ATP:O2A	2.43	0.47
4:B:402:ATP:H5'1	4:B:402:ATP:H8	1.79	0.47
1:A:168:ARG:NH1	4:A:402:ATP:O3G	2.34	0.47
1:B:174[A]:MET:SD	4:B:402:ATP:O4'	2.73	0.47
2:D:41:LEU:O	2:D:45:ILE:HG12	2.15	0.46
7:A:648:HOH:O	6:D:101:PNS:H432	2.16	0.46
1:B:187:SER:O	1:B:191[B]:GLU:HG2	2.16	0.45
1:B:49[A]:ARG:HD3	1:B:310:LEU:O	2.16	0.45
2:C:42:MET:HE1	2:C:46:GLU:OE2	2.16	0.45
1:B:93[B]:CYS:SG	1:B:119:LEU:HD13	2.56	0.45
2:C:29:TYR:CE2	2:C:64:ALA:HA	2.52	0.44
1:B:220:ARG:HH12	2:C:42:MET:HE1	1.82	0.44
1:A:202:ARG:HD2	7:A:632:HOH:O	2.17	0.44
4:A:402:ATP:H5'1	4:A:402:ATP:C8	2.48	0.44
1:B:286:ARG:HH11	4:B:402:ATP:H2'	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:LEU:HD22	1:B:125:VAL:HG21	2.00	0.43
2:D:39:VAL:O	2:D:43:LEU:HG	2.18	0.43
1:B:144:LEU:HD11	1:B:180:ILE:HG22	2.01	0.43
1:B:212:TYR:CD1	1:B:233:GLN:HG2	2.54	0.42
2:C:4:THR:OG1	2:C:5:ILE:N	2.53	0.42
2:C:57:LEU:HD13	2:C:72:THR:HG23	2.01	0.42
1:B:235:LYS:NZ	4:B:402:ATP:O2A	2.48	0.42
2:C:58:LEU:HD23	2:C:58:LEU:HA	1.76	0.42
1:B:73:ARG:HB2	1:B:125:VAL:HG13	2.03	0.41
2:C:28:LEU:HG	2:C:65:SER:HA	2.02	0.41
1:A:115:TRP:HH2	1:B:91[B]:CYS:SG	2.44	0.41
1:A:18:ASP:HA	1:A:19:PRO:HD3	1.90	0.41
2:C:0:HIS:O	2:C:4:THR:HG23	2.20	0.41
2:C:35:SER:O	2:C:39:VAL:HG23	2.21	0.41
1:A:160[A]:ARG:HD2	1:B:96:HIS:NE2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/346 (87%)	296 (99%)	4 (1%)	0	100	100
1	B	304/346 (88%)	301 (99%)	3 (1%)	0	100	100
2	C	75/103 (73%)	69 (92%)	6 (8%)	0	100	100
2	D	59/103 (57%)	58 (98%)	1 (2%)	0	100	100
All	All	738/898 (82%)	724 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/279 (85%)	235 (99%)	2 (1%)	81	86
1	B	244/279 (88%)	241 (99%)	3 (1%)	71	77
2	C	51/84 (61%)	51 (100%)	0	100	100
2	D	30/84 (36%)	30 (100%)	0	100	100
All	All	562/726 (77%)	557 (99%)	5 (1%)	78	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	VAL
1	A	156	ASP
1	B	134	VAL
1	B	156	ASP
1	B	173	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	PNS	C	101	2	15,20,21	0.54	0	18,26,29	0.68	1 (5%)
6	PNS	D	101	2	15,20,21	0.44	0	18,26,29	0.55	0
4	ATP	B	402	-	28,33,33	1.61	2 (7%)	34,52,52	1.77	7 (20%)
4	ATP	A	402	5	28,33,33	1.74	2 (7%)	34,52,52	1.73	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PNS	C	101	2	-	4/24/26/27	-
6	PNS	D	101	2	-	1/24/26/27	-
4	ATP	B	402	-	-	5/18/38/38	0/3/3/3
4	ATP	A	402	5	-	7/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	402	ATP	PB-O3A	6.07	1.66	1.59
4	A	402	ATP	PA-O3A	6.00	1.66	1.59
4	B	402	ATP	PB-O3A	5.86	1.65	1.59
4	B	402	ATP	PA-O3A	5.31	1.65	1.59

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	402	ATP	O3A-PA-O1A	-4.77	96.34	110.70
4	A	402	ATP	O3A-PB-O1B	-4.60	96.87	110.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	402	ATP	O3A-PB-O1B	-4.42	97.40	110.70
4	B	402	ATP	O2B-PB-O3B	-4.37	95.45	107.27
4	B	402	ATP	O3A-PA-O1A	-4.20	98.07	110.70
4	A	402	ATP	O2B-PB-O3B	-3.14	98.78	107.27
4	B	402	ATP	O2B-PB-O1B	3.00	126.41	112.44
4	A	402	ATP	O2B-PB-O1B	2.98	126.31	112.44
4	B	402	ATP	O2A-PA-O5'	-2.91	94.39	107.57
4	A	402	ATP	O2A-PA-O5'	-2.73	95.21	107.57
4	B	402	ATP	O2A-PA-O1A	2.60	124.55	112.44
4	A	402	ATP	O2A-PA-O1A	2.55	124.30	112.44
4	B	402	ATP	C5-C6-N6	2.19	123.65	120.31
6	C	101	PNS	C43-C42-N41	-2.04	107.67	112.31

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	402	ATP	PB-O3B-PG-O2G
4	A	402	ATP	C5'-O5'-PA-O1A
4	B	402	ATP	C5'-O5'-PA-O1A
6	C	101	PNS	N41-C42-C43-S44
6	D	101	PNS	N41-C42-C43-S44
4	A	402	ATP	PB-O3B-PG-O3G
6	C	101	PNS	O27-C28-C29-C30
6	C	101	PNS	O27-C28-C29-C31
6	C	101	PNS	O27-C28-C29-C32
4	A	402	ATP	C5'-O5'-PA-O2A
4	A	402	ATP	C5'-O5'-PA-O3A
4	B	402	ATP	C5'-O5'-PA-O2A
4	B	402	ATP	C5'-O5'-PA-O3A
4	A	402	ATP	PB-O3A-PA-O2A
4	B	402	ATP	PB-O3A-PA-O2A
4	A	402	ATP	O4'-C4'-C5'-O5'
4	B	402	ATP	PB-O3A-PA-O1A

There are no ring outliers.

3 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	D	101	PNS	1	0
4	B	402	ATP	4	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	402	ATP	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/346 (85%)	-0.13	2 (0%) 87 89	12, 24, 53, 67	0
1	B	296/346 (85%)	-0.30	3 (1%) 82 85	14, 26, 45, 63	0
2	C	77/103 (74%)	2.41	46 (59%) 0 0	52, 69, 90, 96	0
2	D	63/103 (61%)	6.39	62 (98%) 0 0	74, 108, 120, 126	0
All	All	731/898 (81%)	0.63	113 (15%) 2 2	12, 28, 107, 126	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	45	ILE	18.7
2	D	3	ALA	13.1
2	D	73	VAL	11.1
2	D	5	ILE	10.7
2	D	32	GLY	10.5
2	D	48	ALA	10.4
2	D	53	PHE	10.2
2	D	63	PHE	9.3
2	D	31	ALA	9.0
2	D	49	PHE	8.9
2	D	41	LEU	8.6
2	D	69	ILE	8.3
2	D	51	ILE	8.2
2	D	10	ALA	7.8
2	D	8	ILE	7.8
2	D	74	LYS	7.7
2	D	61	LYS	7.5
2	D	54	PRO	7.5
2	D	68	ALA	7.4
2	D	39	VAL	7.3
2	D	29	TYR	7.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	75	LEU	7.2
2	D	43	LEU	7.1
2	D	13	GLY	7.0
2	D	4	THR	6.7
2	D	50	ASP	6.5
2	D	12	PHE	6.4
2	D	30	ALA	6.0
2	D	52	GLU	6.0
2	C	21	THR	5.9
1	B	17	ALA	5.9
2	D	47	GLU	5.9
2	D	62	SER	5.8
2	D	15	LEU	5.8
2	C	48	ALA	5.7
2	C	17	THR	5.7
2	D	6	ARG	5.7
2	D	64	ALA	5.5
2	D	44	GLY	5.4
2	D	57	LEU	5.2
2	D	60	ARG	5.1
2	D	36	PHE	5.0
2	D	40	GLN	5.0
2	D	42	MET	5.0
2	D	33	LEU	4.9
2	D	66	ILE	4.9
2	D	9	LEU	4.8
2	D	72	THR	4.8
2	D	46	GLU	4.7
2	C	16	PRO	4.7
2	D	71	ASP	4.6
2	D	58	LEU	4.6
2	D	55	ASP	4.6
2	D	70	GLU	4.5
2	D	2	ASN	4.5
2	D	34	SER	4.4
2	D	16	PRO	4.3
2	D	65	SER	4.3
2	C	54	PRO	4.3
2	D	14	GLN	4.3
2	C	57	LEU	4.1
2	D	56	ASN	4.1
2	D	28	LEU	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	73	VAL	4.1
2	D	11	LYS	4.1
2	C	22	ILE	4.0
2	C	75	LEU	4.0
2	C	30	ALA	4.0
2	C	12	PHE	3.9
2	D	7	GLU	3.9
2	C	15	LEU	3.8
2	C	19	VAL	3.8
2	C	20	ASP	3.8
2	C	31	ALA	3.7
2	C	55	ASP	3.7
2	C	69	ILE	3.6
2	C	53	PHE	3.5
1	A	224	MET	3.5
2	C	72	THR	3.4
2	C	64	ALA	3.3
2	C	58	LEU	3.2
2	C	47	GLU	3.2
2	D	38	SER	3.1
2	D	67	LYS	3.1
2	C	25	GLU	3.1
2	C	49	PHE	3.0
2	C	26	ALA	2.9
2	C	50	ASP	2.9
2	C	56	ASN	2.9
2	C	33	LEU	2.9
2	C	46	GLU	2.8
2	C	63	PHE	2.8
2	C	18	PRO	2.8
2	C	76	ILE	2.8
2	C	71	ASP	2.6
2	C	51	ILE	2.6
2	C	9	LEU	2.5
2	C	74	LYS	2.5
2	C	11	LYS	2.4
2	C	0	HIS	2.4
2	C	52	GLU	2.4
2	D	37	ALA	2.4
1	B	219	GLY	2.4
2	D	35	SER	2.3
2	C	28	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	18	ASP	2.2
2	C	5	ILE	2.2
2	C	1	MET	2.2
2	C	27	ASP	2.1
2	C	29	TYR	2.1
2	C	65	SER	2.0
1	A	225	LEU	2.0
2	C	39	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	PNS	D	101	21/22	0.84	0.20	53,60,75,85	0
4	ATP	A	402	31/31	0.87	0.16	14,35,61,64	31
4	ATP	B	402	31/31	0.89	0.15	16,32,60,69	31
6	PNS	C	101	21/22	0.91	0.23	53,61,70,79	0
5	MG	A	403	1/1	0.91	0.44	51,51,51,51	0
3	ZN	A	401	1/1	0.97	0.07	38,38,38,38	1
3	ZN	B	401	1/1	0.99	0.09	32,32,32,32	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PNS D 101:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ATP B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.