



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 20, 2024 – 05:45 AM EST

PDB ID : 4LO2  
Title : HA17-HA33-Lac  
Authors : Lee, K.; Gu, S.; Jin, L.; Le, T.T.; Cheng, L.W.; Strotmeier, J.; Krueel, A.M.; Yao, G.; Perry, K.; Rummel, A.; Jin, R.  
Deposited on : 2013-07-12  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	
1	B	296	
2	C	147	
3	D	2	
3	E	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
3	BGC	E	1	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6126 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HA-33.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2334	1477	399	455	3	8	1	0
1	B	286	2334	1477	399	455	3	10	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	294	PRO	-	expression tag	UNP Q45871
A	295	GLY	-	expression tag	UNP Q45871
A	296	SER	-	expression tag	UNP Q45871
A	297	ALA	-	expression tag	UNP Q45871
B	294	PRO	-	expression tag	UNP Q45871
B	295	GLY	-	expression tag	UNP Q45871
B	296	SER	-	expression tag	UNP Q45871
B	297	ALA	-	expression tag	UNP Q45871

- Molecule 2 is a protein called HA-17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	144	1187	767	189	226	5	3	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	GLY	-	expression tag	UNP Q45878
C	1	PRO	-	expression tag	UNP Q45878

- Molecule 3 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	D	2	Total	C	O	0	0	0
			23	12	11			
3	E	2	Total	C	O	0	0	0
			23	12	11			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	102	Total	O	0	0
			102	102		
4	C	60	Total	O	0	0
			60	60		
4	B	63	Total	O	0	0
			63	63		


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HA-33




- Molecule 1: HA-33



- Molecule 2: HA-17



- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



- Molecule 3: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



BGCI  
GAL2

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.49Å 119.15Å 163.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.06 – 2.25 32.06 – 2.25	Depositor EDS
% Data completeness (in resolution range)	97.3 (32.06-2.25) 97.3 (32.06-2.25)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.09 (at 2.26Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.7.2_869)	Depositor
R, $R_{free}$	0.166 , 0.211 0.161 , 0.204	Depositor DCC
$R_{free}$ test set	2430 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	38.1	Xtrriage
Anisotropy	0.576	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 53.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6126	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/2393	0.53	0/3267
1	B	0.37	0/2393	0.52	0/3267
2	C	0.40	0/1217	0.54	0/1653
All	All	0.38	0/6003	0.53	0/8187

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2334	0	2250	17	0
1	B	2334	0	2250	9	0
2	C	1187	0	1150	7	0
3	D	23	0	21	0	0
3	E	23	0	21	0	0
4	A	102	0	0	4	0
4	B	63	0	0	0	0
4	C	60	0	0	0	0
All	All	6126	0	5692	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:ASN:N	4:A:479:HOH:O	2.26	0.68
1:B:176:ASN:HA	1:B:224:ARG:HG2	1.78	0.66
1:B:210:ASN:ND2	1:B:228:SER:OG	2.35	0.59
1:A:265:TYR:O	1:A:268:GLN:HG2	2.06	0.55
1:A:186:ARG:HB2	1:A:207:ILE:HD13	1.88	0.55
2:C:106:ASN:O	2:C:107:GLU:HG2	2.10	0.52
1:B:257:ASP:HB3	1:B:260:LYS:HG3	1.90	0.52
1:A:193:ARG:NH2	4:A:497:HOH:O	2.36	0.51
1:A:223:VAL:HG13	1:A:276:GLN:HA	1.93	0.50
2:C:55:CYS:SG	2:C:102:LEU:HD13	2.52	0.49
1:A:131:THR:HB	1:B:115:ASN:HD21	1.80	0.47
1:B:241:ASN:ND2	1:B:251:THR:OG1	2.45	0.47
1:A:216:ILE:HB	1:A:222:THR:OG1	2.15	0.46
2:C:6:THR:OG1	2:C:7:PHE:N	2.48	0.46
1:A:152:ASN:N	1:A:192:ILE:O	2.50	0.45
2:C:18:ILE:HG12	2:C:140:MET:HB2	1.97	0.45
1:A:172:VAL:HG13	1:A:173:THR:HG23	1.98	0.45
1:A:165:LYS:HE3	1:A:181:THR:HG23	2.00	0.44
1:A:246:THR:HB	4:A:474:HOH:O	2.18	0.44
2:C:83:ASN:HA	2:C:86:TYR:CE2	2.53	0.44
1:A:281:HIS:CE1	1:A:283:ASP:HB2	2.53	0.43
1:A:207:ILE:HD12	1:A:207:ILE:HA	1.78	0.43
1:A:248:GLU:O	1:A:289:ASN:HA	2.19	0.42
1:B:241:ASN:HD22	1:B:251:THR:HG1	1.68	0.42
1:B:223:VAL:HG22	1:B:275:ILE:O	2.20	0.42
1:A:268:GLN:HG2	1:A:268:GLN:H	1.69	0.41
2:C:100:LEU:HD23	2:C:111:ALA:HB2	2.01	0.41
1:A:102:GLY:HA2	1:B:9:ASN:HB2	2.03	0.40
1:A:104:ASN:ND2	4:A:498:HOH:O	2.54	0.40
2:C:105:VAL:HB	2:C:110:TYR:CD2	2.56	0.40
1:B:215:TRP:CZ3	1:B:256:ARG:HD3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/296 (96%)	277 (97%)	8 (3%)	0	100	100
1	B	285/296 (96%)	275 (96%)	10 (4%)	0	100	100
2	C	142/147 (97%)	139 (98%)	3 (2%)	0	100	100
All	All	712/739 (96%)	691 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	263/270 (97%)	249 (95%)	14 (5%)	22	23
1	B	263/270 (97%)	254 (97%)	9 (3%)	37	45
2	C	136/138 (99%)	133 (98%)	3 (2%)	52	61
All	All	662/678 (98%)	636 (96%)	26 (4%)	32	38

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	ASP
1	A	30	GLN
1	A	68	ASN
1	A	80	HIS
1	A	123	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	143	GLU
1	A	180	TYR
1	A	186	ARG
1	A	198	LYS
1	A	209	SER
1	A	223	VAL
1	A	232	ASN
1	A	244	SER
1	A	268	GLN
2	C	26	ASN
2	C	79	ASP
2	C	122	LEU
1	B	13	ASP
1	B	30	GLN
1	B	103	ASN
1	B	198	LYS
1	B	210	ASN
1	B	241	ASN
1	B	260	LYS
1	B	281	HIS
1	B	292	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	229	ASN
1	B	67	HIS
1	B	68	ASN
1	B	115	ASN
1	B	210	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates i

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	BGC	D	1	3	12,12,12	1.61	2 (16%)	17,17,17	1.55	4 (23%)
3	GAL	D	2	3	11,11,12	1.86	4 (36%)	15,15,17	1.27	2 (13%)
3	BGC	E	1	3	12,12,12	1.66	2 (16%)	17,17,17	1.13	2 (11%)
3	GAL	E	2	3	11,11,12	2.02	5 (45%)	15,15,17	1.33	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	D	1	3	-	2/2/22/22	0/1/1/1
3	GAL	D	2	3	-	1/2/19/22	0/1/1/1
3	BGC	E	1	3	-	0/2/22/22	0/1/1/1
3	GAL	E	2	3	-	0/2/19/22	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1	BGC	C3-C2	-3.96	1.42	1.52
3	D	1	BGC	C3-C2	-3.87	1.42	1.52
3	E	2	GAL	C4-C3	-3.79	1.42	1.52
3	D	2	GAL	C4-C3	-3.66	1.43	1.52
3	E	2	GAL	O5-C5	3.37	1.50	1.43
3	D	2	GAL	O5-C5	2.78	1.49	1.43
3	E	2	GAL	C2-C3	-2.63	1.48	1.52
3	E	1	BGC	O5-C1	2.63	1.49	1.42
3	D	2	GAL	O5-C1	-2.43	1.39	1.43
3	D	1	BGC	O5-C1	2.36	1.48	1.42
3	E	2	GAL	O2-C2	-2.25	1.38	1.43

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	2	GAL	O2-C2	-2.24	1.38	1.43
3	E	2	GAL	O5-C1	-2.10	1.40	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	BGC	C3-C4-C5	3.10	115.77	110.24
3	D	1	BGC	O5-C5-C4	3.10	115.32	109.69
3	D	2	GAL	C1-C2-C3	3.00	113.36	109.67
3	E	1	BGC	O3-C3-C2	-2.83	103.80	110.35
3	D	1	BGC	O3-C3-C2	-2.65	104.23	110.35
3	E	2	GAL	O3-C3-C2	-2.48	105.24	109.99
3	D	1	BGC	C4-C3-C2	2.38	114.97	110.82
3	E	2	GAL	C3-C4-C5	2.22	114.20	110.24
3	D	2	GAL	O3-C3-C2	-2.19	105.79	109.99
3	E	1	BGC	O6-C6-C5	2.06	118.36	111.29

There are no chirality outliers.


All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	1	BGC	C4-C5-C6-O6
3	D	1	BGC	O5-C5-C6-O6
3	D	2	GAL	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/296 (96%)	-0.12	8 (2%) 53 55	23, 45, 80, 121	2 (0%)
1	B	286/296 (96%)	0.59	44 (15%) 2 2	25, 58, 122, 153	3 (1%)
2	C	144/147 (97%)	0.09	7 (4%) 29 32	25, 45, 70, 90	1 (0%)
All	All	716/739 (96%)	0.21	59 (8%) 11 12	23, 48, 107, 153	6 (0%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	ASN	9.0
1	B	161	LEU	7.8
1	B	229	ASN	7.7
1	B	9	ASN	7.3
1	B	228	SER	7.0
1	B	283	ASP	6.4
1	B	230	ASP	6.0
1	B	217	PHE	6.0
1	A	232	ASN	5.9
1	B	231	GLN	5.9
1	B	265	TYR	5.5
1	B	278	PHE	4.8
1	B	160	ILE	4.6
1	B	245	ASP	4.6
1	B	67	HIS	4.6
1	B	162	ASP	4.6
1	A	229	ASN	4.3
1	B	233	ASN	4.3
1	B	234	ASP	4.2
1	B	215	TRP	4.1
2	C	94	VAL	4.0
1	B	284	ASP	3.9
1	A	294	PRO	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	264	LEU	3.6
1	B	244	SER	3.6
1	B	197	GLU	3.6
1	B	280	TYR	3.5
1	B	268	GLN	3.5
1	A	197	GLU	3.4
1	B	10	SER	3.2
1	B	243	VAL	3.1
1	A	228	SER	2.9
2	C	39	SER	2.9
1	B	163	LEU	2.9
1	B	216	ILE	2.9
1	B	246	THR	2.9
1	B	282	GLY	2.9
1	B	269	THR	2.8
1	B	68	ASN	2.7
1	B	258	THR	2.7
1	B	219	ASN	2.6
1	B	32	ALA	2.6
1	B	218	SER	2.6
1	A	230	ASP	2.6
1	A	9	ASN	2.6
1	B	281	HIS	2.5
1	A	231	GLN	2.5
2	C	100	LEU	2.5
1	B	270	ALA	2.4
1	B	213	LEU	2.3
2	C	88	PHE	2.3
1	B	220	GLY	2.2
2	C	40	ALA	2.2
1	B	222	THR	2.2
1	B	164	ASN	2.1
2	C	20	SER	2.1
1	B	66	ILE	2.0
1	B	227	SER	2.0
2	C	64	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains


There are no non-standard protein/DNA/RNA residues in this entry.


### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BGC	E	1	12/12	0.61	0.47	121,133,136,137	0
3	GAL	E	2	11/12	0.83	0.25	98,108,112,114	0
3	BGC	D	1	12/12	0.91	0.21	72,86,97,103	0
3	GAL	D	2	11/12	0.96	0.08	51,54,60,60	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.