



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 19, 2024 – 01:11 AM EDT

PDB ID : 4NGE
Title : Crystal Structure of Human Presequence Protease in Complex with Amyloid-beta (1-40)
Authors : King, J.V.; Liang, W.G.; Tang, W.J.
Deposited on : 2013-11-01
Resolution : 2.70 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1014	
2	B	40	
2	E	40	
3	C	7	
3	F	7	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	1014	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	ACT	A	1103	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 16054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Presequence protease, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	As	C	N	O				S
1	A	985	7910	4	5046	1337	1483	40	0	2	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	MET	-	EXPRESSION TAG	UNP Q5JRX3
A	25	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	26	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	27	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	28	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	29	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	30	HIS	-	EXPRESSION TAG	UNP Q5JRX3
A	31	ALA	-	EXPRESSION TAG	UNP Q5JRX3
A	32	ALA	-	EXPRESSION TAG	UNP Q5JRX3
A	107	GLN	GLU	ENGINEERED MUTATION	UNP Q5JRX3
A	328	VAL	ILE	SEE REMARK 999	UNP Q5JRX3
A	397	VAL	ALA	SEE REMARK 999	UNP Q5JRX3
A	1037	ARG	GLN	SEE REMARK 999	UNP Q5JRX3

- Molecule 2 is a protein called Beta-amyloid protein 40.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	9	77	52	11	14	0	0	0
2	E	5	43	31	5	7	0	0	0

- Molecule 3 is a protein called Beta-amyloid protein 40.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	4	Total	C	N	O	0	0	0
			20	12	4	4			
3	F	1	Total	C	N	O	0	0	0
			5	3	1	1			

- Molecule 4 is a protein called Presequence protease, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	987	Total	As	C	N	O	S	0	1	0
			7931	1	5062	1345	1483	40			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	24	MET	-	EXPRESSION TAG	UNP Q5JRX3
D	25	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	26	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	27	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	28	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	29	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	30	HIS	-	EXPRESSION TAG	UNP Q5JRX3
D	31	ALA	-	EXPRESSION TAG	UNP Q5JRX3
D	32	ALA	-	EXPRESSION TAG	UNP Q5JRX3
D	107	GLN	GLU	ENGINEERED MUTATION	UNP Q5JRX3
D	328	VAL	ILE	SEE REMARK 999	UNP Q5JRX3
D	397	VAL	ALA	SEE REMARK 999	UNP Q5JRX3
D	1037	ARG	GLN	SEE REMARK 999	UNP Q5JRX3

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Zn	0	0
			1	1		
5	E	1	Total	Zn	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		
6	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	11	Total	O	0	0
			11	11		
8	D	28	Total	O	0	0
			28	28		
8	E	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Presequence protease, mitochondrial





- Molecule 2: Beta-amyloid protein 40



- Molecule 2: Beta-amyloid protein 40





• Molecule 3: Beta-amyloid protein 40



• Molecule 3: Beta-amyloid protein 40



• Molecule 4: Presequence protease, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	247.28Å 86.18Å 158.61Å 90.00° 127.56° 90.00°	Depositor
Resolution (Å)	39.62 – 2.70 39.62 – 2.70	Depositor EDS
% Data completeness (in resolution range)	88.0 (39.62-2.70) 88.4 (39.62-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.83 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.191 , 0.232 0.195 , 0.233	Depositor DCC
R_{free} test set	3371 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.017 for -h-2*1,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	16054	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLY, ZN, CAS, ACT, GOL, MLZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/7877	0.43	0/10688
2	B	0.46	0/78	1.00	0/103
2	E	0.63	0/44	1.14	0/58
4	D	0.26	0/7885	0.43	1/10709 (0.0%)
All	All	0.26	0/15884	0.44	1/21558 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	87	LEU	CA-CB-CG	5.35	127.59	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7910	0	7788	92	0
2	B	77	0	73	31	0
2	E	43	0	37	9	0
3	C	20	0	9	5	0
3	F	5	0	3	3	0
4	D	7931	0	7847	104	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	1	0	0	0	0
5	E	1	0	0	0	0
6	A	6	0	8	0	0
6	D	12	0	16	0	0
7	A	8	0	6	1	0
8	A	11	0	0	1	0
8	D	28	0	0	5	0
8	E	1	0	0	0	0
All	All	16054	0	15787	217	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 217 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:16:LYS:HD3	2:B:17:LEU:CD1	1.62	1.27
4:D:575:ASP:OD2	8:D:1218:HOH:O	1.54	1.20
2:B:16:LYS:HG2	2:B:17:LEU:N	1.57	1.13
4:D:562:VAL:O	4:D:565:ILE:HG22	1.54	1.06
2:B:16:LYS:HG2	2:B:17:LEU:H	0.92	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	958/1014 (94%)	933 (97%)	25 (3%)	0	100 100
2	B	7/40 (18%)	3 (43%)	4 (57%)	0	100 100
2	E	3/40 (8%)	3 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
4	D	960/1014 (95%)	933 (97%)	26 (3%)	1 (0%)	51 78
All	All	1928/2108 (92%)	1872 (97%)	55 (3%)	1 (0%)	51 78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	230	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	857/880 (97%)	854 (100%)	3 (0%)	91 97
2	B	8/31 (26%)	4 (50%)	4 (50%)	0 0
2	E	4/31 (13%)	1 (25%)	3 (75%)	0 0
4	D	859/879 (98%)	852 (99%)	7 (1%)	81 93
All	All	1728/1821 (95%)	1711 (99%)	17 (1%)	76 91

5 of 17 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	E	19	PHE
2	E	22	GLU
4	D	115	GLN
4	D	194	THR
4	D	565	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	690	ASN
1	A	1025	ASN
2	B	15	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	D	325	GLN
4	D	852	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

43 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLZ	A	946	1	8,9,10	1.19	1 (12%)	4,9,11	0.74	0
4	MLY	D	642	4	9,10,11	0.48	0	6,11,13	0.84	0
1	MLY	A	251	1	9,10,11	0.55	0	6,11,13	0.83	0
1	MLY	A	278	1	9,10,11	0.51	0	6,11,13	0.86	0
1	MLY	A	363	1	9,10,11	0.54	0	6,11,13	0.87	0
4	MLY	D	972	4	9,10,11	0.52	0	6,11,13	0.79	0
4	CAS	D	556	4	5,8,9	1.32	0	1,9,11	0.66	0
1	MLY	A	199	1	9,10,11	0.51	0	6,11,13	0.82	0
4	MLY	D	943	4	9,10,11	0.60	0	6,11,13	0.75	0
1	MLY	A	41	1	9,10,11	0.54	0	6,11,13	0.82	0
1	CAS	A	171	1	5,8,9	1.45	1 (20%)	1,9,11	1.55	0
4	MLY	D	946	4	9,10,11	0.52	0	6,11,13	0.80	0
1	CAS	A	112	1	5,8,9	1.32	0	1,9,11	1.07	0
1	MLY	A	525	1	9,10,11	0.50	0	6,11,13	0.85	0
4	MLY	D	287	4	9,10,11	0.52	0	6,11,13	0.97	0
4	MLY	D	769	4	9,10,11	0.60	0	6,11,13	0.84	0
4	MLY	D	125	4	9,10,11	0.55	0	6,11,13	0.79	0
1	MLY	A	911	1	9,10,11	0.53	0	6,11,13	0.84	0
4	MLY	D	854	4	9,10,11	0.51	0	6,11,13	0.96	0
1	MLY	A	466	1	9,10,11	0.51	0	6,11,13	0.95	0
4	MLY	D	764	4	9,10,11	0.53	0	6,11,13	0.79	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	MLY	A	854	1	9,10,11	0.45	0	6,11,13	1.01	0
4	MLY	D	704	4	9,10,11	0.50	0	6,11,13	0.84	0
1	MLY	A	902	1	9,10,11	0.53	0	6,11,13	0.73	0
4	MLY	D	525	4	9,10,11	0.88	0	6,11,13	0.91	0
1	MLY	A	624	1	9,10,11	0.58	0	6,11,13	0.70	0
4	MLY	D	363	4	9,10,11	0.53	0	6,11,13	0.84	0
4	MLY	D	488	4	9,10,11	0.53	0	6,11,13	0.89	0
4	MLY	D	540	4	9,10,11	0.56	0	6,11,13	0.74	0
4	MLY	D	624	4	9,10,11	0.58	0	6,11,13	0.84	0
1	MLY	A	431	1	9,10,11	0.55	0	6,11,13	0.78	0
1	MLY	A	66	1	9,10,11	0.55	0	6,11,13	0.71	0
4	MLY	D	116	4	9,10,11	0.50	0	6,11,13	0.97	0
1	MLY	A	513	1	9,10,11	0.48	0	6,11,13	0.97	0
1	CAS	A	34	1	5,8,9	1.33	0	1,9,11	1.55	0
4	MLY	D	207	4	9,10,11	0.55	0	6,11,13	0.83	0
4	MLY	D	794	4	9,10,11	0.69	0	6,11,13	0.75	0
1	MLY	A	642	1	9,10,11	0.52	0	6,11,13	0.82	0
4	MLY	D	884	4	9,10,11	0.53	0	6,11,13	0.80	0
1	MLY	A	943	1	9,10,11	0.57	0	6,11,13	0.63	0
4	MLY	D	937	4	9,10,11	0.57	0	6,11,13	0.86	0
4	MLY	D	290	4	9,10,11	0.50	0	6,11,13	0.86	0
1	CAS	A	556	1	5,8,9	1.02	0	1,9,11	1.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	MLZ	A	946	1	-	3/7/8/10	-
4	MLY	D	642	4	-	4/8/9/11	-
1	MLY	A	251	1	-	0/8/9/11	-
1	MLY	A	278	1	-	0/8/9/11	-
1	MLY	A	363	1	-	1/8/9/11	-
4	MLY	D	972	4	-	1/8/9/11	-
4	CAS	D	556	4	-	0/0/7/9	-
1	MLY	A	199	1	-	0/8/9/11	-
4	MLY	D	943	4	-	3/8/9/11	-
1	MLY	A	41	1	-	2/8/9/11	-
1	CAS	A	171	1	-	0/0/7/9	-
4	MLY	D	946	4	-	3/8/9/11	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	112	1	-	0/0/7/9	-
1	MLY	A	525	1	-	0/8/9/11	-
4	MLY	D	287	4	-	2/8/9/11	-
4	MLY	D	769	4	-	2/8/9/11	-
4	MLY	D	125	4	-	4/8/9/11	-
1	MLY	A	911	1	-	2/8/9/11	-
4	MLY	D	854	4	-	0/8/9/11	-
1	MLY	A	466	1	-	0/8/9/11	-
4	MLY	D	764	4	-	1/8/9/11	-
1	MLY	A	854	1	-	2/8/9/11	-
4	MLY	D	704	4	-	5/8/9/11	-
1	MLY	A	902	1	-	1/8/9/11	-
4	MLY	D	525	4	-	4/8/9/11	-
1	MLY	A	624	1	-	1/8/9/11	-
4	MLY	D	363	4	-	0/8/9/11	-
4	MLY	D	488	4	-	3/8/9/11	-
4	MLY	D	540	4	-	1/8/9/11	-
4	MLY	D	624	4	-	1/8/9/11	-
1	MLY	A	431	1	-	1/8/9/11	-
1	MLY	A	66	1	-	1/8/9/11	-
4	MLY	D	116	4	-	1/8/9/11	-
1	MLY	A	513	1	-	0/8/9/11	-
1	CAS	A	34	1	-	0/0/7/9	-
4	MLY	D	207	4	-	5/8/9/11	-
4	MLY	D	794	4	-	2/8/9/11	-
1	MLY	A	642	1	-	0/8/9/11	-
4	MLY	D	884	4	-	0/8/9/11	-
1	MLY	A	943	1	-	1/8/9/11	-
4	MLY	D	937	4	-	1/8/9/11	-
4	MLY	D	290	4	-	0/8/9/11	-
1	CAS	A	556	1	-	0/0/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	171	CAS	CA-N	-2.20	1.41	1.48
1	A	946	MLZ	CA-N	-2.02	1.42	1.48

There are no bond angle outliers.

There are no chirality outliers.

5 of 58 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	41	MLY	O-C-CA-CB
1	A	946	MLZ	C-CA-CB-CG
4	D	116	MLY	O-C-CA-CB
4	D	125	MLY	O-C-CA-CB
4	D	207	MLY	N-CA-CB-CG

There are no ring outliers.

12 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	972	MLY	1	0
4	D	943	MLY	1	0
4	D	769	MLY	1	0
1	A	466	MLY	2	0
4	D	704	MLY	1	0
1	A	902	MLY	1	0
1	A	624	MLY	1	0
4	D	363	MLY	1	0
4	D	488	MLY	1	0
4	D	540	MLY	1	0
4	D	794	MLY	1	0
1	A	556	CAS	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	ACT	A	1104	-	3,3,3	0.91	0	3,3,3	0.53	0
6	GOL	D	1101	-	5,5,5	0.21	0	5,5,5	0.26	0
7	ACT	A	1103	-	3,3,3	0.82	0	3,3,3	1.35	0
6	GOL	D	1102	-	5,5,5	0.38	0	5,5,5	0.20	0
6	GOL	A	1102	-	5,5,5	0.36	0	5,5,5	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	D	1101	-	-	4/4/4/4	-
6	GOL	A	1102	-	-	2/4/4/4	-
6	GOL	D	1102	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1102	GOL	O1-C1-C2-C3
6	D	1101	GOL	C1-C2-C3-O3
6	D	1102	GOL	O1-C1-C2-O2
6	D	1102	GOL	O1-C1-C2-C3
6	A	1102	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1104	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	964/1014 (95%)	0.10	37 (3%) 40 39	25, 42, 66, 102	0
2	B	9/40 (22%)	2.92	5 (55%) 0 0	26, 78, 89, 91	0
2	E	5/40 (12%)	2.18	3 (60%) 0 0	71, 73, 93, 99	0
3	C	0/7	-	-	-	-
3	F	0/7	-	-	-	-
4	D	965/1014 (95%)	0.02	23 (2%) 59 60	26, 40, 65, 93	0
All	All	1943/2122 (91%)	0.08	68 (3%) 44 44	25, 41, 67, 102	0

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	17	LEU	8.0
4	D	512	GLU	5.7
4	D	838	VAL	5.2
2	B	15	GLN	5.1
1	A	479	GLN	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	A	34	9/10	0.80	0.18	44,50,84,149	0
4	MLY	D	972	11/12	0.81	0.29	37,54,67,68	0
1	CAS	A	171	9/10	0.82	0.25	33,47,64,174	0
4	CAS	D	556	9/10	0.86	0.32	47,57,84,151	0
4	MLY	D	794	11/12	0.88	0.21	39,47,69,71	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MLY	D	525	11/12	0.88	0.31	20,20,20,20	0
1	CAS	A	556	9/10	0.89	0.29	38,62,116,225	0
4	MLY	D	363	11/12	0.89	0.17	28,38,55,60	0
1	MLY	A	513	11/12	0.89	0.33	43,53,59,59	0
4	MLY	D	290	11/12	0.91	0.18	33,46,56,59	0
4	MLY	D	704	11/12	0.91	0.18	32,38,71,71	0
1	MLY	A	431	11/12	0.91	0.19	29,37,60,62	0
4	MLY	D	937	11/12	0.91	0.20	26,31,48,54	0
1	MLY	A	911	11/12	0.91	0.21	34,40,72,73	0
1	MLY	A	363	11/12	0.92	0.15	34,39,49,50	0
1	MLY	A	41	11/12	0.93	0.19	28,36,48,50	0
1	MLY	A	943	11/12	0.93	0.16	30,41,64,66	0
1	MLY	A	902	11/12	0.93	0.17	27,32,45,51	0
4	MLY	D	943	11/12	0.93	0.15	28,33,52,61	0
4	MLY	D	624	11/12	0.93	0.18	26,34,42,47	0
1	MLY	A	854	11/12	0.94	0.21	42,44,51,51	0
1	MLY	A	199	11/12	0.94	0.18	31,34,53,67	0
1	MLY	A	466	11/12	0.94	0.14	42,51,59,64	0
4	MLY	D	642	11/12	0.94	0.18	32,40,47,51	0
1	MLY	A	278	11/12	0.94	0.24	29,36,54,58	0
4	MLY	D	116	11/12	0.94	0.15	23,33,45,50	0
4	MLY	D	884	11/12	0.94	0.20	24,34,44,55	0
1	MLY	A	525	11/12	0.94	0.20	38,42,57,61	0
1	CAS	A	112	9/10	0.94	0.17	44,46,70,168	0
4	MLY	D	946	11/12	0.94	0.17	32,39,56,60	0
4	MLY	D	488	11/12	0.94	0.17	39,43,55,60	0
1	MLY	A	642	11/12	0.95	0.14	27,37,40,46	0
4	MLY	D	125	11/12	0.95	0.17	27,34,52,57	0
4	MLY	D	207	11/12	0.95	0.17	33,40,58,60	0
1	MLZ	A	946	10/11	0.95	0.15	35,40,53,54	0
4	MLY	D	854	11/12	0.95	0.16	27,37,52,53	0
4	MLY	D	540	11/12	0.96	0.25	31,38,51,54	0
1	MLY	A	251	11/12	0.96	0.20	34,39,49,54	0
1	MLY	A	66	11/12	0.96	0.22	22,35,61,63	0
1	MLY	A	624	11/12	0.96	0.13	29,33,38,56	0
4	MLY	D	287	11/12	0.96	0.18	28,34,44,45	0
4	MLY	D	764	11/12	0.96	0.12	38,47,51,56	0
4	MLY	D	769	11/12	0.96	0.27	36,44,56,60	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	ACT	A	1103	4/4	0.47	0.42	50,73,77,83	0
6	GOL	A	1102	6/6	0.60	0.26	48,61,72,75	0
7	ACT	A	1104	4/4	0.71	0.27	33,59,60,65	0
6	GOL	D	1101	6/6	0.76	0.21	56,65,70,75	0
6	GOL	D	1102	6/6	0.76	0.25	44,58,58,65	0
5	ZN	E	101	1/1	0.99	0.10	45,45,45,45	0
5	ZN	A	1101	1/1	0.99	0.13	44,44,44,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.