

# Full wwPDB EM Validation Report (i)

Apr 29, 2024 – 09:48 pm BST

PDB ID : 4V3A

EMDB ID : EMD-2794

Title: Membrane bound pleurotolysin prepore (TMH1 lock) trapped with engineered

disulphide cross-link

Authors: Lukoyanova, N.; Kondos, S.C.; Farabella, I.; Law, R.H.P.; Reboul, C.F.;

CaradocDavies, T.T.; Spicer, B.A.; Kleifeld, O.; Perugini, M.; Ekkel, S.; Hatfaludi, T.; Oliver, K.; Hotze, E.M.; Tweten, R.K.; Whisstock, J.C.; Topf, M.;

Dunstone, M.A.; Saibil, H.R.

Deposited on : 2014-10-17

Resolution : 15.00 Å(reported)

Based on initial model : 4OEB

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at  $\frac{\text{https://www.wwpdb.org/validation/2017/EMValidationReportHelp}}{\text{with specific help available everywhere you see the (i) symbol.}$ 

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92

MolProbity : 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $MapQ \quad : \quad 1.9.13$ 

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $ELECTRON\ MICROSCOPY$ 

The reported resolution of this entry is 15.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1-A	135	100%
1	1-B	135	100%
1	10-A	135	100%
1	10-B	135	100%
1	11-A	135	100%
1	11-B	135	100%
1	12-A	135	100%
1	12-B	135	100%
1	13-A	135	100%
1	13-B	135	100%



Continued from previous page...

Mol	Chain	Length	Quality of chain
1	14-A	135	100%
1	14-B	135	100%
1	15-A	135	100%
1	15-B	135	100%
1			
	16-A	135	100%
1	16-B	135	100%
1	17-A	135	100%
1	17-B	135	100%
1	18-A	135	100%
1	18-B	135	100%
1	19-A	135	100%
1	19-B	135	100%
1	2-A	135	100%
1	2-B	135	100%
1	20-A	135	100%
1	20-B	135	100%
1	3-A	135	
			100%
1	3-B	135	100%
1	4-A	135	100%
1	4-B	135	100%
1	5-A	135	100%
1	5-B	135	100%
1	6-A	135	100%
1	6-B	135	100%
1	7-A	135	100%



Continued from previous page...

Mol	Chain	$oxed{ egin{array}{c} { m length} \\ { m Length} \\ \end{array} }$	Quality of chain
1	7-B	135	100%
1	8-A	135	100%
1	8-B	135	100%
1	9-A	135	100%
1	9-B	135	100%
2	1-C	388	100%
2	10-C	388	100%
2	11-C	388	100%
2	12-C	388	100%
$\frac{2}{2}$	13-C		
		388	100%
2	14-C	388	100%
2	15-C	388	100%
2	16-C	388	100%
2	17-C	388	100%
2	18-C	388	100%
2	19-C	388	100%
2	2-C	388	100%
2	20-C	388	100%
2	3-C	388	100%
2	4-C	388	100%
2	5-C	388	100%
2	6-C	388	100%
2	7-C	388	100%
2	8-C	388	100%
2	9-C	388	100%



## 2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 13160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called PLEUROTOLYSIN A.

Mol	Chain	Residues	Atoms	AltConf	Trace
1	1-A	135	Total C 135 135	0	135
1	2-A	135	Total C 135 135	0	135
1	3-A	135	Total C 135 135	0	135
1	4-A	135	Total C 135 135	0	135
1	5-A	135	Total C 135 135	0	135
1	6-A	135	Total C 135 135	0	135
1	7-A	135	Total C 135 135	0	135
1	8-A	135	Total C 135 135	0	135
1	9-A	135	Total C 135 135	0	135
1	10-A	135	Total C 135 135	0	135
1	11-A	135	Total C 135 135	0	135
1	12-A	135	Total C 135 135	0	135
1	13-A	135	Total C 135 135	0	135
1	14-A	135	Total C 135 135	0	135
1	15-A	135	Total C 135 135	0	135
1	16-A	135	Total C 135 135	0	135
1	17-A	135	Total C 135 135	0	135



 $Continued\ from\ previous\ page...$ 

Mol	Chain	$oxed{ \mathbf{Residues} }$	Aton	ns	AltConf	Trace
1			Total	С	0	195
1	18-A	135	135	135	0	135
1	19-A	135	Total	С	0	135
1	19-A	155	135	135	U	100
1	20-A	135	Total	С	0	135
	20 11	100	135	135	Ü	100
1	1-B	135	Total	С	0	135
	1.5	100	135	135	Ü	100
1	2-B	135	Total	С	0	135
		100	135	135	Ü	
1	3-B	135	Total	С	0	135
			135	135	_	
1	4-B	135	Total	С	0	135
			135	135	_	
1	5-B	135	Total	С	0	135
			135	135		
1	6-B	135	Total	C	0	135
			135	135		
1	7-B	135	Total	C	0	135
			135	135		
1	8-B	135	Total	C	0	135
			135	135		
1	9-B	135	Total	C	0	135
			135	135		
1	10-B	135	Total 135	C	0	135
				135 C		
1	11-B	135	Total 135		0	135
				135		
1	12-B	135	Total 135	C 135	0	135
			Total	C		
1	13-B	135	135	135	0	135
			Total	C		
1	14-B	135	135	135	0	135
			Total	C		
1	15-B	135	135	135	0	135
			Total	C		
1	16-B	135	135	135	0	135
			Total	C		135
1	17-B	135	135	135	0	
			Total	C		
1	18-B	135	135	135	0	135
			100	100		



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	AltConf	Trace
1	19-B	135	Total C 135 135	0	135
1	20-B	135	Total C 135 135	0	135

### $\bullet$ Molecule 2 is a protein called PLEUROTOLYSIN B.

Mol	Chain	Residues	Atoms	AltConf	Trace
2	1-C	388	Total C 388 388	0	388
2	2-C	388	Total C 388 388	0	388
2	3-C	388	Total C 388 388	0	388
2	4-C	388	Total C 388 388	0	388
2	5-C	388	Total C 388 388	0	388
2	6-C	388	Total C 388 388	0	388
2	7-C	388	Total C 388 388	0	388
2	8-C	388	Total C 388 388	0	388
2	9-C	388	Total C 388 388	0	388
2	10-C	388	Total C 388 388	0	388
2	11-C	388	Total C 388 388	0	388
2	12-C	388	Total C 388 388	0	388
2	13-C	388	Total C 388 388	0	388
2	14-C	388	Total C 388 388	0	388
2	15-C	388	Total C 388 388	0	388
2	16-C	388	Total C 388 388	0	388
2	17-C	388	Total C 388 388	0	388



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms	AltConf	Trace
2	18-C	388	Total C 388 388	0	388
2	19-C	388	Total C 388 388	0	388
2	20-C	388	Total C 388 388	0	388

There are 87 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	?	-	GLU	deletion	UNP Q5W9E8
C	?	_	ALA	deletion	UNP Q5W9E8
С	?	_	GLU	deletion	UNP Q5W9E8
С	?	_	PHE	deletion	UNP Q5W9E8
С	?	_	THR	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	LEU	deletion	UNP Q5W9E8
С	?	-	MET	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	TYR	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	VAL	deletion	UNP Q5W9E8
С	?	-	LYS	deletion	UNP Q5W9E8
С	?	-	VAL	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	PRO	deletion	UNP Q5W9E8
С	?	-	PHE	deletion	UNP Q5W9E8
С	?	-	ILE	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	TYR	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	PHE	deletion	UNP Q5W9E8
С	?	-	LYS	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	ARG	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	VAL	deletion	UNP Q5W9E8
С	?	-	LYS	deletion	UNP Q5W9E8
С	?	-	GLN	deletion	UNP Q5W9E8
С	?	-	ASP	deletion	UNP Q5W9E8
С	?	-	VAL	deletion	UNP Q5W9E8
С	?	-	LYS	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	_	LEU	deletion	UNP Q5W9E8
С	?	-	GLU	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	VAL	deletion	UNP Q5W9E8
С	?	-	LYS	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	TRP	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	ALA	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	_	HIS	deletion	UNP Q5W9E8




### Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	GLN	deletion	UNP Q5W9E8
С	?	-	GLY	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	ILE	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	THR	deletion	UNP Q5W9E8
С	?	-	SER	deletion	UNP Q5W9E8
С	?	-	GLN	deletion	UNP Q5W9E8
С	?	-	ASN	deletion	UNP Q5W9E8
С	?	-	ARG	deletion	UNP Q5W9E8



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A



Clasies 4 A	
Chain 4-A:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 4-B:	100%
There are no outlier residues recorded for	
	tins cham.
• Molecule 1: PLEUROTOLYSIN A	
Chain 5-A:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 5-B:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 6-A:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
• Molecule 1. 1 Electro 1 Old 511 V A	
Chain 6-B:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Cl. : 7 A	
Chain 7-A:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 7-B:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 8-A:	100%
There are no outlier residues recorded for	this chain.
• Molecule 1: PLEUROTOLYSIN A	
Chain 8-B:	1000/
Спаш о-D.	100%



There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 9-A: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 9-B:
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 10-A: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 10-B: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 11-A: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 11-B: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 12-A: 100%
There are no outlier residues recorded for this chain.
• Molecule 1: PLEUROTOLYSIN A
Chain 12-B: 100%
There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 13-A: 100%

There are no outlier residues recorded for this chain.



• Molecule 1:	PLEUROTOLYSIN A
Chain 13-B:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 14-A:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 14-B:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 15-A:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 15-B:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 16-A:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 16-B:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 17-A:	100%
There are no o	outlier residues recorded for this chain.
• Molecule 1:	PLEUROTOLYSIN A
Chain 17-B:	100%

There are no outlier residues recorded for this chain.

 $\bullet$  Molecule 1: PLEUROTOLYSIN A



Chain 18-A: 100%

There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 18-B:

100%

There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 19-A:

100%

There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 19-B:

100%

There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 20-A:

100%

There are no outlier residues recorded for this chain.

• Molecule 1: PLEUROTOLYSIN A

Chain 20-B:

100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 1-C:

100%



• Molecule 2: PLEUROTOLYSIN B

Chain 2-C:

100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 3-C:

100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B



Chain 4-C:	
Cham 4-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 5-C:	100%
There are no outlier residues recorded for	
	oms cham.
• Molecule 2: PLEUROTOLYSIN B	
Chain 6-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 7-C:	1000
	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 8-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 9-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 10-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 11-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 12-C:	100%
There are no outlier residues recorded for	this chain.
• Molecule 2: PLEUROTOLYSIN B	
Chain 13-C:	100%



Thoma	0.70	10.0	antlian	rocidina	recorded	$f_{\alpha n}$	+hia	ahain
There	are	по	outher	residues	recorded	101	ums	cham.

• Molecule 2: PLEUROTOLYSIN B

Chain 14-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 15-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 16-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 17-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 18-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 19-C: 100%

There are no outlier residues recorded for this chain.

• Molecule 2: PLEUROTOLYSIN B

Chain 20-C: 100%

There are no outlier residues recorded for this chain.



# 4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C13	Depositor
Number of particles used	1150	Depositor
Resolution determination method	Not provided	
CTF correction method	ESTIMATED WITH CTFFIND3, THEN	Depositor
	PHASES FLIPPED FOR EACH PARTI-	
	CLE	
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	25	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	3600	Depositor
Magnification	76148	Depositor
Image detector	GATAN ULTRASCAN 4000 (4k x 4k)	Depositor
Maximum map value	0.002	Depositor
Minimum map value	-0.002	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.000	Depositor
Recommended contour level	0.0002	Depositor
Map size (Å)	400.0, 400.0, 400.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.0, 2.0, 2.0	Depositor



## 5 Model quality (i)

### 5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1-A	135	0	0	0	0
1	1-B	135	0	0	0	0
1	2-A	135	0	0	0	0
1	2-B	135	0	0	0	0
1	3-A	135	0	0	0	0
1	3-B	135	0	0	0	0
1	4-A	135	0	0	0	0
1	4-B	135	0	0	0	0
1	5-A	135	0	0	0	0
1	5-B	135	0	0	0	0
1	6-A	135	0	0	0	0
1	6-B	135	0	0	0	0
1	7-A	135	0	0	0	0
1	7-B	135	0	0	0	0
1	8-A	135	0	0	0	0
1	8-B	135	0	0	0	0
1	9-A	135	0	0	0	0
1	9-B	135	0	0	0	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	10-A	135	0	0	0	0
1	10-B	135	0	0	0	0
1	11-A	135	0	0	0	0
1	11-B	135	0	0	0	0
1	12-A	135	0	0	0	0
1	12-B	135	0	0	0	0
1	13-A	135	0	0	0	0
1	13-B	135	0	0	0	0
1	14-A	135	0	0	0	0
1	14-B	135	0	0	0	0
1	15-A	135	0	0	0	0
1	15-B	135	0	0	0	0
1	16-A	135	0	0	0	0
1	16-B	135	0	0	0	0
1	17-A	135	0	0	0	0
1	17-B	135	0	0	0	0
1	18-A	135	0	0	0	0
1	18-B	135	0	0	0	0
1	19-A	135	0	0	0	0
1	19-B	135	0	0	0	0
1	20-A	135	0	0	0	0
1	20-B	135	0	0	0	0
2	1-C	388	0	0	0	0
2	2-C	388	0	0	0	0
2	3-C	388	0	0	0	0
2	4-C	388	0	0	0	0
2	5-C	388	0	0	0	0
2	6-C	388	0	0	0	0
2	7-C	388	0	0	0	0
2	8-C	388	0	0	0	0
2	9-C	388	0	0	0	0
2	10-C	388	0	0	0	0
2	11-C	388	0	0	0	0
2	12-C	388	0	0	0	0
2	13-C	388	0	0	0	0
2	14-C	388	0	0	0	0
2	15-C	388	0	0	0	0
2	16-C	388	0	0	0	0
2	17-C	388	0	0	0	0
2	18-C	388	0	0	0	0
2	19-C	388	0	0	0	0
2	20-C	388	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	13160	0	0	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

There are no protein backbone outliers to report in this entry.

#### 5.3.2 Protein sidechains (i)

There are no protein residues with a non-rotameric sidechain to report in this entry.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

There are no ligands in this entry.

### 5.7 Other polymers (i)

There are no such residues in this entry.



## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




## 6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-2794. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections (i)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices (i)

#### 6.2.1 Primary map






The images above show central slices of the map in three orthogonal directions.

### 6.3 Largest variance slices (i)


#### 6.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) (i)

#### 6.4.1 Primary map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.



### 6.5 Orthogonal surface views (i)

#### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0002. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.6 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 7.2 Volume estimate (i)




The volume at the recommended contour level is  $2675~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $2417~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



## 7.3 Rotationally averaged power spectrum (i)



<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.067  $\rm \mathring{A}^{-1}$ 



# 8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.





## 9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-2794 and PDB model 4V3A. Per-residue inclusion information can be found in section 3 on page 11.

## 9.1 Map-model overlay (i)








The images above show the 3D surface view of the map at the recommended contour level 0.0002 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



### 9.2 Q-score mapped to coordinate model (i)

This section was not generated.


## 9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0002).



## 9.4 Atom inclusion (i)



At the recommended contour level, 93% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.



## 9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.0002) and Q-score for the entire model and for each chain.

Chain	Atom inclusion
All	0.9320
A	0.8810
В	0.9180
С	0.9540



