

wwPDB X-ray Structure Validation Summary Report (i)

Jun 16, 2024 – 08:41 AM EDT

PDB ID : 5EI8

Title: Rapid Discovery of Pyrido[3,4-d]pyrimidine Inhibitors of Monopolar Spindle

kinase 1 (MPS1) Using a Structure-Based Hydridization Approach

Authors: Innocenti, P.; Woodward, H.L.; Solanki, S.; Naud, N.; Westwood, I.M.; Cronin,

N.; Hayes, A.; Roberts, J.; Henley, A.T.; Baker, R.; Faisal, A.; Mak, G.; Box, G.; Valenti, M.; De Haven Brandon, A.; O'Fee, L.; Saville, J.; Schmitt, J.; Burke, R.; van Montfort, R.L.M.; Raymaud, F.I.; Eccles, S.A.; Linardopoulos,

S.; Blagg, J.; Hoelder, S.

Deposited on : 2015-10-29

Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$


CCP4 : 7.0.044 (Gargrove)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Whole archive Similar resolution Metric (#Entries) (#Entries, resolution range(Å)) R_{free} 6864 (2.20-2.16) 130704 Clashscore 141614 7689 (2.20-2.16) Ramachandran outliers 138981 7564 (2.20-2.16) Sidechain outliers 138945 7564 (2.20-2.16) RSRZ outliers 127900 6738 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			7%		
1	A	313	72%	11%	17%

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.37.1



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 2123 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dual specificity protein kinase TTK.

Mol	Chain	Residues		Atoms			ZeroOcc	AltConf	Trace	
1	Λ	261	Total	С	N	О	S	0	0	0
1	Α	201	2040	1317	333	377	13	0	0	U

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	496	MET	-	initiating methionine	UNP P33981
A	497	HIS	-	expression tag	UNP P33981
A	498	HIS	-	expression tag	UNP P33981
A	499	HIS	-	expression tag	UNP P33981
A	500	HIS	-	expression tag	UNP P33981
A	501	HIS	-	expression tag	UNP P33981
A	502	HIS	-	expression tag	UNP P33981
A	503	SER	-	expression tag	UNP P33981
A	504	SER	-	expression tag	UNP P33981
A	505	GLY	-	expression tag	UNP P33981
A	506	VAL	-	expression tag	UNP P33981
A	507	ASP	-	expression tag	UNP P33981
A	508	LEU	-	expression tag	UNP P33981
A	509	GLY	-	expression tag	UNP P33981
A	510	THR	-	expression tag	UNP P33981
A	511	GLU	-	expression tag	UNP P33981
A	512	ASN	-	expression tag	UNP P33981
A	513	LEU	-	expression tag	UNP P33981
A	514	TYR	-	expression tag	UNP P33981
A	515	PHE	-	expression tag	UNP P33981
A	516	GLN	-	expression tag	UNP P33981
A	517	SER	-	expression tag	UNP P33981
A	518	MET	-	expression tag	UNP P33981
A	754	ILE	-	expression tag	UNP P33981
A	755	GLU	-	expression tag	UNP P33981
A	756	PHE	-	expression tag	UNP P33981
A	757	PRO	-	expression tag	UNP P33981

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	758	ASP	-	expression tag	UNP P33981
A	759	ILE	-	expression tag	UNP P33981
A	760	PRO	-	expression tag	UNP P33981
A	761	GLU	-	expression tag	UNP P33981
A	762	LYS	-	expression tag	UNP P33981
A	763	ASP	-	expression tag	UNP P33981
A	764	LEU	-	expression tag	UNP P33981
A	765	GLN	-	expression tag	UNP P33981
A	766	ASP	-	expression tag	UNP P33981
A	767	VAL	-	expression tag	UNP P33981
A	768	LEU	-	expression tag	UNP P33981
A	769	LYS	-	expression tag	UNP P33981
A	770	CYS	-	expression tag	UNP P33981
A	771	CYS	-	expression tag	UNP P33981
A	772	LEU	-	expression tag	UNP P33981
A	773	LYS	-	expression tag	UNP P33981
A	774	ARG	-	expression tag	UNP P33981
A	775	ASP	-	expression tag	UNP P33981
A	776	PRO	-	expression tag	UNP P33981
A	777	LYS	-	expression tag	UNP P33981
A	778	GLN	-	expression tag	UNP P33981
A	779	ARG	-	expression tag	UNP P33981
A	780	ILE	-	expression tag	UNP P33981
A	781	SER	-	expression tag	UNP P33981
A	782	ILE	-	expression tag	UNP P33981
A	783	PRO	-	expression tag	UNP P33981
A	784	GLU	-	expression tag	UNP P33981
A	785	LEU	-	expression tag	UNP P33981
A	786	LEU	-	expression tag	UNP P33981
A	787	ALA	-	expression tag	UNP P33981
A	788	HIS	-	expression tag	UNP P33981
A	789	PRO	-	expression tag	UNP P33981
A	790	TYR	-	expression tag	UNP P33981
A	791	VAL	-	expression tag	UNP P33981
A	792	GLN	-	expression tag	UNP P33981
A	793	ILE	-	expression tag	UNP P33981
A	794	GLN	-	expression tag	UNP P33981
A	795	THR	-	expression tag	UNP P33981
A	796	HIS	-	expression tag	UNP P33981
A	797	PRO	-	expression tag	UNP P33981
A	798	VAL	-	expression tag	UNP P33981
A	799	ASN	-	expression tag	UNP P33981


Continued on next page...



n previous	paae
	n previous

Chain	Residue	Modelled	Actual	Comment	Reference
A	800	GLN	-	expression tag	UNP P33981
A	801	MET	-	expression tag	UNP P33981
A	802	ALA	-	expression tag	UNP P33981
A	803	LYS	-	expression tag	UNP P33981
A	804	GLY	-	expression tag	UNP P33981
A	805	THR	-	expression tag	UNP P33981
A	806	THR	-	expression tag	UNP P33981
A	807	GLU	_	expression tag	UNP P33981
A	808	GLU	-	expression tag	UNP P33981


• Molecule 2 is $\{N\}$ -[2-methoxy-4-(1-methylpyrazol-4-yl)phenyl]-8-(1-methylpyrazol-4-yl)pyr ido[3,4-d]pyrimidin-2-amine (three-letter code: 5OE) (formula: $C_{22}H_{20}N_8O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 31	C 22	N 8	O 1	0	0

• Molecule 3 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C_2H_6OS).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O S 4 2 1 1	0	0
3	A	1	Total C O S 4 2 1 1	0	0
3	A	1	Total C O S 4 2 1 1	0	0

• Molecule 4 is water.


\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	40	Total O 40 40	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dual specificity protein kinase TTK





4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	70.00Å 101.98Å 110.87Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.53 - 2.17	Depositor
resolution (A)	37.53 - 2.18	EDS
% Data completeness	99.2 (37.53-2.17)	Depositor
(in resolution range)	99.2 (37.53-2.18)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.19 (at 2.18Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.214 , 0.244	Depositor
it, it free	0.249 , 0.277	DCC
R_{free} test set	1092 reflections (5.22%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtriage
Anisotropy	0.403	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 62.1	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2123	wwPDB-VP
Average B, all atoms $(Å^2)$	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.24% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, 5OE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		$\mathbf{lengths}$	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.45	0/2086	0.61	0/2835	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2040	0	1952	31	0
2	A	31	0	0	1	0
3	A	12	0	18	0	0
4	A	40	0	0	1	0
All	All	2123	0	1970	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)	
1:A:574:TYR:CD1	1:A:643:ILE:HD11	2.00	0.96	
1:A:788:HIS:ND1	1:A:789:PRO:HD2	1.87	0.90	
1:A:738:ILE:H	1:A:738:ILE:HD12	1.41	0.84	
1:A:788:HIS:CE1	1:A:789:PRO:HD2	2.18	0.78	
1:A:530:GLN:NE2	1:A:674:ASP:OD2	2.15	0.78	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/313 (82%)	249 (98%)	6 (2%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	213/288 (74%)	213 (100%)	0	100	100	

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	pe Chain Ro	Res	Link	Bond lengths			Bond angles		
MIOI	туре		nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	5OE	A	901	-	28,35,35	0.76	1 (3%)	38,50,50	0.83	2 (5%)
3	DMS	A	902	-	3,3,3	0.32	0	3,3,3	0.32	0
3	DMS	A	904	-	3,3,3	0.30	0	3,3,3	0.23	0
3	DMS	A	903	-	3,3,3	0.26	0	3,3,3	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5OE	A	901	-	-	0/10/14/14	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}({ ext{ iny A}})$
2	A	901	5OE	C18-C17	-3.27	1.44	1.49



All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	A	901	5OE	C15-C1-C2	2.34	126.82	122.63
2	A	901	5OE	C16-N5-C17	2.20	119.58	117.99

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	901	5OE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$		$OWAB(A^2)$	Q < 0.9
1	A	261/313 (83%)	0.48	21 (8%) 12	12	40, 62, 97, 120	0

The worst 5 of 21 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	622	TRP	4.6
1	A	751	ASN	4.3
1	A	560	ALA	4.3
1	A	674	ASP	3.8
1	A	536	SER	3.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	DMS	A	903	4/4	0.79	0.28	110,110,110,110	0
3	DMS	A	902	4/4	0.87	0.29	105,106,106,107	0


Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	DMS	A	904	4/4	0.90	0.34	113,113,113,113	0
2	5OE	A	901	31/31	0.93	0.12	36,43,57,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers (i)

There are no such residues in this entry.

