



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 01:37 PM EDT

PDB ID : 5ENO  
Title : MBX2319 bound structure of bacterial efflux pump.  
Authors : Sjuts, H.; Ornik, A.R.; Pos, K.M.  
Deposited on : 2015-11-09  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	609	 3% 90% 5% 5%
1	B	609	 % 89% 5% 5%
1	C	609	 2% 90% 5% 5%
2	D	169	 4% 91% 5% .
2	E	169	 5% 88% 5% 8%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	169	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '7%', a large green segment in the middle labeled '91%', and a yellow segment on the right labeled '5%'. A small grey dot is visible at the end of the bar.</p>

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 17885 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multidrug efflux pump subunit AcrB, Multidrug efflux pump subunit AcrB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	580	4419	2774	749	874	22	0	0	0
1	B	578	4405	2766	746	871	22	0	0	0
1	C	577	4396	2761	744	869	22	0	0	0

There are 27 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	552	GLY	-	linker	UNP P31224
A	553	GLY	-	linker	UNP P31224
A	554	SER	-	linker	UNP P31224
A	555	GLY	-	linker	UNP P31224
A	556	GLY	-	linker	UNP P31224
A	557	SER	-	linker	UNP P31224
A	558	GLY	-	linker	UNP P31224
A	559	GLY	-	linker	UNP P31224
A	560	SER	-	linker	UNP P31224
B	552	GLY	-	linker	UNP P31224
B	553	GLY	-	linker	UNP P31224
B	554	SER	-	linker	UNP P31224
B	555	GLY	-	linker	UNP P31224
B	556	GLY	-	linker	UNP P31224
B	557	SER	-	linker	UNP P31224
B	558	GLY	-	linker	UNP P31224
B	559	GLY	-	linker	UNP P31224
B	560	SER	-	linker	UNP P31224
C	552	GLY	-	linker	UNP P31224
C	553	GLY	-	linker	UNP P31224
C	554	SER	-	linker	UNP P31224
C	555	GLY	-	linker	UNP P31224

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	556	GLY	-	linker	UNP P31224
C	557	SER	-	linker	UNP P31224
C	558	GLY	-	linker	UNP P31224
C	559	GLY	-	linker	UNP P31224
C	560	SER	-	linker	UNP P31224

- Molecule 2 is a protein called DARPin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	162	Total	C	N	O	S	0	0	0
			1237	777	224	235	1			
2	E	156	Total	C	N	O	S	0	0	0
			1177	741	206	229	1			
2	F	162	Total	C	N	O	S	0	0	0
			1237	777	224	235	1			

- Molecule 3 is 3,3-dimethyl-8-morpholin-4-yl-6-(2-phenylethylsulfanyl)-1,4-dihydropyrido[3,4-c]pyridine-5-carbonitrile (three-letter code: 5QG) (formula: C<sub>23</sub>H<sub>27</sub>N<sub>3</sub>O<sub>2</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	C	1	Total	C	N	O	S	0	0
			29	23	3	2	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	292	Total 292	O 292	0	0
4	B	309	Total 309	O 309	0	0
4	C	294	Total 294	O 294	0	0
4	D	36	Total 36	O 36	0	0
4	E	27	Total 27	O 27	0	0
4	F	27	Total 27	O 27	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multidrug efflux pump subunit AcrB, Multidrug efflux pump subunit AcrB




- Molecule 1: Multidrug efflux pump subunit AcrB, Multidrug efflux pump subunit AcrB




- Molecule 1: Multidrug efflux pump subunit AcrB, Multidrug efflux pump subunit AcrB





- Molecule 2: DARPin






- Molecule 2: DARPin



- Molecule 2: DARPin





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	108.76Å 145.15Å 174.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.20 49.62 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-2.20) 99.7 (49.62-2.20)	Depositor EDS
$R_{merge}$	0.31	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.68 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.199 , 0.247 0.205 , 0.249	Depositor DCC
$R_{free}$ test set	6897 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.5	Xtrriage
Anisotropy	0.431	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 29.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	17885	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 5QG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.87	3/4494 (0.1%)	0.89	6/6089 (0.1%)
1	B	0.87	2/4480 (0.0%)	0.96	15/6070 (0.2%)
1	C	0.82	0/4471	0.86	7/6058 (0.1%)
2	D	0.76	0/1262	0.80	1/1716 (0.1%)
2	E	0.76	0/1196	0.86	0/1626
2	F	0.76	0/1262	0.81	2/1716 (0.1%)
All	All	0.83	5/17165 (0.0%)	0.89	31/23275 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	99	ASP	CB-CG	5.75	1.63	1.51
1	A	634	TRP	CB-CG	5.67	1.60	1.50
1	A	84	SER	CB-OG	5.61	1.49	1.42
1	B	84	SER	CB-OG	5.44	1.49	1.42
1	A	776	GLU	CD-OE1	-5.10	1.20	1.25

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	99	ASP	CB-CG-OD2	16.80	133.42	118.30
1	B	765	ARG	NE-CZ-NH1	9.95	125.28	120.30
1	B	765	ARG	NE-CZ-NH2	-8.07	116.27	120.30
1	B	99	ASP	CB-CG-OD1	-7.19	111.83	118.30
1	C	815	ARG	NE-CZ-NH1	7.17	123.88	120.30
1	A	765	ARG	NE-CZ-NH1	7.03	123.81	120.30
1	A	168	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	B	586	ARG	NE-CZ-NH1	6.14	123.37	120.30
1	C	239	ARG	NE-CZ-NH2	-6.14	117.23	120.30
2	F	77	ASP	CB-CG-OD1	6.05	123.74	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	765	ARG	NE-CZ-NH2	-5.97	117.32	120.30
1	B	815	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	C	815	ARG	NE-CZ-NH2	-5.92	117.34	120.30
1	B	815	ARG	NE-CZ-NH2	-5.81	117.39	120.30
1	B	653	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	B	653	ARG	NE-CZ-NH1	5.71	123.15	120.30
1	A	767	ARG	NE-CZ-NH2	5.69	123.14	120.30
1	B	792	ARG	NE-CZ-NH1	5.66	123.13	120.30
1	B	792	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	B	586	ARG	NE-CZ-NH2	-5.37	117.62	120.30
1	C	761	ASP	CB-CG-OD2	-5.34	113.49	118.30
1	A	767	ARG	NE-CZ-NH1	-5.24	117.68	120.30
1	B	307	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	C	761	ASP	CB-CG-OD1	5.18	122.96	118.30
1	B	795	ASP	CB-CG-OD1	5.12	122.90	118.30
2	F	44	ASP	CB-CG-OD2	-5.10	113.71	118.30
2	D	72	ASP	CB-CG-OD1	5.09	122.88	118.30
1	B	808	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	A	620	ARG	NE-CZ-NH2	-5.09	117.76	120.30
1	C	808	ARG	NE-CZ-NH1	5.07	122.84	120.30
1	C	780	ARG	NE-CZ-NH1	5.03	122.82	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4419	0	4356	11	0
1	B	4405	0	4343	14	0
1	C	4396	0	4335	14	0
2	D	1237	0	1201	3	0
2	E	1177	0	1159	3	0
2	F	1237	0	1201	1	0
3	C	29	0	0	2	0
4	A	292	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	309	0	0	2	0
4	C	294	0	0	3	0
4	D	36	0	0	0	0
4	E	27	0	0	0	0
4	F	27	0	0	1	0
All	All	17885	0	16595	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:667:ASN:O	1:A:668:LEU:HG	1.92	0.70
1:A:733:GLN:OE1	1:A:743:ILE:HD11	1.96	0.65
1:A:743:ILE:HD11	4:A:945:HOH:O	1.98	0.63
1:B:238:THR:HG21	4:B:1204:HOH:O	2.00	0.61
1:C:70:ASN:HB3	4:C:1081:HOH:O	2.02	0.60
1:B:219:LEU:HD23	1:C:754:TRP:CZ3	2.39	0.57
2:D:153:SER:HB3	2:D:161:LEU:HD23	1.89	0.55
1:A:298:ASN:HD22	1:A:301:ASP:H	1.55	0.53
1:B:238:THR:CG2	4:B:1204:HOH:O	2.56	0.53
1:C:617:PHE:N	4:C:1005:HOH:O	2.41	0.53
1:B:115:MET:CE	1:C:116:PRO:HG2	2.40	0.51
1:B:213:GLN:HE22	1:B:238:THR:HG22	1.76	0.51
2:D:42:ALA:O	2:D:50:PRO:HD3	2.12	0.50
1:B:247:GLY:HA2	1:B:268:ILE:HD13	1.93	0.49
1:A:41:PRO:HB3	1:A:295:THR:HG21	1.95	0.49
1:B:775:SER:HB3	1:B:780:ARG:HD3	1.94	0.49
2:E:51:LEU:HD11	2:E:63:VAL:HG13	1.95	0.49
1:C:705:GLU:HB3	1:C:847:LEU:HD22	1.94	0.49
1:C:714:THR:HG22	4:C:1164:HOH:O	2.13	0.48
1:B:764:ASP:OD1	1:B:765:ARG:HD3	2.14	0.47
1:C:100:ALA:HB1	1:C:131:LYS:HD2	1.97	0.47
1:C:178:PHE:HB3	3:C:901:5QG:C15	2.45	0.47
1:B:153:ASP:OD1	1:B:182:TYR:OH	2.34	0.46
1:A:223:PRO:HD3	1:B:275:TYR:CD1	2.50	0.46
1:B:111:LEU:HD21	1:B:127:VAL:HG11	1.96	0.46
2:F:80:GLY:O	2:F:110:ASP:HA	2.16	0.45
1:A:109:ASN:CG	1:C:108:GLN:HG2	2.36	0.45
1:C:178:PHE:HB3	3:C:901:5QG:C14	2.48	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:42:ALA:O	2:E:50:PRO:HD3	2.17	0.44
1:A:578:LEU:HD12	1:A:578:LEU:N	2.32	0.43
1:B:777:ALA:O	1:B:781:MET:HG2	2.19	0.42
1:A:247:GLY:HA2	1:A:268:ILE:CD1	2.49	0.42
1:C:259:ARG:HD2	4:F:202:HOH:O	2.18	0.42
1:A:667:ASN:O	1:A:667:ASN:CG	2.57	0.42
1:C:185:ARG:HD3	1:C:272:GLY:O	2.20	0.42
1:C:777:ALA:O	1:C:781:MET:HG2	2.20	0.42
1:B:763:ILE:HD11	1:C:59:ASP:HB3	2.02	0.41
1:A:713:LEU:HG	1:A:843:LEU:HD23	2.03	0.41
2:D:51:LEU:HD11	2:D:63:VAL:HG13	2.02	0.41
1:B:247:GLY:HA2	1:B:268:ILE:CD1	2.51	0.41
2:E:105:ASP:C	2:E:105:ASP:OD1	2.58	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	574/609 (94%)	555 (97%)	18 (3%)	1 (0%)	47	55
1	B	572/609 (94%)	554 (97%)	16 (3%)	2 (0%)	41	46
1	C	571/609 (94%)	556 (97%)	15 (3%)	0	100	100
2	D	160/169 (95%)	156 (98%)	3 (2%)	1 (1%)	25	26
2	E	154/169 (91%)	151 (98%)	3 (2%)	0	100	100
2	F	160/169 (95%)	156 (98%)	3 (2%)	1 (1%)	25	26
All	All	2191/2334 (94%)	2128 (97%)	58 (3%)	5 (0%)	47	55

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	633	ASP
1	B	147	GLY
2	F	9	HIS
2	D	12	SER
1	B	617	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	471/492 (96%)	464 (98%)	7 (2%)	65	78
1	B	470/492 (96%)	465 (99%)	5 (1%)	73	85
1	C	469/492 (95%)	460 (98%)	9 (2%)	57	71
2	D	126/132 (96%)	125 (99%)	1 (1%)	81	90
2	E	120/132 (91%)	117 (98%)	3 (2%)	47	60
2	F	126/132 (96%)	122 (97%)	4 (3%)	39	50
All	All	1782/1872 (95%)	1753 (98%)	29 (2%)	62	76

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	TYR
1	A	133	SER
1	A	151	GLN
1	A	314	GLU
1	A	575	MET
1	A	715	SER
1	A	801	PHE
1	B	49	TYR
1	B	148	THR
1	B	300	LEU
1	B	586	ARG
1	B	801	PHE
1	C	40	PRO
1	C	49	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	135	SER
1	C	256	ASP
1	C	274	ASN
1	C	610	PHE
1	C	630	SER
1	C	645	GLU
1	C	801	PHE
2	D	23	ARG
2	E	97	GLU
2	E	134	LYS
2	E	139	VAL
2	F	20	GLU
2	F	27	ASP
2	F	112	ASN
2	F	139	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	284	GLN
1	A	298	ASN
1	A	622	GLN
1	A	687	GLN
1	B	70	ASN
1	B	74	ASN
1	C	81	ASN
1	C	284	GLN
2	D	5	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	5QG	C	901	-	32,32,32	2.04	6 (18%)	38,45,45	2.05	6 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5QG	C	901	-	-	0/11/31/31	0/4/4/4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	901	5QG	C2-C1	-7.54	1.30	1.44
3	C	901	5QG	O8-C30	-5.15	1.41	1.45
3	C	901	5QG	C7-S8	-2.97	1.72	1.76
3	C	901	5QG	C5-N17	2.79	1.44	1.37
3	C	901	5QG	C7-N6	2.67	1.36	1.33
3	C	901	5QG	C6-C3	2.13	1.53	1.51

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	901	5QG	C9-S8-C7	5.74	110.87	101.82
3	C	901	5QG	C2-C7-S8	5.45	123.84	117.75
3	C	901	5QG	C2-C7-N6	-5.34	119.07	123.12

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	901	5QG	C4-C5-N6	-4.06	116.35	122.61
3	C	901	5QG	C5-N6-C7	3.55	125.88	116.30
3	C	901	5QG	C3-C2-C1	2.76	123.18	119.57

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	901	5QG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	580/609 (95%)	-0.15	17 (2%) 51 49	14, 26, 61, 94	0
1	B	578/609 (94%)	-0.31	6 (1%) 82 81	14, 24, 53, 128	0
1	C	577/609 (94%)	-0.18	15 (2%) 56 53	12, 27, 65, 88	0
2	D	162/169 (95%)	-0.08	6 (3%) 41 39	21, 35, 67, 113	0
2	E	156/169 (92%)	0.14	8 (5%) 28 26	22, 36, 71, 109	0
2	F	162/169 (95%)	0.33	12 (7%) 14 13	23, 40, 88, 131	0
All	All	2215/2334 (94%)	-0.14	64 (2%) 51 49	12, 28, 64, 131	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	11	GLY	15.5
1	B	617	PHE	9.5
2	F	8	HIS	8.3
2	F	11	GLY	7.4
2	F	5	HIS	5.3
2	D	11	GLY	4.9
2	E	31	ARG	4.9
2	E	12	SER	4.6
2	F	9	HIS	4.2
1	A	837	THR	4.1
1	A	39	ALA	4.1
1	A	677	ALA	4.0
2	F	10	HIS	3.9
1	A	831	ALA	3.9
2	D	28	ASP	3.8
1	A	255	GLN	3.8
1	C	678	THR	3.8
2	D	12	SER	3.7
1	C	635	ALA	3.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	F	13	ASP	3.4
1	A	634	TRP	3.4
1	C	133	SER	3.3
2	E	34	MET	3.2
2	F	31	ARG	3.2
1	B	833	PRO	3.2
1	C	255	GLN	3.2
1	A	834	GLY	3.2
1	C	634	TRP	3.1
1	A	865	GLN	3.1
2	E	35	ALA	3.1
2	F	6	HIS	3.1
2	F	163	GLU	3.1
1	A	678	THR	3.0
1	A	617	PHE	3.0
2	F	12	SER	2.9
1	A	836	SER	2.9
2	D	31	ARG	2.9
1	A	257	GLY	2.8
1	C	638	PRO	2.8
1	B	255	GLN	2.8
1	A	569	GLN	2.8
1	C	831	ALA	2.6
1	B	711	ASP	2.6
1	C	170	SER	2.6
2	E	28	ASP	2.6
1	C	299	ALA	2.5
2	F	28	ASP	2.5
2	D	10	HIS	2.4
1	C	834	GLY	2.4
1	A	636	ASP	2.3
1	C	257	GLY	2.2
2	F	76	TYR	2.2
1	C	833	PRO	2.2
1	C	837	THR	2.2
1	B	712	MET	2.2
2	D	35	ALA	2.2
1	C	640	GLU	2.2
2	E	163	GLU	2.1
1	B	831	ALA	2.1
1	A	633	ASP	2.1
1	A	832	ALA	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	E	30	VAL	2.1
1	A	640	GLU	2.0
1	C	618	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	5QG	C	901	29/29	0.90	0.17	39,51,63,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.