



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 17, 2021 – 06:46 PM EDT

PDB ID : 5RP7
Title : PanDDA analysis group deposition – Proteinase K changed state model for fragment Frag Xtal Screen D12a
Authors : Lima, G.M.A.; Talibov, V.; Benz, L.S.; Jagudin, E.; Mueller, U.
Deposited on : 2020-09-23
Resolution : 1.15 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.20
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.20


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1492 (1.18-1.10)
Clashscore	141614	1537 (1.18-1.10)
Ramachandran outliers	138981	1483 (1.18-1.10)
Sidechain outliers	138945	1480 (1.18-1.10)
RSRZ outliers	127900	1464 (1.18-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	F91	A	1002[L]	-	-	-	X

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2362 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proteinase K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	279	2023	1244	355	414	10	0	17	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	207	ASP	SER	conflict	UNP P06873

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

- Molecule 3 is N-(pyridin-4-ylmethyl)-2,3-dihydro-1,4-benzodioxin-6-amine (three-letter code: F91) (formula: C₁₄H₁₄N₂O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	18	14	2	2	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	316	316	316	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proteinase K



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	68.04Å 68.04Å 102.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.11 – 1.15 48.11 – 1.15	Depositor EDS
% Data completeness (in resolution range)	99.0 (48.11-1.15) 99.0 (48.11-1.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.15Å)	Xtrriage
Refinement program	PHENIX 1.19.1	Depositor
R, R_{free}	0.186 , 0.197 0.188 , 0.198	Depositor DCC
R_{free} test set	4187 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	11.7	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2362	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, F91

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2062	0.61	0/2804

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2023	0	1900	4	0
2	A	5	0	0	0	0
3	A	18	0	14	0	0
4	A	316	0	0	1	2
All	All	2362	0	1914	4	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ILE:HD12	1:A:131[A]:LEU:HD11	1.92	0.52
1:A:48:GLU:HB3	1:A:79:SER:HB2	2.01	0.42
1:A:250[A]:ARG:NH1	4:A:1104:HOH:O	2.40	0.42
1:A:212:TRP:HB3	1:A:216[A]:SER:OG	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1295:HOH:O	4:A:1295:HOH:O[8_665]	1.07	1.13
4:A:1329:HOH:O	4:A:1385:HOH:O[3_644]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	277/279 (99%)	269 (97%)	8 (3%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/213 (99%)	207 (99%)	3 (1%)	67 29

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	TYR
1	A	154[A]	MET
1	A	225	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	F91	A	1002[L]	-	20,20,20	2.42	7 (35%)	26,26,26	1.98	6 (23%)
2	SO4	A	1001	-	4,4,4	0.18	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	F91	A	1002[L]	-	-	1/5/12/12	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1002[L]	F91	O2-C1	6.43	1.44	1.37
3	A	1002[L]	F91	O5-C6	5.48	1.43	1.37
3	A	1002[L]	F91	C9-N10	2.90	1.47	1.38
3	A	1002[L]	F91	O2-C3	-2.82	1.36	1.43
3	A	1002[L]	F91	O5-C4	-2.78	1.36	1.43
3	A	1002[L]	F91	C11-C12	2.24	1.56	1.51
3	A	1002[L]	F91	C8-C9	2.23	1.43	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1002[L]	F91	O5-C6-C1	-4.44	118.05	122.03
3	A	1002[L]	F91	C3-O2-C1	4.30	121.06	113.65
3	A	1002[L]	F91	O2-C1-C6	-4.22	118.25	122.03
3	A	1002[L]	F91	C12-C11-N10	-3.22	105.39	113.77
3	A	1002[L]	F91	O2-C1-C13	3.15	121.90	117.05
3	A	1002[L]	F91	O5-C6-C7	2.95	121.58	116.85

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1002[L]	F91	C12-C11-N10-C9

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/279 (100%)	2.85	226 (81%) 0 0	8, 11, 17, 28	12 (4%)

All (226) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	ALA	12.8
1	A	198	VAL	6.3
1	A	64	ARG	5.9
1	A	34[A]	CYS	5.7
1	A	13[A]	ILE	5.3
1	A	278	GLN	4.9
1	A	201	ILE	4.8
1	A	131[A]	LEU	4.8
1	A	213	ILE	4.6
1	A	252	ILE	4.6
1	A	237	LEU	4.5
1	A	107	ILE	4.5
1	A	90	LEU	4.5
1	A	104	TYR	4.4
1	A	93	VAL	4.4
1	A	5	ASN	4.4
1	A	82	TYR	4.3
1	A	180	VAL	4.2
1	A	36	TYR	4.2
1	A	266	PHE	4.2
1	A	84	VAL	4.2
1	A	60	TYR	4.2
1	A	212	TRP	4.2
1	A	95	VAL	4.2
1	A	77	VAL	4.1
1	A	251	TYR	4.1
1	A	169	TYR	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	99	ASN	4.0
1	A	56	VAL	4.0
1	A	272	LEU	4.0
1	A	233	LEU	4.0
1	A	73	CYS	4.0
1	A	106	THR	4.0
1	A	261	LEU	3.9
1	A	89	GLN	3.9
1	A	1	ALA	3.9
1	A	178	CYS	3.9
1	A	192	PHE	3.9
1	A	114	VAL	3.9
1	A	127	VAL	3.9
1	A	155	VAL	3.8
1	A	35	VAL	3.8
1	A	240	LEU	3.7
1	A	31[A]	GLN	3.7
1	A	59	TYR	3.7
1	A	38	ILE	3.7
1	A	238[A]	MET	3.6
1	A	21	SER	3.6
1	A	128	VAL	3.6
1	A	42	ILE	3.6
1	A	81	THR	3.6
1	A	2	ALA	3.6
1	A	8	TRP	3.6
1	A	133	LEU	3.6
1	A	271	LEU	3.6
1	A	113	PHE	3.6
1	A	148	LEU	3.6
1	A	208	ILE	3.5
1	A	61	TYR	3.5
1	A	202	PHE	3.5
1	A	186	TYR	3.5
1	A	24	TYR	3.5
1	A	112	ASP	3.5
1	A	10	LEU	3.5
1	A	11	ALA	3.4
1	A	245	ALA	3.4
1	A	126	GLY	3.4
1	A	153	VAL	3.4
1	A	269	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	91[A]	PHE	3.4
1	A	16	THR	3.4
1	A	76	THR	3.4
1	A	129	ALA	3.4
1	A	182	ALA	3.4
1	A	220	ILE	3.4
1	A	37	VAL	3.4
1	A	177	VAL	3.4
1	A	49	PHE	3.4
1	A	23	TYR	3.3
1	A	195	TYR	3.3
1	A	236	TYR	3.3
1	A	206	THR	3.3
1	A	158	ALA	3.3
1	A	235	ALA	3.3
1	A	100	GLY	3.3
1	A	101	SER	3.2
1	A	102	GLY	3.2
1	A	123	CYS	3.2
1	A	115	ALA	3.2
1	A	196	GLY	3.2
1	A	204	PRO	3.2
1	A	74	ALA	3.1
1	A	18	PRO	3.1
1	A	141	VAL	3.1
1	A	164	ALA	3.1
1	A	103	GLN	3.1
1	A	20	THR	3.1
1	A	105	SER	3.1
1	A	85	ALA	3.1
1	A	258	LYS	3.1
1	A	249	CYS	3.1
1	A	25	TYR	3.1
1	A	39	ASP	3.0
1	A	156	ALA	3.0
1	A	253	ALA	3.0
1	A	256	ALA	3.0
1	A	234	ALA	3.0
1	A	231	ALA	2.9
1	A	222	GLY	2.9
1	A	221	SER	2.9
1	A	54[A]	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	137	TYR	2.9
1	A	29	ALA	2.9
1	A	244	THR	2.9
1	A	157	VAL	2.9
1	A	230	VAL	2.9
1	A	108	ILE	2.9
1	A	88	THR	2.9
1	A	6	ALA	2.9
1	A	44	ALA	2.9
1	A	209	LEU	2.8
1	A	32	GLY	2.8
1	A	58	THR	2.8
1	A	179	THR	2.8
1	A	211	THR	2.8
1	A	223	THR	2.8
1	A	96	LEU	2.8
1	A	219[A]	SER	2.8
1	A	68	GLY	2.8
1	A	22	THR	2.8
1	A	224	SER	2.8
1	A	152	GLY	2.8
1	A	274	TYR	2.8
1	A	109	ALA	2.7
1	A	163	ASN	2.7
1	A	273	ALA	2.7
1	A	255	THR	2.7
1	A	78	GLY	2.7
1	A	199	LEU	2.7
1	A	30	GLY	2.7
1	A	181	GLY	2.7
1	A	277	TYR	2.7
1	A	7	PRO	2.6
1	A	171	PRO	2.6
1	A	66	GLY	2.6
1	A	203	GLY	2.6
1	A	210	SER	2.6
1	A	226	ALA	2.6
1	A	161[A]	ASN	2.6
1	A	110	GLY	2.6
1	A	159	ALA	2.6
1	A	75	GLY	2.6
1	A	184	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	200	ASP	2.6
1	A	227	THR	2.6
1	A	140[A]	SER	2.6
1	A	53	ALA	2.6
1	A	67	ASN	2.6
1	A	189	ARG	2.5
1	A	216[A]	SER	2.5
1	A	14	SER	2.5
1	A	92	GLY	2.5
1	A	130	SER	2.5
1	A	132	SER	2.5
1	A	264	ILE	2.5
1	A	160	GLY	2.5
1	A	167	ARG	2.5
1	A	217	THR	2.5
1	A	225	MET	2.5
1	A	146	ALA	2.4
1	A	172	ALA	2.4
1	A	248	ALA	2.4
1	A	154[A]	MET	2.4
1	A	116	SER	2.4
1	A	4	THR	2.4
1	A	239	THR	2.4
1	A	268	THR	2.4
1	A	19	GLY	2.4
1	A	214	GLY	2.4
1	A	257	ASN	2.4
1	A	9	GLY	2.4
1	A	205	GLY	2.4
1	A	122	ASN	2.3
1	A	276	ASN	2.3
1	A	183	SER	2.3
1	A	175	PRO	2.3
1	A	228	PRO	2.3
1	A	254	ASP	2.3
1	A	72	HIS	2.3
1	A	70	GLY	2.3
1	A	259	GLY	2.3
1	A	71	THR	2.3
1	A	215	GLY	2.3
1	A	125	LYS	2.3
1	A	143[A]	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	ALA	2.2
1	A	98	ASP	2.2
1	A	162	ASN	2.2
1	A	166	ALA	2.2
1	A	262	SER	2.2
1	A	144	ALA	2.2
1	A	120	ASN	2.2
1	A	270	ASN	2.2
1	A	51	GLY	2.2
1	A	55	MET	2.2
1	A	260	ASP	2.2
1	A	242[A]	LYS	2.1
1	A	63	SER	2.1
1	A	150	SER	2.1
1	A	40	THR	2.1
1	A	168	ASN	2.1
1	A	33	SER	2.1
1	A	165	ASP	2.1
1	A	17	SER	2.1
1	A	232	GLY	2.1
1	A	46	HIS	2.1
1	A	124	PRO	2.1
1	A	241	GLY	2.1
1	A	3	GLN	2.1
1	A	83	GLY	2.0
1	A	47	PRO	2.0
1	A	111	MET	2.0
1	A	134	GLY	2.0
1	A	229	HIS	2.0
1	A	250[A]	ARG	2.0
1	A	275	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	F91	A	1002[L]	18/18	0.06	0.74	16,25,34,34	18
2	SO4	A	1001	5/5	0.90	0.15	11,13,14,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.