



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 06:05 PM JST

PDB ID : 6JNR
Title : RXRa structure complexed with CU-6PMN and SRC1 peptide.
Authors : Kawasaki, M.; Nakano, S.; Motoyama, T.; Yamada, S.; Watanabe, M.; Takamura, Y.; Fujihara, M.; Tokiwa, H.; Kakuta, H.; Ito, S.
Deposited on : 2019-03-18
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	 77% 9% • 12%
1	B	243	 77% 9% • 12%
2	C	12	 33% 33% 8% 25%
2	D	12	 42% 8% 33% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3709 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Retinoic acid receptor RXR-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	214	1690	1085	291	304	10	0	0	0
1	B	215	1697	1090	292	305	10	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	237	GLY	-	expression tag	UNP P19793
A	238	SER	-	expression tag	UNP P19793
A	239	SER	-	expression tag	UNP P19793
A	240	HIS	-	expression tag	UNP P19793
B	236	GLY	-	expression tag	UNP P19793
B	237	SER	-	expression tag	UNP P19793
B	238	SER	-	expression tag	UNP P19793
B	239	HIS	-	expression tag	UNP P19793

- Molecule 2 is a protein called HIS-LYS-ILE-LEU-HIS-ARG-LEU-LEU-GLN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	9	81	53	18	10	0	0	0
2	D	10	90	58	19	13	0	0	0

- Molecule 3 is 7-oxidanyl-2-oxidanylidene-6-(3,5,5,8,8-pentamethyl-6,7-dihydronaphthalen-2-yl)chromene-3-carboxylic acid (three-letter code: WY5) (formula: C₂₅H₂₆O₅) (labeled as "Ligand of Interest" by depositor).

4 Data and refinement statistics i

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	64.39Å 64.39Å 111.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.39 – 2.30 32.20 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (64.39-2.30) 99.8 (32.20-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.41 (at 2.31Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.179 , 0.238 0.186 , 0.248	Depositor DCC
R_{free} test set	943 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 14.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.489 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3709	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: WY5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.46	0/1725	1.24	13/2334 (0.6%)
1	B	0.46	0/1733	1.27	18/2345 (0.8%)
2	C	0.42	0/82	1.18	1/108 (0.9%)
2	D	0.39	0/91	1.33	1/120 (0.8%)
All	All	0.46	0/3631	1.25	33/4907 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
2	C	0	1
All	All	0	2

There are no bond length outliers.

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	358	ARG	NE-CZ-NH2	-9.71	115.44	120.30
1	A	316	ARG	NE-CZ-NH2	-9.06	115.77	120.30
1	A	379	ASP	CB-CG-OD2	8.84	126.26	118.30
1	A	358	ARG	NE-CZ-NH2	-8.75	115.92	120.30
1	A	359	ASP	CB-CG-OD2	-8.62	110.54	118.30
1	A	359	ASP	CB-CG-OD1	8.06	125.55	118.30
1	B	421	ARG	NE-CZ-NH2	-7.99	116.31	120.30
1	B	379	ASP	CB-CG-OD2	7.69	125.22	118.30
1	B	316	ARG	NE-CZ-NH2	-7.49	116.56	120.30
1	A	379	ASP	CB-CG-OD1	-7.25	111.78	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	371	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	B	379	ASP	CB-CG-OD1	-7.09	111.92	118.30
2	D	6	ARG	NE-CZ-NH2	-7.05	116.77	120.30
1	A	322	ASP	CB-CG-OD2	6.93	124.54	118.30
1	B	322	ASP	CB-CG-OD2	6.11	123.80	118.30
1	B	371	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	B	451	LEU	CB-CG-CD1	5.79	120.84	111.00
1	B	324	ILE	CA-CB-CG2	5.73	122.36	110.90
1	B	456	GLU	OE1-CD-OE2	-5.71	116.45	123.30
1	B	359	ASP	CB-CG-OD2	-5.66	113.21	118.30
1	A	330	LEU	CB-CG-CD2	-5.54	101.57	111.00
1	B	334	ARG	NE-CZ-NH2	5.47	123.03	120.30
1	A	363	ASP	CB-CG-OD2	5.35	123.11	118.30
1	B	336	SER	CB-CA-C	5.31	120.20	110.10
1	A	336	SER	CB-CA-C	5.29	120.15	110.10
1	B	285	ARG	NE-CZ-NH2	-5.28	117.66	120.30
1	B	316	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	309	LEU	CB-CG-CD1	-5.22	102.12	111.00
2	C	6	ARG	NE-CZ-NH2	-5.13	117.73	120.30
1	A	444	ASP	CB-CG-OD2	-5.13	113.69	118.30
1	B	427	SER	N-CA-CB	5.07	118.10	110.50
1	A	405	LYS	CD-CE-NZ	5.01	123.23	111.70
1	B	359	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	262	ASN	Peptide
2	C	8	LEU	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1690	0	1722	8	0
1	B	1697	0	1730	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	81	0	94	1	0
2	D	90	0	100	5	0
3	A	30	0	0	1	0
3	B	30	0	0	0	0
4	A	47	0	0	0	0
4	B	44	0	0	0	0
All	All	3709	0	3646	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1:HIS:N	2:C:1:HIS:ND1	2.42	0.67
1:A:444:ASP:C	1:A:444:ASP:OD1	2.37	0.63
1:A:263:ASP:HB3	1:A:266:THR:HG23	1.81	0.61
2:D:1:HIS:ND1	2:D:1:HIS:N	2.49	0.59
1:A:318:ILE:O	1:A:358:ARG:HD3	2.02	0.58
2:D:9:GLN:O	2:D:10:GLU:HB3	2.05	0.56
1:A:266:THR:HG22	1:A:445:THR:HG23	1.92	0.51
1:B:263:ASP:HB3	1:B:266:THR:HG23	1.91	0.51
1:A:369:CYS:HB2	1:A:400:LEU:HD13	1.93	0.50
1:A:259:VAL:HB	1:A:278:THR:CG2	2.42	0.49
1:B:276:LEU:HD21	1:B:305:TRP:CE3	2.48	0.49
1:A:276:LEU:HD21	1:A:305:TRP:CE3	2.49	0.47
1:B:258:VAL:HB	1:B:278:THR:CG2	2.45	0.46
1:B:251:ILE:O	1:B:254:ALA:HB3	2.16	0.46
1:B:322:ASP:HB3	1:B:334:ARG:HG2	1.97	0.45
2:D:9:GLN:O	2:D:10:GLU:CB	2.65	0.45
2:D:10:GLU:CD	2:D:10:GLU:C	2.77	0.44
1:B:318:ILE:O	1:B:358:ARG:HD3	2.19	0.43
1:A:309:LEU:HB3	3:A:501:WY5:OAA	2.18	0.43
2:D:6:ARG:O	2:D:9:GLN:HB2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/243 (86%)	202 (96%)	8 (4%)	0	100	100
1	B	211/243 (87%)	206 (98%)	5 (2%)	0	100	100
2	C	7/12 (58%)	7 (100%)	0	0	100	100
2	D	8/12 (67%)	7 (88%)	1 (12%)	0	100	100
All	All	436/510 (86%)	422 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/209 (88%)	175 (95%)	9 (5%)	25	35
1	B	185/209 (88%)	177 (96%)	8 (4%)	29	40
2	C	9/11 (82%)	6 (67%)	3 (33%)	0	0
2	D	10/11 (91%)	6 (60%)	4 (40%)	0	0
All	All	388/440 (88%)	364 (94%)	24 (6%)	18	25

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	THR
1	A	312	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	334	ARG
1	A	336	SER
1	A	350	LEU
1	A	410	GLU
1	A	444	ASP
1	A	447	ILE
1	A	450	PHE
1	B	324	ILE
1	B	334	ARG
1	B	336	SER
1	B	350	LEU
1	B	427	SER
1	B	447	ILE
1	B	448	ASP
1	B	450	PHE
2	C	1	HIS
2	C	2	LYS
2	C	9	GLN
2	D	1	HIS
2	D	2	LYS
2	D	9	GLN
2	D	10	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	9	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	WY5	A	501	-	33,33,33	1.62	6 (18%)	52,53,53	2.05	15 (28%)
3	WY5	B	501	-	33,33,33	1.63	7 (21%)	52,53,53	2.20	22 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	WY5	A	501	-	-	3/8/27/27	0/4/4/4
3	WY5	B	501	-	-	3/8/27/27	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	WY5	CAK-CAT	4.73	1.48	1.40
3	B	501	WY5	CAK-CAT	4.34	1.48	1.40
3	B	501	WY5	CAA-CAI	4.01	1.49	1.41
3	B	501	WY5	CAV-CAW	3.83	1.47	1.40
3	A	501	WY5	CAV-CAW	3.79	1.47	1.40
3	A	501	WY5	CAA-CAI	3.53	1.48	1.41
3	B	501	WY5	CAF-CAG	2.94	1.49	1.40
3	A	501	WY5	CAL-CAK	-2.67	1.49	1.53
3	A	501	WY5	CAF-CAG	2.65	1.48	1.40
3	B	501	WY5	OAC-CAD	2.58	1.26	1.21
3	B	501	WY5	CAL-CAK	-2.42	1.50	1.53
3	A	501	WY5	OAC-CAD	2.16	1.26	1.21
3	B	501	WY5	CAA-CAB	-2.12	1.39	1.43

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	WY5	CAE-CAF-CAG	6.06	125.00	117.55
3	A	501	WY5	CAI-CAA-CAB	5.52	122.33	118.04
3	B	501	WY5	CAX-CAW-CAY	5.00	128.73	119.49
3	A	501	WY5	CAE-CAF-CAG	4.65	123.27	117.55
3	A	501	WY5	CAX-CAW-CAY	4.29	127.42	119.49
3	B	501	WY5	CAH-CAG-CAF	-4.25	116.85	120.83
3	B	501	WY5	CAX-CAW-CAV	-4.16	115.27	122.33
3	A	501	WY5	OAA-CAI-CAH	4.09	121.92	115.79
3	B	501	WY5	CAM-CAL-CAK	3.86	116.38	110.16
3	B	501	WY5	CAI-CAA-CAB	3.63	120.87	118.04
3	A	501	WY5	OAA-CAI-CAA	-3.51	116.26	120.55
3	A	501	WY5	CAH-CAG-CAF	-3.46	117.60	120.83
3	A	501	WY5	CAX-CAW-CAV	-3.35	116.64	122.33
3	A	501	WY5	OAA-CAD-OAC	3.18	120.42	116.22
3	A	501	WY5	OAE-CAJ-OAD	-3.07	116.58	123.61
3	B	501	WY5	CAA-CAB-CAC	-3.05	119.16	122.10
3	B	501	WY5	OAA-CAD-OAC	3.01	120.19	116.22
3	B	501	WY5	OAA-CAI-CAH	2.96	120.23	115.79
3	B	501	WY5	CAN-CAO-CAT	2.94	114.90	110.16
3	B	501	WY5	OAE-CAJ-OAD	-2.80	117.21	123.61
3	B	501	WY5	OAA-CAI-CAA	-2.72	117.22	120.55
3	A	501	WY5	CAA-CAB-CAC	-2.71	119.49	122.10
3	B	501	WY5	CAF-CAE-CAA	-2.66	117.06	121.65
3	A	501	WY5	OAA-CAD-CAC	2.56	119.48	117.04
3	B	501	WY5	CAV-CAF-CAG	-2.49	117.48	123.14
3	A	501	WY5	CAU-CAT-CAK	-2.43	116.94	119.76
3	B	501	WY5	CAU-CAT-CAK	-2.34	117.04	119.76
3	B	501	WY5	CAQ-CAO-CAT	-2.33	106.09	110.10
3	A	501	WY5	OAE-CAJ-CAC	2.33	121.61	115.83
3	B	501	WY5	CAR-CAL-CAK	-2.29	106.17	110.10
3	B	501	WY5	CAB-CAC-CAJ	-2.27	117.87	119.79
3	B	501	WY5	CAI-OAA-CAD	2.23	124.41	122.23
3	B	501	WY5	CAY-CAW-CAV	-2.21	115.11	118.78
3	A	501	WY5	CAY-CAW-CAV	-2.19	115.15	118.78
3	B	501	WY5	CAO-CAT-CAK	-2.17	120.36	122.70
3	A	501	WY5	CAE-CAA-CAB	-2.09	119.11	122.77
3	B	501	WY5	OAB-CAG-CAH	2.07	125.00	119.46

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	WY5	CAD-CAC-CAJ-OAD

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	501	WY5	CAD-CAC-CAJ-OAD
3	A	501	WY5	CAB-CAC-CAJ-OAD
3	B	501	WY5	CAB-CAC-CAJ-OAD
3	A	501	WY5	CAB-CAC-CAJ-OAE
3	B	501	WY5	CAB-CAC-CAJ-OAE

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	WY5	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/243 (88%)	-0.75	0 100 100	21, 32, 54, 83	0
1	B	215/243 (88%)	-0.75	0 100 100	21, 32, 51, 99	0
2	C	9/12 (75%)	-0.77	0 100 100	28, 32, 47, 63	0
2	D	10/12 (83%)	-0.73	0 100 100	29, 34, 66, 80	0
All	All	448/510 (87%)	-0.75	0 100 100	21, 32, 54, 99	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	WY5	A	501	30/30	0.96	0.11	23,27,32,34	0
3	WY5	B	501	30/30	0.96	0.11	23,28,33,34	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.