



# wwPDB X-ray Structure Validation Summary Report

Jun 23, 2024 – 12:04 AM EDT

PDB ID : 6KXU  
Title : BON1  
Authors : Wang, Q.C.; Jiang, M.Q.; Isupov, M.N.; Sun, L.F.; Wu, Y.K.  
Deposited on : 2019-09-12  
Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	76% 20% ..
1	C	553	76% 20% ..
1	D	553	74% 22% ..
1	F	553	75% 21% ..
2	B	528	71% 21% .. 5%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	548	
4	G	526	
5	H	553	

## 2 Entry composition i

There are 10 unique types of molecules in this entry. The entry contains 33483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein BONZAI 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	546	4242	2699	711	819	13	0	0	0
1	C	553	4299	2735	719	832	13	0	0	0
1	D	553	4299	2735	719	832	13	0	0	0
1	F	553	4299	2735	719	832	13	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	GLY	-	expression tag	UNP Q941L3
A	-8	THR	-	expression tag	UNP Q941L3
A	-7	SER	-	expression tag	UNP Q941L3
A	-6	SER	-	expression tag	UNP Q941L3
A	-5	MET	-	expression tag	UNP Q941L3
A	-4	ALA	-	expression tag	UNP Q941L3
A	-3	ASP	-	expression tag	UNP Q941L3
A	-2	ILE	-	expression tag	UNP Q941L3
A	-1	GLY	-	expression tag	UNP Q941L3
A	0	SER	-	expression tag	UNP Q941L3
C	-9	GLY	-	expression tag	UNP Q941L3
C	-8	THR	-	expression tag	UNP Q941L3
C	-7	SER	-	expression tag	UNP Q941L3
C	-6	SER	-	expression tag	UNP Q941L3
C	-5	MET	-	expression tag	UNP Q941L3
C	-4	ALA	-	expression tag	UNP Q941L3
C	-3	ASP	-	expression tag	UNP Q941L3
C	-2	ILE	-	expression tag	UNP Q941L3
C	-1	GLY	-	expression tag	UNP Q941L3
C	0	SER	-	expression tag	UNP Q941L3
D	-9	GLY	-	expression tag	UNP Q941L3

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-8	THR	-	expression tag	UNP Q941L3
D	-7	SER	-	expression tag	UNP Q941L3
D	-6	SER	-	expression tag	UNP Q941L3
D	-5	MET	-	expression tag	UNP Q941L3
D	-4	ALA	-	expression tag	UNP Q941L3
D	-3	ASP	-	expression tag	UNP Q941L3
D	-2	ILE	-	expression tag	UNP Q941L3
D	-1	GLY	-	expression tag	UNP Q941L3
D	0	SER	-	expression tag	UNP Q941L3
F	-9	GLY	-	expression tag	UNP Q941L3
F	-8	THR	-	expression tag	UNP Q941L3
F	-7	SER	-	expression tag	UNP Q941L3
F	-6	SER	-	expression tag	UNP Q941L3
F	-5	MET	-	expression tag	UNP Q941L3
F	-4	ALA	-	expression tag	UNP Q941L3
F	-3	ASP	-	expression tag	UNP Q941L3
F	-2	ILE	-	expression tag	UNP Q941L3
F	-1	GLY	-	expression tag	UNP Q941L3
F	0	SER	-	expression tag	UNP Q941L3

- Molecule 2 is a protein called Protein BONZAI 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	504	3932	2507	660	753	12	0	0	0

- Molecule 3 is a protein called Protein BONZAI 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	534	4158	2648	696	802	12	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-4	ALA	-	expression tag	UNP Q941L3
E	-3	ASP	-	expression tag	UNP Q941L3
E	-2	ILE	-	expression tag	UNP Q941L3
E	-1	GLY	-	expression tag	UNP Q941L3
E	0	SER	-	expression tag	UNP Q941L3

- Molecule 4 is a protein called Protein BONZAI 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	G	504	3930	2503	661	754	12	0	0	0


- Molecule 5 is a protein called Protein BONZAI 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	H	543	4227	2691	706	817	13	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-9	GLY	-	expression tag	UNP Q941L3
H	-8	THR	-	expression tag	UNP Q941L3
H	-7	SER	-	expression tag	UNP Q941L3
H	-6	SER	-	expression tag	UNP Q941L3
H	-5	MET	-	expression tag	UNP Q941L3
H	-4	ALA	-	expression tag	UNP Q941L3
H	-3	ASP	-	expression tag	UNP Q941L3
H	-2	ILE	-	expression tag	UNP Q941L3
H	-1	GLY	-	expression tag	UNP Q941L3
H	0	THR	-	expression tag	UNP Q941L3

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	D	1	Total C O 4 2 2	0	0
6	F	1	Total C O 4 2 2	0	0
6	F	1	Total C O 4 2 2	0	0
6	G	1	Total C O 4 2 2	0	0
6	H	1	Total C O 4 2 2	0	0
6	H	1	Total C O 4 2 2	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 6 3 3	0	0

- Molecule 8 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Mn 1 1	0	0
8	B	1	Total Mn 1 1	0	0
8	C	1	Total Mn 1 1	0	0
8	D	2	Total Mn 2 2	0	0
8	E	1	Total Mn 1 1	0	0
8	F	2	Total Mn 2 2	0	0
8	G	1	Total Mn 1 1	0	0
8	H	1	Total Mn 1 1	0	0

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	C	1	Total Cl 1 1	0	0
9	D	1	Total Cl 1 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	4	Total O 4 4	0	0
10	B	6	Total O 6 6	0	0
10	C	7	Total O 7 7	0	0
10	D	3	Total O 3 3	0	0
10	E	1	Total O 1 1	0	0
10	H	2	Total O 2 2	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein BONZAI 1




- Molecule 1: Protein BONZAI 1




- Molecule 1: Protein BONZAI 1







• Molecule 1: Protein BONZAI 1





• Molecule 2: Protein BONZAI 1







• Molecule 3: Protein BONZAI 1




• Molecule 4: Protein BONZAI 1



• Molecule 5: Protein BONZAI 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.41Å 118.70Å 222.32Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	82.31 – 2.83 82.31 – 2.83	Depositor EDS
% Data completeness (in resolution range)	98.2 (82.31-2.83) 98.2 (82.31-2.83)	Depositor EDS
$R_{merge}$	0.19	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 2.82Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.204 , 0.249 0.207 , 0.249	Depositor DCC
$R_{free}$ test set	7469 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	69.9	Xtriage
Anisotropy	0.576	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 38.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.000 for k,h,-l 0.000 for -k,-h,-l 0.246 for h,-k,-l	Xtriage
Reported twinning fraction	0.733 for H, K, L 0.267 for -h,-k,l	Depositor
Outliers	7 of 148582 reflections (0.005%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	33483	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	92.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.70 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.1000e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL, GOL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/4323	0.76	5/5847 (0.1%)
1	C	0.41	0/4382	0.77	3/5929 (0.1%)
1	D	0.44	1/4382 (0.0%)	0.81	7/5929 (0.1%)
1	F	0.40	2/4382 (0.0%)	0.77	4/5929 (0.1%)
2	B	0.41	0/4004	0.80	7/5408 (0.1%)
3	E	0.39	0/4234	0.75	2/5726 (0.0%)
4	G	0.41	0/4001	0.75	2/5408 (0.0%)
5	H	0.39	0/4305	0.76	3/5818 (0.1%)
All	All	0.41	3/34013 (0.0%)	0.77	33/45994 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
1	F	0	1
2	B	0	2
All	All	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	392	GLU	CD-OE2	8.62	1.35	1.25
1	F	447	THR	CB-OG1	5.18	1.53	1.43
1	F	322	SER	CB-OG	5.03	1.48	1.42

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	38	LYS	CB-CA-C	10.71	131.82	110.40
1	F	447	THR	CA-CB-OG1	10.28	130.59	109.00
1	F	447	THR	N-CA-CB	10.24	129.75	110.30
1	A	540	ARG	CG-CD-NE	-8.69	93.54	111.80
1	D	540	ARG	CG-CD-NE	-7.86	95.29	111.80

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	PHE	Peptide
2	B	184	LYS	Peptide
2	B	296	THR	Peptide
1	D	12	LYS	Peptide
1	F	11	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4242	0	4246	76	0
1	C	4299	0	4297	79	0
1	D	4299	0	4298	91	0
1	F	4299	0	4295	96	0
2	B	3932	0	3944	76	0
3	E	4158	0	4172	75	0
4	G	3930	0	3945	82	0
5	H	4227	0	4233	81	0
6	A	12	0	18	0	0
6	B	4	0	6	0	0
6	C	16	0	24	0	0
6	D	4	0	6	1	0
6	F	8	0	12	0	0
6	G	4	0	6	1	0
6	H	8	0	12	0	0
7	A	6	0	8	2	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	1	0	0	0	0
8	D	2	0	0	0	0
8	E	1	0	0	0	0
8	F	2	0	0	0	0
8	G	1	0	0	0	0
8	H	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
10	A	4	0	0	0	0
10	B	6	0	0	0	0
10	C	7	0	0	1	0
10	D	3	0	0	0	0
10	E	1	0	0	0	0
10	H	2	0	0	1	0
All	All	33483	0	33522	633	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 633 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:427:MET:HG3	4:G:184:LYS:N	1.61	1.15
1:F:318:ASP:OD2	1:F:447:THR:OG1	1.65	1.14
1:F:142:PHE:O	1:F:144:PRO:HD3	1.49	1.12
5:H:102:ARG:NH2	10:H:701:HOH:O	1.90	1.02
1:C:195:VAL:CG2	1:C:198:GLY:O	2.16	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	542/553 (98%)	504 (93%)	35 (6%)	3 (1%)	25	46
1	C	551/553 (100%)	503 (91%)	40 (7%)	8 (2%)	10	22
1	D	551/553 (100%)	510 (93%)	30 (5%)	11 (2%)	7	16
1	F	551/553 (100%)	510 (93%)	32 (6%)	9 (2%)	9	21
2	B	494/528 (94%)	463 (94%)	26 (5%)	5 (1%)	15	31
3	E	526/548 (96%)	487 (93%)	31 (6%)	8 (2%)	10	22
4	G	494/526 (94%)	465 (94%)	27 (6%)	2 (0%)	34	56
5	H	535/553 (97%)	494 (92%)	38 (7%)	3 (1%)	25	46
All	All	4244/4367 (97%)	3936 (93%)	259 (6%)	49 (1%)	13	28

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	278	ALA
2	B	280	GLN
1	C	18	PHE
1	C	146	ALA
1	D	144	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	475/481 (99%)	434 (91%)	41 (9%)	10	22
1	C	481/481 (100%)	441 (92%)	40 (8%)	11	24
1	D	481/481 (100%)	442 (92%)	39 (8%)	11	24
1	F	481/481 (100%)	441 (92%)	40 (8%)	11	24
2	B	441/461 (96%)	398 (90%)	43 (10%)	8	16
3	E	467/477 (98%)	428 (92%)	39 (8%)	11	23
4	G	442/460 (96%)	405 (92%)	37 (8%)	11	23
5	H	474/481 (98%)	432 (91%)	42 (9%)	9	20

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	3742/3803 (98%)	3421 (91%)	321 (9%)	10	22

5 of 321 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	397	GLN
5	H	39	SER
1	F	504	SER
4	G	248	SER
5	H	227	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 50 such sidechains are listed below:

Mol	Chain	Res	Type
3	E	281	ASN
1	F	281	ASN
5	H	534	GLN
3	E	312	ASN
1	F	65	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 12 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	GOL	A	604	-	5,5,5	0.19	0	5,5,5	0.50	0
6	EDO	C	603	-	3,3,3	0.09	0	2,2,2	0.18	0
6	EDO	G	601	-	3,3,3	0.10	0	2,2,2	0.66	0
6	EDO	F	602	-	3,3,3	0.96	0	2,2,2	1.48	0
6	EDO	H	602	-	3,3,3	0.16	0	2,2,2	0.33	0
6	EDO	B	601	-	3,3,3	0.19	0	2,2,2	0.25	0
6	EDO	D	601	-	3,3,3	0.05	0	2,2,2	0.23	0
6	EDO	F	601	-	3,3,3	0.07	0	2,2,2	0.19	0
6	EDO	C	601	-	3,3,3	0.44	0	2,2,2	0.50	0
6	EDO	C	604	-	3,3,3	0.14	0	2,2,2	0.06	0
6	EDO	H	601	-	3,3,3	0.13	0	2,2,2	0.17	0
6	EDO	C	602	-	3,3,3	0.34	0	2,2,2	0.26	0
6	EDO	A	601	-	3,3,3	0.41	0	2,2,2	1.39	0
6	EDO	A	603	-	3,3,3	0.29	0	2,2,2	0.39	0
6	EDO	A	602	-	3,3,3	0.08	0	2,2,2	0.00	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. <sup>1,2</sup> means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	604	-	-	2/4/4/4	-
6	EDO	C	603	-	-	0/1/1/1	-
6	EDO	G	601	-	-	1/1/1/1	-
6	EDO	F	602	-	-	1/1/1/1	-
6	EDO	H	602	-	-	1/1/1/1	-
6	EDO	B	601	-	-	1/1/1/1	-
6	EDO	D	601	-	-	0/1/1/1	-
6	EDO	F	601	-	-	1/1/1/1	-
6	EDO	C	601	-	-	0/1/1/1	-
6	EDO	C	604	-	-	1/1/1/1	-
6	EDO	H	601	-	-	1/1/1/1	-
6	EDO	C	602	-	-	1/1/1/1	-
6	EDO	A	601	-	-	1/1/1/1	-
6	EDO	A	603	-	-	1/1/1/1	-
6	EDO	A	602	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	604	GOL	O1-C1-C2-C3
7	A	604	GOL	O1-C1-C2-O2
6	F	602	EDO	O1-C1-C2-O2
6	G	601	EDO	O1-C1-C2-O2
6	A	601	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	604	GOL	2	0
6	G	601	EDO	1	0
6	D	601	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	546/553 (98%)	0.01	5 (0%) 84 83	44, 78, 129, 185	0
1	C	553/553 (100%)	0.03	10 (1%) 68 63	29, 77, 141, 196	0
1	D	553/553 (100%)	0.02	12 (2%) 62 57	25, 80, 147, 220	0
1	F	553/553 (100%)	0.07	19 (3%) 45 37	46, 84, 156, 239	0
2	B	504/528 (95%)	0.18	30 (5%) 21 15	38, 94, 167, 236	0
3	E	534/548 (97%)	0.04	19 (3%) 42 35	50, 86, 162, 206	0
4	G	504/526 (95%)	0.57	56 (11%) 5 3	37, 83, 208, 271	0
5	H	543/553 (98%)	0.10	23 (4%) 36 28	48, 87, 169, 235	0
All	All	4290/4367 (98%)	0.12	174 (4%) 37 29	25, 83, 166, 271	0

The worst 5 of 174 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	G	208	VAL	21.4
4	G	288	LEU	12.4
4	G	263	LEU	12.2
4	G	207	GLU	11.0
4	G	275	PRO	10.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	MN	F	603	1/1	0.68	0.08	127,127,127,127	1
8	MN	F	604	1/1	0.71	0.10	91,91,91,91	1
6	EDO	C	603	4/4	0.75	0.21	88,99,109,110	0
7	GOL	A	604	6/6	0.75	0.15	87,99,99,109	0
8	MN	G	602	1/1	0.77	0.14	106,106,106,106	1
8	MN	E	601	1/1	0.79	0.08	117,117,117,117	1
6	EDO	D	601	4/4	0.82	0.17	89,91,92,103	0
8	MN	C	606	1/1	0.83	0.09	129,129,129,129	1
6	EDO	F	601	4/4	0.84	0.16	94,106,106,113	0
6	EDO	C	601	4/4	0.87	0.19	67,78,83,84	0
8	MN	A	605	1/1	0.88	0.11	133,133,133,133	1
6	EDO	B	601	4/4	0.88	0.22	56,64,68,79	0
9	CL	D	602	1/1	0.88	0.10	73,73,73,73	0
8	MN	H	603	1/1	0.89	0.13	111,111,111,111	1
8	MN	D	603	1/1	0.89	0.25	164,164,164,164	1
6	EDO	H	601	4/4	0.90	0.14	89,94,105,105	0
9	CL	C	605	1/1	0.91	0.19	76,76,76,76	0
6	EDO	A	602	4/4	0.91	0.13	85,92,94,95	0
8	MN	D	604	1/1	0.92	0.10	113,113,113,113	1
6	EDO	C	602	4/4	0.92	0.17	58,60,67,72	0
6	EDO	C	604	4/4	0.93	0.18	80,82,86,90	0
6	EDO	A	603	4/4	0.93	0.12	69,75,84,88	0
6	EDO	H	602	4/4	0.95	0.22	68,84,84,89	0
6	EDO	G	601	4/4	0.96	0.21	60,68,71,71	0
8	MN	B	602	1/1	0.96	0.04	141,141,141,141	1
6	EDO	A	601	4/4	0.97	0.18	33,37,38,42	0
6	EDO	F	602	4/4	0.98	0.21	21,29,30,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around MN F 603:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN F 604:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN G 602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN E 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN C 606:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around MN A 605:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN H 603:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around MN D 603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









## 6.5 Other polymers [i](#)

There are no such residues in this entry.