



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 19, 2023 – 05:28 PM JST

PDB ID : 6LTW
Title : Crystal structure of Apo form of I122A/I330A variant of S-adenosylmethionine synthetase from *Cryptosporidium hominis*
Authors : Singh, R.K.; Michailidou, F.; Rentmeister, A.; Kuemmel, D.
Deposited on : 2020-01-23
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	 4% 80% 12% 8%
1	B	414	 4% 82% 9% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	A	503	-	X	-	-
3	PO4	B	504	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6547 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S-adenosylmethionine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	380	2992	1889	514	571	18	0	5	0
1	B	379	2951	1866	500	566	19	0	3	0


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP A0A0S4TKQ5
A	-6	ALA	-	expression tag	UNP A0A0S4TKQ5
A	-5	HIS	-	expression tag	UNP A0A0S4TKQ5
A	-4	HIS	-	expression tag	UNP A0A0S4TKQ5
A	-3	HIS	-	expression tag	UNP A0A0S4TKQ5
A	-2	HIS	-	expression tag	UNP A0A0S4TKQ5
A	-1	HIS	-	expression tag	UNP A0A0S4TKQ5
A	0	HIS	-	expression tag	UNP A0A0S4TKQ5
A	122	ALA	ILE	variant	UNP A0A0S4TKQ5
A	330	ALA	ILE	variant	UNP A0A0S4TKQ5
B	-7	MET	-	initiating methionine	UNP A0A0S4TKQ5
B	-6	ALA	-	expression tag	UNP A0A0S4TKQ5
B	-5	HIS	-	expression tag	UNP A0A0S4TKQ5
B	-4	HIS	-	expression tag	UNP A0A0S4TKQ5
B	-3	HIS	-	expression tag	UNP A0A0S4TKQ5
B	-2	HIS	-	expression tag	UNP A0A0S4TKQ5
B	-1	HIS	-	expression tag	UNP A0A0S4TKQ5
B	0	HIS	-	expression tag	UNP A0A0S4TKQ5
B	122	ALA	ILE	variant	UNP A0A0S4TKQ5
B	330	ALA	ILE	variant	UNP A0A0S4TKQ5

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Mg 2 2	0	0
2	B	2	Total Mg 2 2	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	305	Total O 305 305	0	0
4	B	275	Total O 275 275	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: S-adenosylmethionine synthase



- Molecule 1: S-adenosylmethionine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.88Å 100.36Å 66.84Å 90.00° 95.86° 90.00°	Depositor
Resolution (Å)	48.08 – 1.65 48.07 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.08-1.65) 99.0 (48.07-1.65)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.8.0218	Depositor
R, R_{free}	0.156 , 0.191 0.169 , 0.201	Depositor DCC
R_{free} test set	4768 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtrriage
Anisotropy	0.524	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6547	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.28	7/3046 (0.2%)	1.21	18/4104 (0.4%)
1	B	1.25	8/3004 (0.3%)	1.21	22/4050 (0.5%)
All	All	1.27	15/6050 (0.2%)	1.21	40/8154 (0.5%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	32	SER	CB-OG	-11.38	1.27	1.42
1	A	32	SER	CB-OG	-8.14	1.31	1.42
1	A	133	GLU	CD-OE1	7.46	1.33	1.25
1	A	195	GLU	CD-OE1	7.28	1.33	1.25
1	B	339	TYR	CE1-CZ	-7.14	1.29	1.38
1	B	168	GLU	CD-OE1	-6.09	1.19	1.25
1	B	167	ARG	CZ-NH1	5.97	1.40	1.33
1	B	164	SER	CB-OG	-5.67	1.34	1.42
1	A	171	TYR	CG-CD1	-5.59	1.31	1.39
1	A	97	TYR	CE1-CZ	5.46	1.45	1.38
1	A	150	GLU	CB-CG	-5.34	1.42	1.52
1	A	399	TRP	CG-CD1	-5.29	1.29	1.36
1	B	28	GLU	CB-CG	-5.22	1.42	1.52
1	B	225	GLU	CB-CG	-5.10	1.42	1.52
1	B	272	ARG	N-CA	5.10	1.56	1.46

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	167	ARG	NE-CZ-NH2	-10.80	114.90	120.30
1	A	139	ASP	CB-CG-OD2	-8.36	110.78	118.30
1	A	139	ASP	CB-CG-OD1	8.27	125.74	118.30
1	B	304	TYR	CB-CG-CD1	7.96	125.78	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	167	ARG	NE-CZ-NH1	7.95	124.27	120.30
1	B	180	ARG	NE-CZ-NH1	7.80	124.20	120.30
1	A	170	ASP	CB-CG-OD2	-7.73	111.34	118.30
1	A	187	ASP	CB-CG-OD1	6.95	124.55	118.30
1	B	139	ASP	CB-CG-OD2	-6.81	112.17	118.30
1	B	187	ASP	CB-CG-OD1	6.80	124.42	118.30
1	B	48	ASP	CB-CG-OD2	-6.77	112.20	118.30
1	B	142[A]	MET	CG-SD-CE	-6.71	89.46	100.20
1	B	142[B]	MET	CG-SD-CE	-6.71	89.46	100.20
1	A	207	ARG	NE-CZ-NH1	6.59	123.60	120.30
1	B	370	LEU	CB-CG-CD1	-6.54	99.88	111.00
1	B	336	LEU	CB-CG-CD2	6.53	122.09	111.00
1	B	180	ARG	NE-CZ-NH2	-6.42	117.09	120.30
1	B	218	ASP	CB-CG-OD1	6.36	124.02	118.30
1	B	36	ASP	CB-CG-OD1	6.06	123.75	118.30
1	A	307	ARG	NE-CZ-NH1	-6.04	117.28	120.30
1	A	400	GLU	OE1-CD-OE2	-6.03	116.06	123.30
1	B	370	LEU	CA-CB-CG	5.93	128.93	115.30
1	A	187	ASP	CB-CG-OD2	-5.84	113.04	118.30
1	A	218	ASP	CB-CG-OD1	5.84	123.56	118.30
1	A	242	ASP	CB-CG-OD1	5.84	123.56	118.30
1	A	352	ASP	CB-CG-OD2	5.82	123.54	118.30
1	A	406	GLN	CA-C-O	-5.77	107.99	120.10
1	B	258	PHE	CB-CG-CD1	5.67	124.77	120.80
1	A	242	ASP	CB-CG-OD2	-5.54	113.31	118.30
1	A	146	TYR	CB-CG-CD1	-5.49	117.71	121.00
1	A	48	ASP	CB-CG-OD2	-5.48	113.37	118.30
1	B	139	ASP	CB-CG-OD1	5.47	123.22	118.30
1	B	299	ASP	CB-CG-OD1	-5.44	113.41	118.30
1	A	105	ASP	CB-CG-OD1	5.43	123.19	118.30
1	B	370	LEU	N-CA-CB	-5.36	99.68	110.40
1	A	54	ASP	CB-CG-OD1	5.28	123.05	118.30
1	B	71	MET	CG-SD-CE	5.24	108.58	100.20
1	A	370	LEU	CA-CB-CG	5.17	127.19	115.30
1	B	258	PHE	CB-CG-CD2	-5.06	117.26	120.80
1	B	364	ASP	CB-CG-OD1	5.03	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2992	0	3002	16	0
1	B	2951	0	2955	11	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	10	0	0	0	0
3	B	10	0	0	0	0
4	A	305	0	0	2	0
4	B	275	0	0	2	0
All	All	6547	0	5957	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:ILE:CG2	1:B:181:VAL:HG11	2.12	0.79
1:B:100:GLU:HB2	4:B:806:HOH:O	1.89	0.70
1:B:340:ILE:HD11	1:B:359:VAL:HG21	1.85	0.58
1:A:94:GLU:OE1	1:A:382:LYS:NZ	2.37	0.58
1:B:142[A]:MET:HB3	1:B:327[A]:SER:OG	2.06	0.56
1:A:133:GLU:HG3	4:A:639:HOH:O	2.06	0.55
1:A:117:GLN:O	1:A:118:GLN:CB	2.57	0.52
1:A:227[B]:ARG:NH2	1:A:250:ILE:O	2.43	0.51
1:B:66:LYS:HG3	1:B:67:THR:H	1.76	0.51
1:B:72:VAL:HG22	1:B:113:ILE:HD13	1.93	0.50
1:A:117:GLN:O	1:A:118:GLN:HB3	2.12	0.49
1:A:94:GLU:HG3	4:A:722:HOH:O	2.11	0.49
1:B:172:ILE:HG21	1:B:181:VAL:HG11	1.93	0.49
1:B:167:ARG:HD3	4:B:841:HOH:O	2.14	0.47
1:A:172:ILE:HG23	1:A:181:VAL:HG11	1.96	0.47
1:B:76:ILE:HD12	1:B:115:LEU:HD13	1.96	0.46
1:A:83:ASN:HD22	1:A:86:ARG:HH21	1.64	0.45
1:A:117:GLN:HB3	1:A:118:GLN:HE21	1.82	0.44
1:B:172:ILE:HG22	1:B:181:VAL:HG11	1.94	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:66:LYS:O	1:B:67:THR:C	2.56	0.44
1:A:201:GLY:O	1:A:321:ARG:HD3	2.19	0.43
1:A:99:SER:HG	1:A:101:GLU:CD	2.23	0.42
1:A:227[A]:ARG:HG2	1:A:252:ILE:HB	2.01	0.41
1:A:315:PHE:HE2	1:A:404:ILE:HD11	1.86	0.41
1:A:268:GLY:C	1:A:269:LEU:HD12	2.41	0.41
1:A:148:THR:O	1:A:154:LEU:HA	2.21	0.40
1:A:75:GLU:OE2	1:A:118:GLN:HA	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	381/414 (92%)	371 (97%)	10 (3%)	0	100	100
1	B	378/414 (91%)	367 (97%)	11 (3%)	0	100	100
All	All	759/828 (92%)	738 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/354 (93%)	324 (98%)	6 (2%)	59	36

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	325/354 (92%)	316 (97%)	9 (3%)	43	18
All	All	655/708 (92%)	640 (98%)	15 (2%)	52	25

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	GLU
1	A	22	GLN
1	A	142	MET
1	A	143	MET
1	A	199	LYS
1	A	403	ILE
1	B	22	GLN
1	B	101	GLU
1	B	142[A]	MET
1	B	142[B]	MET
1	B	143	MET
1	B	181	VAL
1	B	242	ASP
1	B	257	ARG
1	B	406	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	118	GLN
1	A	159	HIS
1	B	373	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	PO4	B	504	2	4,4,4	1.46	1 (25%)	6,6,6	3.30	4 (66%)
3	PO4	A	503	2	4,4,4	1.19	1 (25%)	6,6,6	3.17	4 (66%)
3	PO4	B	503	2	4,4,4	1.55	1 (25%)	6,6,6	1.49	1 (16%)
3	PO4	A	504	2	4,4,4	1.20	1 (25%)	6,6,6	1.35	1 (16%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	504	PO4	P-O2	-2.34	1.47	1.54
3	B	503	PO4	P-O1	2.28	1.56	1.50
3	B	504	PO4	P-O2	-2.23	1.47	1.54
3	A	503	PO4	P-O2	-2.16	1.48	1.54

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	B	504	PO4	O4-P-O1	-5.44	90.98	110.89
3	A	503	PO4	O4-P-O1	-5.24	91.73	110.89
3	B	504	PO4	O2-P-O1	4.77	128.36	110.89
3	A	503	PO4	O3-P-O2	3.33	118.65	107.97
3	A	503	PO4	O4-P-O2	3.26	118.42	107.97
3	A	503	PO4	O3-P-O1	-2.55	101.55	110.89
3	B	504	PO4	O3-P-O2	2.45	115.82	107.97
3	A	504	PO4	O2-P-O1	-2.40	102.11	110.89
3	B	503	PO4	O4-P-O2	-2.34	100.45	107.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	504	PO4	O4-P-O2	-2.30	100.60	107.97


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	380/414 (91%)	0.13	17 (4%) 33 32	14, 26, 47, 81	0
1	B	379/414 (91%)	0.11	17 (4%) 33 32	16, 28, 50, 68	0
All	All	759/828 (91%)	0.12	34 (4%) 33 32	14, 27, 48, 81	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	178	SER	6.8
1	A	179	SER	6.4
1	B	178	SER	5.8
1	A	125	CYS	5.0
1	B	125	CYS	5.0
1	B	179	SER	4.9
1	A	118	GLN	3.9
1	A	128	VAL	3.8
1	B	128	VAL	3.8
1	A	176	GLY	3.4
1	A	180	ARG	3.3
1	A	181	VAL	3.3
1	B	181	VAL	3.2
1	B	176	GLY	3.2
1	A	126	VAL	3.1
1	B	86	ARG	3.0
1	B	274	ILE	3.0
1	A	274	ILE	2.9
1	B	406	GLN	2.8
1	A	127	HIS	2.7
1	B	127	HIS	2.7
1	B	81	ASN	2.7
1	A	117	GLN	2.6
1	B	275	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	86	ARG	2.3
1	A	21	GLU	2.3
1	A	81	ASN	2.2
1	B	99	SER	2.2
1	B	180	ARG	2.2
1	A	275	ILE	2.1
1	A	200	HIS	2.1
1	B	106	TYR	2.1
1	B	117	GLN	2.1
1	B	202	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	A	502	1/1	0.88	0.25	39,39,39,39	0
2	MG	B	502	1/1	0.97	0.31	47,47,47,47	0
3	PO4	A	504	5/5	0.97	0.09	21,23,24,27	0
3	PO4	B	503	5/5	0.98	0.07	18,21,24,25	0
3	PO4	B	504	5/5	0.98	0.08	23,25,28,37	0
2	MG	B	501	1/1	0.99	0.04	18,18,18,18	0
2	MG	A	501	1/1	0.99	0.05	20,20,20,20	0
3	PO4	A	503	5/5	0.99	0.09	24,25,30,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 A 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around PO4 B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around PO4 B 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG B 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.