

Full wwPDB X-ray Structure Validation Report (i)

Jun 11, 2024 – 06:17 PM EDT

PDB ID	:	6MXX
Title	:	Structure of 53BP1 tandem Tudor domains in complex with small molecule
		UNC2991
Authors	:	Cui, G.; Botuyan, M.V.; Mer, G.
Deposited on	:	2018-10-31
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
R _{free}	130704	5042 (2.30-2.30)		
Clashscore	141614	5643 (2.30-2.30)		
Ramachandran outliers	138981	5575(2.30-2.30)		
Sidechain outliers	138945	5575(2.30-2.30)		
RSRZ outliers	127900	4938 (2.30-2.30)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
		100	3%		
1	А	123	91%	7% •	
			7%		_
1	В	123	88%	9% •	
			7%		
1	С	123	93%	5%•	1
			8%		
1	D	123	88%	10% •	1
			14%		
1	E	123	89%	9% •	



Mol	Chain	Length	Quality of chain	
1	F	193		
1	T	120	91% //	%•
1	G	123	91% 5%	
1	ц	192	11%	
1	11	120	89% 7%	••
1	Ι	123	89% 7%	•
1	т	102	9%	
	յ	123	93%	•••

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	В	1703	-	-	-	Х
2	PO4	С	1702	-	-	-	Х
2	PO4	D	1703	-	-	-	Х
2	PO4	Е	1702	-	-	-	Х
3	K6P	В	1701[A]	-	-	-	Х
3	K6P	В	1701[B]	-	-	-	Х
3	K6P	J	1701[A]	-	-	-	Х
3	K6P	J	1701[B]	-	-	-	Х



6MXX

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 20865 atoms, of which 10103 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues			Atom	ıs			ZeroOcc	AltConf	Trace
1	Δ	120	Total	С	Η	Ν	Ο	\mathbf{S}	6	13	0
1	11	120	2051	659	1021	174	194	3	0	10	0
1	В	110	Total	С	Η	Ν	Ο	\mathbf{S}	11	7	0
	D	115	1968	637	978	165	185	3	11		0
1	С	191	Total	С	Η	Ν	Ο	\mathbf{S}	10	14	0
1	U	121	2066	667	1026	175	195	3	10	14	0
1	Л	120	Total	С	Η	Ν	Ο	\mathbf{S}	11	13	0
1	D	120	2045	659	1019	171	193	3	11	10	U
1	F	120	Total	С	Η	Ν	Ο	\mathbf{S}	19	6	0
L		120	1944	627	965	163	186	3	12	0	0
1	F	120	Total	С	Η	Ν	Ο	\mathbf{S}	10	14	0
1	I.	120	2066	662	1031	176	193	4	19	14	0
1	С	110	Total	С	Η	Ν	0	\mathbf{S}	91	12	0
	G	115	1974	644	966	165	195	4	21	10	0
1	ц	110	Total	С	Η	Ν	0	\mathbf{S}	8	10	0
	11	115	1993	643	990	167	188	5	0	10	0
1	т	118	Total	С	Η	Ν	0	\mathbf{S}	16	2	0
	1	110	1831	596	901	157	174	3	10		
1	т	120	Total	С	Н	Ν	0	S	93	10	0
	J	120	2005	648	992	170	192	3	20	10	

• Molecule 1 is a protein called TP53-binding protein 1.

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1481	GLY	-	expression tag	UNP Q12888
А	1482	HIS	-	expression tag	UNP Q12888
А	1483	MET	-	expression tag	UNP Q12888
В	1481	GLY	-	expression tag	UNP Q12888
В	1482	HIS	-	expression tag	UNP Q12888
В	1483	MET	-	expression tag	UNP Q12888
С	1481	GLY	-	expression tag	UNP Q12888
С	1482	HIS	-	expression tag	UNP Q12888
C	1483	MET	-	expression tag	UNP Q12888



6MXX

Chain	Residue	Modelled	Actual	Comment	Reference
D	1401	CLV	metual		
D	1481	GLY	-	expression tag	UNP Q12888
D	1482	HIS	-	expression tag	UNP Q12888
D	1483	MET	-	expression tag	UNP Q12888
Ε	1481	GLY	-	expression tag	UNP Q12888
Ε	1482	HIS	-	expression tag	UNP Q12888
E	1483	MET	-	expression tag	UNP Q12888
F	1481	GLY	-	expression tag	UNP Q12888
F	1482	HIS	-	expression tag	UNP Q12888
F	1483	MET	-	expression tag	UNP Q12888
G	1481	GLY	-	expression tag	UNP Q12888
G	1482	HIS	-	expression tag	UNP Q12888
G	1483	MET	-	expression tag	UNP Q12888
Н	1481	GLY	-	expression tag	UNP Q12888
Н	1482	HIS	-	expression tag	UNP Q12888
Н	1483	MET	-	expression tag	UNP Q12888
Ι	1481	GLY	-	expression tag	UNP Q12888
Ι	1482	HIS	-	expression tag	UNP Q12888
Ι	1483	MET	-	expression tag	UNP Q12888
J	1481	GLY	-	expression tag	UNP Q12888
J	1482	HIS	-	expression tag	UNP Q12888
J	1483	MET	-	expression tag	UNP Q12888

• Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	А	1	Total 5	0 4	Р 1	0	0

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	С	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	D	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Е	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Е	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	F	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	F	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	G	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	G	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Н	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Н	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Ι	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	Ι	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	J	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0
2	J	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{P} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is N-[3-(tert-butylamino)propyl]-3-iodobenzamide (three-letter code: K6P) (formula: $C_{14}H_{21}IN_2O$).





Mol	Chain	Residues	Atoms	ZeroOcc AltConf
3	В	1	Total C H I N O	0 1
0	D	T	78 28 42 2 4 2	
3	С	1	Total C H I N O	0 1
0	U	T	78 28 42 2 4 2	
3	F	1	Total C H I N O	0 1
0	Ľ	L L	78 28 42 2 4 2	
3	C	1	Total C H I N O	0 1
0	G	T	78 28 42 2 4 2	
3	T	1	Total C H I N O	0 1
5	J	L	78 28 42 2 4 2	

• Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{H} & \text{O} \\ 4 & 1 & 1 & 2 \end{array}$	0	0
4	Н	1	Total C H O 4 1 1 2	0	0
4	Н	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{H} & \text{O} \\ 4 & 1 & 1 & 2 \end{array}$	0	0
4	Н	1	$\begin{array}{ccccc} \text{Total} & \text{C} & \text{H} & \text{O} \\ 4 & 1 & 1 & 2 \end{array}$	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	57	Total O 57 57	0	0
5	В	46	Total O 46 46	0	0
5	С	49	Total O 49 49	0	0
5	D	49	Total O 49 49	0	0
5	Е	36	Total O 36 36	0	0
5	F	44	Total O 44 44	0	0
5	G	27	$\begin{array}{cc} \text{Total} & \text{O} \\ 27 & 27 \end{array}$	0	0
5	Н	43	Total O 43 43	0	0
5	Ι	29	TotalO2929	0	0
5	J	31	Total O 31 31	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: TP53-binding protein 1



• Molecule 1: TP53-binding protein	1	
Chain F:	91%	7%•
GLY HIS MAT NAFT NI499 V1492 V1492 V1498 V1498 V1498 V1498 V1499 V1501 V1512 V1512 V1544 V1544	M1584	
• Molecule 1: TP53-binding protein	1	
Chain G:	91%	5% • •
GLY HIS MET MET MET M1434 F1486 F1486 F1486 A1493 A1493 A1493 A1495 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1496 S1495 S1496 S1495 S1496 S1495 S145 S145 S145 S145 S15 S145 S15 S145 S15 S145 S15 S15 S15 S15 S15 S15 S15 S15 S15 S1	N1532 11532 11532 11532 11567 11597 11597 11502 GLY	
• Molecule 1: TP53-binding protein	1	
Chain H:	89%	7% ••
GLY HIS MET MET MET N1489 F1486 F1486 F1499 K1499 S1496 S1496 S1496 F1501 F1501 F1501 F1501 F1501 F1501 F1513 F1513 F1513	L1528 R1522 K1563 E1564 E1575 L1562 GLY	
• Molecule 1: TP53-binding protein	1	
Chain I:	89%	7% •
GLY HIS MET ASN ASN ASN ASN 1485 71486 71486 71486 71495 71495 71495 71495 71495 71495 71500 71500 71510 71511 71512 71511 71512 71512 71512	Lises G1553 D1531 D1532 D1533 L1582 C1576 C1575 C1576 L1588 L1588 L1588 L1588 L1590 L1590 L1500 L1500 L1500 G1576 G1500 G1500	
• Molecule 1: TP53-binding protein	1	
Chain J:	93%	
GLY H1S M1483 M1483 M1485 F1486 F1486 F1486 F1496 81496 81496 81496 81496 81496 81496 81496 81496 81496 81496 81496 11499 11502 L1517 L1528 L1528 L1528 L1602		





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	68.94Å 159.43Å 181.19Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution (Å)	48.14 - 2.30	Depositor
Resolution (A)	48.14 - 2.30	EDS
% Data completeness	97.5 (48.14-2.30)	Depositor
(in resolution range)	97.5(48.14-2.30)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.45 (at 2.29 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
P. P.	0.206 , 0.232	Depositor
Λ, Λ_{free}	0.206 , 0.231	DCC
R_{free} test set	2000 reflections $(2.29%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	44.9	Xtriage
Anisotropy	0.019	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.41 , 49.6	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	20865	wwPDB-VP
Average B, all atoms $(Å^2)$	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, K6P, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.26	0/1075	0.45	0/1444
1	В	0.25	0/1022	0.45	0/1375
1	С	0.28	0/1093	0.46	0/1472
1	D	0.26	0/1082	0.45	0/1454
1	Е	0.25	0/1025	0.44	0/1378
1	F	0.25	0/1088	0.45	0/1460
1	G	0.25	0/1064	0.44	0/1436
1	Н	0.25	0/1046	0.45	0/1409
1	Ι	0.25	0/954	0.44	0/1285
1	J	0.25	0/1055	0.43	0/1420
All	All	0.26	0/10504	0.45	0/14133

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1030	1021	999	6	0
1	В	990	978	971	6	0
1	C	1040	1026	1004	4	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1026	1019	999	9	0
1	Е	979	965	943	6	0
1	F	1035	1031	1003	5	0
1	G	1008	966	946	5	0
1	Н	1003	990	978	6	0
1	Ι	930	901	898	5	0
1	J	1013	992	970	3	0
2	А	5	0	0	0	0
2	В	10	0	0	0	0
2	С	10	0	0	0	0
2	D	20	0	0	0	0
2	Ε	10	0	0	0	0
2	F	10	0	0	0	0
2	G	10	0	0	0	0
2	Н	10	0	0	0	0
2	Ι	10	0	0	0	0
2	J	10	0	0	0	0
3	В	36	42	0	0	0
3	С	36	42	0	1	0
3	Е	36	42	0	0	0
3	G	36	42	0	0	0
3	J	36	42	0	0	0
4	В	3	1	1	0	0
4	Н	9	3	3	0	0
5	А	57	0	0	1	0
5	В	46	0	0	0	0
5	С	49	0	0	0	0
5	D	49	0	0	0	0
5	Ε	36	0	0	0	0
5	F	44	0	0	0	0
5	G	27	0	0	0	0
5	Н	43	0	0	0	0
5	Ι	29	0	0	0	0
5	J	31	0	0	0	0
All	All	10762	10103	9715	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic	Clash
	1100111-2	distance (Å)	overlap (Å)
1:G:1567:GLU:OE1	1:G:1582:LYS:NZ	2.27	0.68
1:H:1562:ARG:NH1	1:H:1564:GLU:OE2	2.27	0.67
1:I:1530:LYS:O	1:I:1597:ARG:NH1	2.30	0.65
1:A:1534[B]:LEU:HD11	1:A:1602:LEU:HD21	1.79	0.64
1:J:1492:VAL:HG12	1:J:1501:PHE:HB3	1.81	0.63
1:E:1492:VAL:HG12	1:E:1501:PHE:HB3	1.81	0.63
1:F:1492:VAL:HG12	1:F:1501:PHE:HB3	1.82	0.61
1:A:1492:VAL:HG12	1:A:1501:PHE:HB3	1.82	0.61
1:J:1498[A]:ASN:OD1	1:J:1499[A]:GLY:N	2.33	0.61
1:G:1492:VAL:HG12	1:G:1501:PHE:HB3	1.83	0.61
1:F:1530:LYS:O	1:F:1597:ARG:NH1	2.37	0.58
1:B:1490:ARG:NH2	1:B:1520:ASP:OD1	2.37	0.58
1:I:1492:VAL:HG22	1:I:1501:PHE:HB3	1.85	0.58
1:H:1492:VAL:HG12	1:H:1501:PHE:HB3	1.87	0.57
1:D:1505:LYS:O	1:D:1517[B]:LEU:HD12	2.06	0.56
1:B:1517[B]:LEU:HD23	1:B:1519:PHE:CZ	2.44	0.53
1:C:1517[B]:LEU:HD23	1:C:1519:PHE:CZ	2.44	0.52
1:G:1495[B]:TRP:HA	1:G:1495[B]:TRP:CE3	2.47	0.49
1:J:1498[A]:ASN:ND2	1:J:1502:TYR:OH	2.44	0.49
1:D:1517[A]:LEU:HD12	1:D:1527:VAL:HG21	1.95	0.48
1:A:1534[B]:LEU:CD1	1:A:1602:LEU:HD21	2.44	0.48
1:E:1497:SER:OG	1:E:1498:ASN:N	2.46	0.47
1:D:1517[B]:LEU:HD23	1:D:1519:PHE:CZ	2.50	0.46
1:A:1562:ARG:NH1	5:A:1803:HOH:O	2.48	0.46
1:B:1543:GLU:OE1	1:B:1595:ARG:NH1	2.47	0.46
1:F:1544:VAL:HG12	1:F:1588:LEU:CD2	2.46	0.46
1:B:1562:ARG:NH2	1:B:1564:GLU:OE2	2.50	0.45
1:G:1495[B]:TRP:HA	1:G:1495[B]:TRP:HE3	1.81	0.44
1:H:1505:LYS:O	1:H:1517[B]:LEU:HD12	2.17	0.44
1:D:1492[B]:VAL:HG22	1:D:1501:PHE:HB3	1.99	0.44
1:F:1544:VAL:HG12	1:F:1588:LEU:HD21	1.99	0.44
1:I:1498:ASN:ND2	1:I:1502:TYR:OH	2.51	0.44
1:D:1492[A]:VAL:HG12	1:D:1501:PHE:HB3	1.99	0.43
1:D:1493[B]:ALA:HB3	1:D:1517[B]:LEU:CD2	2.49	0.43
1:H:1494[B]:LYS:O	1:H:1495[B]:TRP:O	2.37	0.43
1:B:1494[B]:LYS:NZ	1:B:1499:GLY:O	2.46	0.43
1:C:1521:ASP:HB3	3:C:1701[A]:K6P:C4	2.49	0.42
1:E:1493:ALA:HB3	1:E:1517[A]:LEU:CD2	2.49	0.42
1:B:1494[B]:LYS:HB3	1:B:1531:ASP:HB3	2.01	0.42
1:E:1491:VAL:HG12	1:E:1534:LEU:HD12	2.00	0.42
1:D:1491:VAL:HG12	1:D:1534:LEU:HD12	2.02	0.42
1:H:1517[B]:LEU:HD23	1:H:1519:PHE:CZ	2.54	0.42



Atom-1	Atom-2	Interatomic	Clash
		distance (A)	overlap (A)
1:D:1594:ASN:OD1	1:D:1597:ARG:NH2	2.47	0.41
1:I:1492:VAL:O	1:I:1492:VAL:HG13	2.19	0.41
1:E:1508[A]:ARG:HD3	1:E:1509:ASP:N	2.35	0.41
1:I:1544:VAL:HG12	1:I:1588:LEU:HD21	2.01	0.41
1:F:1517[A]:LEU:HD12	1:F:1527:VAL:HG21	2.02	0.41
1:A:1493[B]:ALA:HB3	1:A:1517[B]:LEU:CD2	2.50	0.41
1:G:1530:LYS:O	1:G:1597:ARG:NH1	2.54	0.40
1:C:1493[B]:ALA:HB3	1:C:1517[B]:LEU:CD2	2.52	0.40
1:C:1527[B]:VAL:HG23	1:C:1532:ILE:HD11	2.04	0.40
1:E:1517[A]:LEU:HD23	1:E:1519:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	131/123~(106%)	128~(98%)	3~(2%)	0	100	100
1	В	124/123~(101%)	120~(97%)	4 (3%)	0	100	100
1	С	133/123~(108%)	131~(98%)	2(2%)	0	100	100
1	D	131/123~(106%)	120~(92%)	9~(7%)	2(2%)	10	10
1	Е	124/123~(101%)	119~(96%)	5 (4%)	0	100	100
1	F	132/123~(107%)	127~(96%)	5 (4%)	0	100	100
1	G	130/123~(106%)	122 (94%)	6 (5%)	2(2%)	10	10
1	Н	127/123~(103%)	120 (94%)	5 (4%)	2(2%)	9	9
1	Ι	118/123~(96%)	114 (97%)	4 (3%)	0	100	100
1	J	128/123~(104%)	122 (95%)	6 (5%)	0	100	100
All	All	$127\overline{8}/1230~(104\%)$	1223 (96%)	49 (4%)	6 (0%)	41	35



All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Н	1495[A]	TRP
1	Н	1495[B]	TRP
1	G	1495[A]	TRP
1	G	1495[B]	TRP
1	D	1499[A]	GLY
1	D	1499[B]	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Rotameric Outlie		Outliers	Percentiles
1	А	110/103~(107%)	110 (100%)	0	100 100
1	В	104/103~(101%)	104 (100%)	0	100 100
1	\mathbf{C}	112/103~(109%)	112 (100%)	0	100 100
1	D	109/103~(106%)	109 (100%)	0	100 100
1	Ε	104/103~(101%)	104 (100%)	0	100 100
1	F	110/103~(107%)	110 (100%)	0	100 100
1	G	106/103~(103%)	106 (100%)	0	100 100
1	Η	108/103~(105%)	108 (100%)	0	100 100
1	Ι	94/103~(91%)	94 (100%)	0	100 100
1	J	107/103~(104%)	107 (100%)	0	100 100
All	All	1064/1030~(103%)	1064 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. There are no such side chains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

35 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	Chain	Dog	Link	Bo	ond leng	ths	Bond angles		les
	туре	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	PO4	D	1701	-	4,4,4	0.92	0	6,6,6	0.43	0
4	FMT	Н	1701	-	2,2,2	0.61	0	1,1,1	0.61	0
3	K6P	G	1701[A]	-	18,18,18	0.17	0	23,24,24	0.45	0
3	K6P	В	1701[A]	-	18,18,18	0.19	0	23,24,24	0.45	0
2	PO4	В	1703	-	4,4,4	0.94	0	6,6,6	0.42	0
2	PO4	F	1701	-	4,4,4	0.91	0	6,6,6	0.43	0
2	PO4	G	1702	-	4,4,4	0.93	0	6,6,6	0.41	0
2	PO4	F	1702	-	4,4,4	0.92	0	6,6,6	0.42	0
2	PO4	Н	1705	-	4,4,4	0.95	0	6,6,6	0.44	0
3	K6P	Е	1701[A]	-	18,18,18	0.18	0	23,24,24	0.46	0
2	PO4	Е	1703	-	4,4,4	0.91	0	6,6,6	0.43	0
4	FMT	Н	1703	-	$2,\!2,\!2$	0.63	0	1,1,1	0.61	0
3	K6P	В	1701[B]	-	18,18,18	0.17	0	23,24,24	0.39	0
2	PO4	С	1703	-	4,4,4	0.92	0	6,6,6	0.43	0
2	PO4	Е	1702	-	4,4,4	0.93	0	6,6,6	0.43	0
3	K6P	G	1701[B]	-	18,18,18	0.18	0	23,24,24	0.42	0
2	PO4	Ι	1701	-	4,4,4	0.92	0	6,6,6	0.43	0
3	K6P	С	1701[A]	-	18,18,18	0.20	0	23,24,24	0.58	0
2	PO4	Н	1704	-	4,4,4	0.92	0	6,6,6	0.43	0
2	PO4	Ι	1702	-	4,4,4	0.92	0	6,6,6	0.43	0
3	K6P	J	1701[A]	-	18,18,18	0.18	0	23,24,24	0.45	0
2	PO4	D	1702	-	4,4,4	0.92	0	6,6,6	0.43	0



Mal	Turne	Chain	Dec	Tinle	Bo	ond leng	$_{\rm ths}$	B	ond ang	les
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
3	K6P	Е	1701[B]	-	18,18,18	0.18	0	23,24,24	0.35	0
2	PO4	D	1703	-	4,4,4	0.92	0	6,6,6	0.45	0
2	PO4	В	1704	-	4,4,4	0.94	0	6,6,6	0.42	0
2	PO4	J	1702	-	4,4,4	0.93	0	6,6,6	0.43	0
4	FMT	В	1702	-	2,2,2	0.61	0	1,1,1	0.61	0
4	FMT	Н	1702	-	2,2,2	0.62	0	1,1,1	0.61	0
3	K6P	С	1701[B]	-	18,18,18	0.17	0	23,24,24	0.34	0
2	PO4	С	1702	-	4,4,4	0.93	0	6,6,6	0.43	0
2	PO4	J	1703	-	4,4,4	0.93	0	6,6,6	0.45	0
2	PO4	D	1704	-	4,4,4	0.93	0	6,6,6	0.43	0
2	PO4	А	1701	-	4,4,4	0.90	0	6,6,6	0.43	0
2	PO4	G	1703	-	4,4,4	0.92	0	6,6,6	0.43	0
3	K6P	J	1701[B]	-	18,18,18	0.18	0	23,24,24	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	K6P	С	1701[A]	-	-	3/13/13/13	0/1/1/1
3	K6P	В	1701[B]	-	-	2/13/13/13	0/1/1/1
3	K6P	J	1701[B]	-	-	0/13/13/13	0/1/1/1
3	K6P	С	1701[B]	-	-	2/13/13/13	0/1/1/1
3	K6P	G	1701[A]	-	-	0/13/13/13	0/1/1/1
3	K6P	J	1701[A]	-	-	3/13/13/13	0/1/1/1
3	K6P	Е	1701[B]	-	-	0/13/13/13	0/1/1/1
3	K6P	В	1701[A]	-	-	2/13/13/13	0/1/1/1
3	K6P	G	1701[B]	-	-	0/13/13/13	0/1/1/1
3	K6P	Е	1701[A]	-	-	0/13/13/13	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	1701[A]	K6P	C6-C7-N1-C8
3	В	1701[A]	K6P	O1-C7-N1-C8
3	С	1701[A]	K6P	C6-C7-N1-C8



Mol	Chain	Res	Type	Atoms
3	С	1701[A]	K6P	O1-C7-N1-C8
3	J	1701[A]	K6P	C6-C7-N1-C8
3	J	1701[A]	K6P	O1-C7-N1-C8
3	С	1701[A]	K6P	N1-C8-C9-C10
3	В	1701[B]	K6P	O1-C7-N1-C8
3	В	1701[B]	K6P	C6-C7-N1-C8
3	С	1701[B]	K6P	O1-C7-N1-C8
3	С	1701[B]	K6P	C6-C7-N1-C8
3	J	1701[A]	K6P	N1-C8-C9-C10

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	С	1701[A]	K6P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	120/123~(97%)	0.40	4 (3%) 46 53	30, 44, 60, 132	2(1%)
1	В	119/123~(96%)	0.47	8 (6%) 17 23	35, 49, 74, 101	3(2%)
1	С	121/123~(98%)	0.53	8 (6%) 18 23	31, 47, 82, 139	1 (0%)
1	D	120/123~(97%)	0.53	10 (8%) 11 15	34, 51, 78, 101	2(1%)
1	Ε	120/123~(97%)	0.68	17 (14%) 2 3	34, 51, 93, 115	3(2%)
1	F	120/123~(97%)	0.43	9 (7%) 14 19	34, 50, 71, 108	5 (4%)
1	G	119/123~(96%)	0.70	12 (10%) 7 9	37, 56, 98, 121	5(4%)
1	Н	119/123~(96%)	0.54	13 (10%) 5 8	37, 53, 85, 123	2(1%)
1	Ι	118/123~(95%)	0.90	20 (16%) 1 2	44, 67, 98, 135	3(2%)
1	J	$12\overline{0/123}~(97\%)$	0.53	11 (9%) 9 12	38, 56, 86, 129	5 (4%)
All	All	1196/1230~(97%)	0.57	112 (9%) 8 11	30, 52, 88, 139	31 (2%)

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type RSRZ	
1	J	1484	ASN	6.1
1	Ι	1512	ALA	6.1
1	А	1603	GLY	6.0
1	Н	1499	GLY	5.4
1	G	1495[A]	TRP	5.1
1	С	1495[A]	TRP	5.1
1	D	1603	GLY	5.0
1	Е	1485	SER	5.0
1	G	1512	ALA	4.8
1	В	1498[A]	ASN	4.6
1	Ι	1602	LEU	4.6
1	G	1513	GLY	4.3
1	D	1499[A]	GLY	4.1



Mol	Chain	Res	Type	RSRZ
1	С	1482	HIS	4.1
1	F	1499[A]	GLY	4.0
1	Е	1484	ASN	3.9
1	F	1603	GLY	3.9
1	F	1512	ALA	3.8
1	Ι	1529	GLY	3.7
1	Ι	1532	ILE	3.7
1	В	1497[A]	SER	3.7
1	Ι	1513	GLY	3.7
1	Е	1511	GLY	3.6
1	Е	1513	GLY	3.6
1	Ι	1528	LEU	3.5
1	Ι	1575	GLU	3.5
1	Ι	1510	VAL	3.5
1	G	1497[A]	SER	3.5
1	J	1528	LEU	3.4
1	J	1483	MET	3.4
1	Е	1495	TRP	3.4
1	J	1495[A]	TRP	3.3
1	Е	1512	ALA	3.3
1	G	1532	ILE	3.3
1	F	1498[A]	ASN	3.3
1	G	1494[A]	LYS	3.3
1	Ι	1486	PHE	3.3
1	J	1602	LEU	3.2
1	D	1498[A]	ASN	3.2
1	G	1498[A]	ASN	3.2
1	Н	1495[A]	TRP	3.1
1	Е	1575	GLU	3.0
1	Е	1510	VAL	3.0
1	G	1527	VAL	3.0
1	Ι	1576	GLY	3.0
1	С	1512	ALA	3.0
1	В	1495[A]	TRP	2.9
1	С	1510	VAL	2.9
1	Е	1486	PHE	2.9
1	D	1495[A]	TRP	2.9
1	G	1486	PHE	2.8
1	А	1484	ASN	2.8
1	G	1575[A]	GLU	2.8
1	Ι	1495	TRP	2.8
1	Н	1498[A]	ASN	2.8



6MXX

Mol	Chain Res		Type RSR2	
1	Н	1496[A]	SER	2.8
1	Н	1497[A]	SER	2.8
1	Н	1575	GLU	2.8
1	G	1515	TYR	2.7
1	Ι	1485	SER	2.7
1	Н	1513	GLY	2.7
1	Ι	1498	ASN	2.7
1	Н	1528	LEU	2.7
1	J	1493[A]	ALA	2.7
1	D	1497[A]	SER	2.6
1	С	1497[A]	SER	2.6
1	А	1495[A]	TRP	2.6
1	Ι	1515	TYR	2.5
1	Е	1494	LYS	2.5
1	Н	1494[A]	LYS	2.4
1	В	1496[A]	SER	2.4
1	D	1496[A]	SER	2.4
1	Ι	1496	SER	2.4
1	В	1512	ALA	2.4
1	Е	1508[A]	ARG	2.4
1	Е	1528	LEU	2.4
1	Ι	1574	LYS	2.3
1	Е	1499	GLY	2.3
1	Ι	1499	GLY	2.3
1	Ι	1601	GLY	2.3
1	Н	1512	ALA	2.3
1	D	1574	LYS	2.3
1	E	1517[A]	LEU	2.3
1	Ι	1487	VAL	2.3
1	D	1492[A]	VAL	2.3
1	H	1486	PHE	2.3
1	A	1517[A]	LEU	2.3
1	E	1535	CYS	2.3
1	J	1552	TYR	2.3
1	C	1484	ASN	2.2
1	C	1517[A]	LEU	2.2
1	E	1498	ASN	2.2
1	E	1496	SER	2.2
1	J	1486	PHE	2.2
1	F	1495[A]	TRP	2.2
1	F	1576	GLY	2.2
1	I	1590	LEU	2.2



Mol	Chain	Res	Type	RSRZ	
1	J	1517[A]	LEU	2.2	
1	F	1584[A]	MET	2.2	
1	В	1517[A]	LEU	2.1	
1	J	1498[A]	ASN	2.1	
1	F	1575	GLU	2.1	
1	F	1513	GLY	2.1	
1	Н	1484	ASN	2.1	
1	Н	1492	VAL	2.1	
1	С	1591	GLU	2.1	
1	D	1517[A]	LEU	2.1	
1	J	1497[A]	SER	2.0	
1	G	1508	ARG	2.0	
1	D	1494[A]	LYS	2.0	
1	В	1590	LEU	2.0	
1	В	1575	GLU	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
2	PO4	D	1703	5/5	0.64	0.43	139,139,140,141	0
2	PO4	G	1703	5/5	0.67	0.29	141,142,143,143	0
2	PO4	Е	1702	5/5	0.68	0.84	260,260,261,262	0
2	PO4	Ι	1702	5/5	0.68	0.29	144,145,145,145	0
2	PO4	В	1703	5/5	0.72	0.52	$152,\!153,\!154,\!155$	0
2	PO4	С	1702	5/5	0.75	0.47	173,173,174,175	0
3	K6P	J	1701[A]	18/18	0.76	0.52	148,183,203,205	39
3	K6P	J	1701[B]	18/18	0.76	0.52	149,183,212,212	39



6MXX

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	$Q{<}0.9$
3	K6P	В	1701[A]	18/18	0.77	0.54	$119,\!146,\!166,\!167$	39
3	K6P	В	1701[B]	18/18	0.77	0.54	121,147,171,171	39
4	FMT	Н	1701	3/3	0.77	0.21	67,68,70,84	0
3	K6P	G	1701[B]	18/18	0.81	0.50	133,166,179,179	39
3	K6P	G	1701[A]	18/18	0.81	0.50	132,166,179,179	39
2	PO4	D	1701	5/5	0.82	0.35	130,131,131,133	0
2	PO4	J	1703	5/5	0.84	0.25	105,109,112,112	0
2	PO4	F	1701	5/5	0.84	0.53	$196,\!198,\!198,\!199$	0
3	K6P	С	1701[B]	18/18	0.85	0.38	87,104,120,122	39
2	PO4	J	1702	5/5	0.85	0.25	139,139,139,139	0
3	K6P	С	1701[A]	18/18	0.85	0.38	82,100,120,127	39
4	FMT	В	1702	3/3	0.86	0.13	72,73,74,87	0
2	PO4	В	1704	5/5	0.87	0.23	93,94,95,96	0
4	FMT	Н	1703	3/3	0.87	0.18	79,80,81,97	0
2	PO4	С	1703	5/5	0.88	0.27	109,112,114,114	0
3	K6P	Е	1701[A]	18/18	0.89	0.36	92,112,115,115	39
3	K6P	Е	1701[B]	18/18	0.89	0.36	92,111,114,116	39
2	PO4	D	1702	5/5	0.89	0.53	180,181,182,184	0
2	PO4	D	1704	5/5	0.90	0.23	100,103,104,106	0
2	PO4	Ι	1701	5/5	0.90	0.18	98,98,98,101	0
2	PO4	Е	1703	5/5	0.91	0.23	93,93,96,97	0
2	PO4	G	1702	5/5	0.91	0.24	96,96,97,101	0
2	PO4	A	1701	5/5	0.92	0.23	93,98,101,102	0
2	PO4	Н	1705	5/5	0.93	0.27	88,89,92,93	0
2	PO4	F	1702	5/5	0.93	0.24	86,87,89,89	0
2	PO4	Н	1704	5/5	0.93	0.19	121,121,122,122	0
4	FMT	Н	1702	3/3	0.97	0.30	52,52,52,63	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

























6.5 Other polymers (i)

There are no such residues in this entry.

