



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2023 – 11:02 PM EDT

PDB ID : 6UIB  
Title : Crystal structure of IL23 bound to peptide 23-652  
Authors : Durbin, J.D.; Wang, J.; Afshar, S.  
Deposited on : 2019-09-30  
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	
2	B	316	
3	C	20	
4	D	2	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-23 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	132	963	619	170	169	5	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	ASP	-	expression tag	UNP Q9NPF7
A	-2	ARG	-	expression tag	UNP Q9NPF7
A	-1	SER	-	expression tag	UNP Q9NPF7
A	0	LEU	-	expression tag	UNP Q9NPF7
A	171	HIS	-	expression tag	UNP Q9NPF7
A	172	HIS	-	expression tag	UNP Q9NPF7
A	173	HIS	-	expression tag	UNP Q9NPF7
A	174	HIS	-	expression tag	UNP Q9NPF7
A	175	HIS	-	expression tag	UNP Q9NPF7
A	176	HIS	-	expression tag	UNP Q9NPF7

- Molecule 2 is a protein called Interleukin-12 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	274	2095	1327	341	416	11	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	ASP	-	expression tag	UNP P29460
B	-2	ARG	-	expression tag	UNP P29460
B	-1	SER	-	expression tag	UNP P29460
B	0	LEU	-	expression tag	UNP P29460
B	307	HIS	-	expression tag	UNP P29460
B	308	HIS	-	expression tag	UNP P29460

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	309	HIS	-	expression tag	UNP P29460
B	310	HIS	-	expression tag	UNP P29460
B	311	HIS	-	expression tag	UNP P29460
B	312	HIS	-	expression tag	UNP P29460

- Molecule 3 is a protein called Peptide 23-652.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
			Total	C	N	O				S
3	C	15	122	81	17	21	3	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	2	28	16	2	10	0	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	O	0	0
			5	5		
5	B	20	Total	O	0	0
			20	20		
5	C	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin-23 subunit alpha



- Molecule 2: Interleukin-12 subunit beta



- Molecule 3: Peptide 23-652



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.59Å 94.67Å 101.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.90 – 2.74 19.73 – 2.74	Depositor EDS
% Data completeness (in resolution range)	97.5 (19.90-2.74) 97.5 (19.73-2.74)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.70 (at 2.75Å)	Xtrriage
Refinement program	BUSTER 2.11.7	Depositor
R, $R_{free}$	0.206 , 0.241 0.218 , 0.254	Depositor DCC
$R_{free}$ test set	1003 reflections (5.39%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	72.3	Xtrriage
Anisotropy	0.532	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 70.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3234	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/984	0.65	0/1341
2	B	0.46	0/2139	0.72	0/2915
3	C	0.50	0/127	0.70	0/171
All	All	0.48	0/3250	0.70	0/4427

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	963	0	930	8	0
2	B	2095	0	1922	8	0
3	C	122	0	97	1	0
4	D	28	0	25	0	0
5	A	5	0	0	0	0
5	B	20	0	0	0	0
5	C	1	0	0	0	0
All	All	3234	0	2974	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:221:LEU:HD13	2:B:303:VAL:HG21	1.68	0.74
2:B:224:LEU:HD12	2:B:229:GLN:HG3	1.86	0.58
2:B:134:VAL:HG22	2:B:190:VAL:HG22	1.89	0.54
2:B:133:SER:HB3	2:B:191:ASP:HB2	1.90	0.53
1:A:8:SER:HB2	1:A:9:PRO:HD2	1.89	0.53
2:B:134:VAL:HG11	2:B:169:VAL:HG11	1.93	0.50
1:A:67:SER:O	1:A:71:LEU:HG	2.15	0.46
1:A:155:ALA:O	1:A:159:ARG:HG3	2.15	0.46
2:B:223:PRO:HA	2:B:230:VAL:HG12	2.01	0.43
2:B:221:LEU:HD13	2:B:303:VAL:CG2	2.43	0.42
1:A:30:PRO:HG3	3:C:7:PHE:CE1	2.55	0.42
1:A:81:TYR:CG	1:A:150:LEU:HD13	2.55	0.41
2:B:221:LEU:HD21	2:B:284:ILE:HD12	2.00	0.41
1:A:138:GLN:H	1:A:138:GLN:HG2	1.58	0.41
1:A:15:GLN:CG	1:A:162:ALA:HA	2.50	0.41
1:A:50:PRO:HG2	1:A:77:GLY:HA2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	126/180 (70%)	120 (95%)	6 (5%)	0	100	100
2	B	258/316 (82%)	243 (94%)	14 (5%)	1 (0%)	34	55
3	C	13/20 (65%)	13 (100%)	0	0	100	100
All	All	397/516 (77%)	376 (95%)	20 (5%)	1 (0%)	41	61

All (1) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
2	B	20	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/154 (63%)	91 (94%)	6 (6%)	18	32
2	B	225/287 (78%)	214 (95%)	11 (5%)	25	43
3	C	12/18 (67%)	10 (83%)	2 (17%)	2	3
All	All	334/459 (73%)	315 (94%)	19 (6%)	20	36

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	17	LEU
1	A	47	ASN
1	A	110	LEU
1	A	138	GLN
1	A	148	ARG
2	B	0	LEU
2	B	8	VAL
2	B	13	LEU
2	B	29	ASP
2	B	103	ASN
2	B	152	THR
2	B	204	SER
2	B	227	SER
2	B	265	ASP
2	B	278	CYS
2	B	295	SER
3	C	6	SER
3	C	14	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	56	GLN
2	B	229	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	D	1	2,4	14,14,15	0.33	0	17,19,21	0.54	0
4	NAG	D	2	4	14,14,15	0.30	0	17,19,21	0.97	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	2	NAG	C1-O5-C5	2.97	116.21	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	132/180 (73%)	0.22	8 (6%) 21 24	53, 82, 123, 142	0
2	B	274/316 (86%)	0.15	13 (4%) 31 34	51, 83, 129, 173	0
3	C	15/20 (75%)	0.15	0 100 100	54, 64, 91, 100	0
All	All	421/516 (81%)	0.17	21 (4%) 28 32	51, 82, 127, 173	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	ASP	4.0
2	B	11	VAL	3.8
2	B	43	SER	3.7
2	B	42	GLN	3.2
2	B	259	SER	3.1
1	A	8	SER	2.9
2	B	103	ASN	2.8
2	B	67	THR	2.8
1	A	65	ASP	2.6
1	A	141	LEU	2.6
2	B	19	ALA	2.6
2	B	288	ALA	2.5
1	A	110	LEU	2.4
1	A	34	HIS	2.3
2	B	258	LYS	2.3
2	B	52	THR	2.3
2	B	128	THR	2.3
1	A	74	ILE	2.3
1	A	114	GLN	2.2
2	B	25	VAL	2.2
2	B	105	THR	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	D	2	14/15	0.93	0.28	93,96,99,102	0
4	NAG	D	1	14/15	0.96	0.14	69,82,89,90	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.