



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 01:47 AM EDT

PDB ID : 6ZD1  
Title : Structure of apo telomerase from *Candida Tropicalis*  
Authors : Zhai, L.; Rety, S.; Chen, W.F.; Auguin, D.; Xi, X.G.  
Deposited on : 2020-06-13  
Resolution : 2.47 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5857 (2.50-2.46)
Clashscore	141614	6594 (2.50-2.46)
Ramachandran outliers	138981	6469 (2.50-2.46)
Sidechain outliers	138945	6471 (2.50-2.46)
RSRZ outliers	127900	5738 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	705	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5305 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Telomerase reverse transcriptase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	645	5291	3410	893	967	10	11	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	880	VAL	-	expression tag	UNP C5MCQ7
A	881	ASP	-	expression tag	UNP C5MCQ7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	14	Total	O	0	0
			14	14		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Telomerase reverse transcriptase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.80Å 65.80Å 387.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.78 – 2.47 96.99 – 2.47	Depositor EDS
% Data completeness (in resolution range)	85.1 (43.78-2.47) 85.1 (96.99-2.47)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 2.45Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874, PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.246 , 0.284 0.246 , 0.284	Depositor DCC
$R_{free}$ test set	1417 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	67.8	Xtrriage
Anisotropy	0.047	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5305	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	82.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/5370	0.47	2/7196 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	245	LEU	C-N-CA	6.15	137.08	121.70
1	A	246	LYS	N-CA-C	5.05	124.63	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	246	LYS	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5291	0	5436	76	0
2	A	14	0	0	0	0
All	All	5305	0	5436	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:676:LEU:HD23	1:A:730:VAL:HG21	1.32	1.08
1:A:415:LYS:HG3	1:A:420:ARG:HG3	1.59	0.84
1:A:366:ARG:HD3	1:A:776:PHE:CZ	2.15	0.80
1:A:248:MSE:HE1	1:A:380:LEU:HD22	1.66	0.78
1:A:363:ASN:H	1:A:366:ARG:HE	1.35	0.74
1:A:624:ASN:HD22	1:A:628:GLN:HB2	1.51	0.73
1:A:245:LEU:C	1:A:246:LYS:HG3	2.12	0.70
1:A:717:ILE:HG12	1:A:727:MSE:HE3	1.73	0.69
1:A:420:ARG:HE	1:A:422:ILE:HD11	1.59	0.67
1:A:676:LEU:HD23	1:A:730:VAL:CG2	2.18	0.67
1:A:415:LYS:HG3	1:A:420:ARG:CG	2.26	0.65
1:A:228:ARG:HD2	1:A:231:LEU:HD23	1.77	0.64
1:A:302:LEU:HD13	1:A:337:GLU:HB3	1.79	0.64
1:A:245:LEU:O	1:A:246:LYS:HG3	1.99	0.62
1:A:482:VAL:HG13	1:A:487:GLU:HB3	1.82	0.62
1:A:365:ASP:OD2	1:A:368:LYS:NZ	2.31	0.62
1:A:873:TRP:O	1:A:876:THR:HG22	2.01	0.61
1:A:363:ASN:N	1:A:366:ARG:HE	2.00	0.60
1:A:777:LEU:HD22	1:A:791:ASN:ND2	2.17	0.59
1:A:379:GLU:OE1	1:A:383:ASN:ND2	2.33	0.58
1:A:553:LEU:HD22	1:A:838:ARG:HA	1.85	0.58
1:A:553:LEU:HA	1:A:843:MSE:HE2	1.84	0.58
1:A:624:ASN:ND2	1:A:628:GLN:HB2	2.20	0.56
1:A:766:LYS:NZ	1:A:770:CYS:SG	2.68	0.56
1:A:556:PHE:O	1:A:560:LYS:N	2.38	0.55
1:A:362:SER:OG	1:A:415:LYS:HG2	2.07	0.55
1:A:565:LEU:HD12	1:A:830:GLU:HA	1.88	0.55
1:A:776:PHE:CE1	1:A:777:LEU:HG	2.43	0.53
1:A:541:GLU:HA	1:A:604:LYS:HD3	1.90	0.53
1:A:247:THR:OG1	1:A:250:ASP:OD2	2.27	0.52
1:A:624:ASN:ND2	1:A:630:ASP:OD2	2.43	0.52
1:A:750:ARG:HG2	1:A:752:THR:H	1.75	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:ASN:H	1:A:366:ARG:NE	2.05	0.50
1:A:539:ASP:OD2	1:A:545:TYR:OH	2.23	0.50
1:A:660:GLU:HB3	1:A:696:LYS:CE	2.41	0.50
1:A:624:ASN:ND2	1:A:628:GLN:OE1	2.45	0.50
1:A:549:GLU:HB2	1:A:600:LEU:HG	1.94	0.50
1:A:313:LYS:H	1:A:315:LYS:NZ	2.10	0.49
1:A:867:LYS:HB2	1:A:871:ILE:HD13	1.95	0.49
1:A:539:ASP:O	1:A:543:THR:OG1	2.28	0.49
1:A:273:ASP:HB2	1:A:315:LYS:HE3	1.95	0.49
1:A:662:LYS:HA	1:A:665:TRP:CZ3	2.49	0.48
1:A:562:VAL:HG22	1:A:563:ARG:H	1.78	0.48
1:A:553:LEU:HG	1:A:557:LEU:HD21	1.96	0.48
1:A:554:ASP:HA	1:A:557:LEU:CD1	2.43	0.48
1:A:407:ASN:ND2	1:A:619:THR:OG1	2.47	0.47
1:A:504:LYS:HD3	1:A:504:LYS:HA	1.74	0.46
1:A:363:ASN:O	1:A:366:ARG:NE	2.49	0.46
1:A:655:THR:HG22	1:A:659:LYS:HE2	1.98	0.45
1:A:368:LYS:HB3	1:A:370:PHE:CZ	2.51	0.45
1:A:309:ILE:H	1:A:309:ILE:HG12	1.64	0.45
1:A:660:GLU:HB3	1:A:696:LYS:NZ	2.32	0.45
1:A:343:VAL:O	1:A:347:LEU:HB2	2.18	0.44
1:A:777:LEU:HD23	1:A:777:LEU:HA	1.79	0.44
1:A:780:PHE:CZ	1:A:785:LEU:HG	2.52	0.44
1:A:554:ASP:HA	1:A:557:LEU:HG	2.00	0.44
1:A:210:PHE:HB3	1:A:211:ARG:H	1.62	0.44
1:A:216:VAL:HG21	1:A:315:LYS:HA	2.00	0.43
1:A:776:PHE:CD1	1:A:777:LEU:HG	2.52	0.43
1:A:597:VAL:HG12	1:A:599:THR:HG23	2.00	0.43
1:A:245:LEU:HA	1:A:251:TYR:OH	2.20	0.42
1:A:502:LYS:HE2	1:A:502:LYS:HB2	1.81	0.42
1:A:554:ASP:N	1:A:554:ASP:OD1	2.51	0.42
1:A:787:THR:O	1:A:791:ASN:ND2	2.52	0.42
1:A:554:ASP:HB3	1:A:558:LYS:HD2	2.02	0.42
1:A:405:PHE:HZ	1:A:448:ASP:HB3	1.85	0.42
1:A:187:VAL:O	1:A:191:ILE:HG13	2.20	0.42
1:A:550:TYR:CE2	1:A:844:GLU:HG3	2.55	0.42
1:A:506:LEU:HD12	1:A:506:LEU:HA	1.92	0.41
1:A:536:PHE:HB3	1:A:604:LYS:HE3	2.01	0.41
1:A:385:ILE:HG13	1:A:412:LEU:HD11	2.01	0.41
1:A:565:LEU:HD11	1:A:833:ILE:HG13	2.03	0.41
1:A:715:VAL:HG12	1:A:727:MSE:HE2	2.03	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:410:ILE:HD11	1:A:620:CYS:SG	2.61	0.41
1:A:376:ILE:HD12	1:A:376:ILE:HA	1.94	0.40
1:A:879:ILE:H	1:A:879:ILE:HG13	1.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	635/705 (90%)	617 (97%)	16 (2%)	2 (0%)	41 59

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	677	VAL
1	A	554	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	599/640 (94%)	584 (98%)	15 (2%)	47 71

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	246	LYS
1	A	249	ILE
1	A	257	ARG
1	A	309	ILE
1	A	415	LYS
1	A	416	ARG
1	A	420	ARG
1	A	475	ASP
1	A	554	ASP
1	A	555	GLU
1	A	629	TYR
1	A	671	ASN
1	A	780	PHE
1	A	789	ARG
1	A	830	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	624	ASN
1	A	628	GLN
1	A	791	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	634/705 (89%)	0.41	34 (5%) <span style="border: 1px solid red; padding: 2px;">25</span> <span style="border: 1px solid red; padding: 2px;">26</span>	48, 79, 117, 144	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	401	SER	8.8
1	A	556	PHE	5.7
1	A	596	LYS	5.3
1	A	753	CYS	4.2
1	A	561	ARG	3.8
1	A	399	PRO	3.5
1	A	752	THR	3.3
1	A	437	ARG	3.2
1	A	403	GLY	3.2
1	A	777	LEU	3.2
1	A	404	GLN	3.1
1	A	553	LEU	2.9
1	A	557	LEU	2.9
1	A	402	LYS	2.9
1	A	628	GLN	2.7
1	A	597	VAL	2.7
1	A	631	LEU	2.6
1	A	594	VAL	2.6
1	A	565	LEU	2.5
1	A	677	VAL	2.5
1	A	564	THR	2.4
1	A	755	PHE	2.4
1	A	421	VAL	2.4
1	A	600	LEU	2.4
1	A	332	ARG	2.3
1	A	376	ILE	2.3
1	A	315	LYS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	563	ARG	2.3
1	A	559	LEU	2.3
1	A	541	GLU	2.3
1	A	554	ASP	2.2
1	A	625	LYS	2.2
1	A	667	ALA	2.1
1	A	567	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.