



wwPDB EM Validation Summary Report ⓘ

Jul 8, 2024 – 02:32 pm BST

PDB ID : 7PBX
EMDB ID : EMD-13308
Title : Cryo-EM structure of the GroEL-GroES complex with ADP bound to both rings ("tight" conformation).
Authors : Pichkur, E.B.; Stanishneva-Konovalova, T.B.
Deposited on : 2021-08-02
Resolution : 3.43 Å(reported)
Based on initial model : 1SX4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Ac	524	
1	Ad	524	
1	Ai	524	
1	Aj	524	
1	Ao	524	
1	Ap	524	
1	Au	524	
1	Av	524	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain			
1	Ba	524	36%	88%	10%	•
1	Bb	524	8%	94%	6%	•
1	Bg	524	36%	88%	10%	•
1	Bh	524	8%	94%	6%	•
1	Bm	524	36%	88%	10%	•
1	Bn	524	8%	94%	6%	•
2	Af	97	30%	65%	26%	5% •
2	Al	97	30%	65%	26%	5% •
2	Ar	97	30%	65%	26%	5% •
2	Ax	97	31%	65%	26%	5% •
2	Bd	97	30%	65%	26%	5% •
2	Bj	97	30%	65%	26%	5% •
2	Bp	97	30%	65%	26%	5% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ADP	Ad	601	X	-	-	-
3	ADP	Aj	601	X	-	-	-
3	ADP	Ap	601	X	-	-	-
3	ADP	Av	601	X	-	-	-
3	ADP	Bb	601	X	-	-	-
3	ADP	Bh	601	X	-	-	-
3	ADP	Bn	601	X	-	-	-

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 59472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60 kDa chaperonin.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	Ac	524	3856	2397	665	774	20	0	0
1	Ad	524	3856	2397	665	774	20	0	0
1	Ai	524	3856	2397	665	774	20	0	0
1	Aj	524	3856	2397	665	774	20	0	0
1	Ao	524	3856	2397	665	774	20	0	0
1	Ap	524	3856	2397	665	774	20	0	0
1	Au	524	3856	2397	665	774	20	0	0
1	Av	524	3856	2397	665	774	20	0	0
1	Ba	524	3856	2397	665	774	20	0	0
1	Bb	524	3856	2397	665	774	20	0	0
1	Bg	524	3856	2397	665	774	20	0	0
1	Bh	524	3856	2397	665	774	20	0	0
1	Bm	524	3856	2397	665	774	20	0	0
1	Bn	524	3856	2397	665	774	20	0	0

- Molecule 2 is a protein called 10 kDa chaperonin.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	Af	97	728	454	127	145	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	Al	97	Total 728	C 454	N 127	O 145	S 2	0	0
2	Ar	97	Total 728	C 454	N 127	O 145	S 2	0	0
2	Ax	97	Total 728	C 454	N 127	O 145	S 2	0	0
2	Bd	97	Total 728	C 454	N 127	O 145	S 2	0	0
2	Bj	97	Total 728	C 454	N 127	O 145	S 2	0	0
2	Bp	97	Total 728	C 454	N 127	O 145	S 2	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
3	Ac	1	Total 27	C 10	N 5	O 10	P 2	0
3	Ad	1	Total 27	C 10	N 5	O 10	P 2	0
3	Ai	1	Total 27	C 10	N 5	O 10	P 2	0
3	Aj	1	Total 27	C 10	N 5	O 10	P 2	0
3	Ao	1	Total 27	C 10	N 5	O 10	P 2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
3	Ap	1	27	10	5	10	2	0
3	Au	1	27	10	5	10	2	0
3	Av	1	27	10	5	10	2	0
3	Ba	1	27	10	5	10	2	0
3	Bb	1	27	10	5	10	2	0
3	Bg	1	27	10	5	10	2	0
3	Bh	1	27	10	5	10	2	0
3	Bm	1	27	10	5	10	2	0
3	Bn	1	27	10	5	10	2	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
4	Ac	1	1	1	0
4	Ad	1	1	1	0
4	Ai	1	1	1	0
4	Aj	1	1	1	0
4	Ao	1	1	1	0
4	Ap	1	1	1	0
4	Au	1	1	1	0
4	Av	1	1	1	0
4	Ba	1	1	1	0
4	Bb	1	1	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
4	Bg	1	Total 1	Mg 1	0
4	Bh	1	Total 1	Mg 1	0
4	Bm	1	Total 1	Mg 1	0
4	Bn	1	Total 1	Mg 1	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin




- Molecule 1: 60 kDa chaperonin







• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin










• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin





• Molecule 1: 60 kDa chaperonin








• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin







• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin




• Molecule 1: 60 kDa chaperonin







• Molecule 2: 10 kDa chaperonin




• Molecule 2: 10 kDa chaperonin




• Molecule 2: 10 kDa chaperonin




• Molecule 2: 10 kDa chaperonin







• Molecule 2: 10 kDa chaperonin



• Molecule 2: 10 kDa chaperonin



• Molecule 2: 10 kDa chaperonin



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	41000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	100	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	5.566	Depositor
Minimum map value	-1.036	Depositor
Average map value	0.023	Depositor
Map value standard deviation	0.187	Depositor
Recommended contour level	1.5	Depositor
Map size (Å)	425.088, 425.088, 425.088	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.107, 1.107, 1.107	Depositor

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	Ac	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Ad	0.64	1/3884 (0.0%)	0.92	10/5243 (0.2%)
1	Ai	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Aj	0.64	1/3884 (0.0%)	0.92	9/5243 (0.2%)
1	Ao	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Ap	0.64	1/3884 (0.0%)	0.92	10/5243 (0.2%)
1	Au	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Av	0.64	1/3884 (0.0%)	0.92	9/5243 (0.2%)
1	Ba	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Bb	0.64	1/3884 (0.0%)	0.92	9/5243 (0.2%)
1	Bg	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Bh	0.64	1/3884 (0.0%)	0.92	9/5243 (0.2%)
1	Bm	1.35	5/3884 (0.1%)	1.15	31/5243 (0.6%)
1	Bn	0.64	1/3884 (0.0%)	0.92	8/5243 (0.2%)
2	Af	4.06	14/732 (1.9%)	2.02	34/983 (3.5%)
2	Al	4.06	14/732 (1.9%)	2.02	34/983 (3.5%)
2	Ar	4.06	14/732 (1.9%)	2.03	34/983 (3.5%)
2	Ax	4.06	14/732 (1.9%)	2.03	34/983 (3.5%)
2	Bd	4.06	14/732 (1.9%)	2.02	34/983 (3.5%)
2	Bj	4.06	14/732 (1.9%)	2.02	34/983 (3.5%)
2	Bp	4.06	14/732 (1.9%)	2.02	34/983 (3.5%)
All	All	1.56	140/59500 (0.2%)	1.16	519/80283 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Ac	0	5
1	Ad	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Ai	0	5
1	Aj	0	1
1	Ao	0	5
1	Ap	0	1
1	Au	0	5
1	Av	0	1
1	Ba	0	5
1	Bb	0	1
1	Bg	0	5
1	Bh	0	1
1	Bm	0	5
1	Bn	0	1
2	Af	0	3
2	Al	0	3
2	Ar	0	3
2	Ax	0	3
2	Bd	0	3
2	Bj	0	3
2	Bp	0	3
All	All	0	63

The worst 5 of 140 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	Bp	96	GLU	CB-CG	82.59	3.09	1.52
2	Ar	96	GLU	CB-CG	82.57	3.09	1.52
2	Ax	96	GLU	CB-CG	82.57	3.09	1.52
2	Af	96	GLU	CB-CG	82.56	3.09	1.52
2	Bj	96	GLU	CB-CG	82.56	3.09	1.52

The worst 5 of 519 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Ar	94	ILE	CB-CA-C	14.60	140.79	111.60
2	Bj	94	ILE	CB-CA-C	14.59	140.78	111.60
2	Af	94	ILE	CB-CA-C	14.59	140.78	111.60
2	Bd	94	ILE	CB-CA-C	14.59	140.78	111.60
2	Al	94	ILE	CB-CA-C	14.59	140.78	111.60

There are no chirality outliers.

5 of 63 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	Ac	213	VAL	Peptide
1	Ac	255	GLU	Peptide
1	Ac	285	ARG	Sidechain
1	Ac	354	GLU	Peptide
1	Ac	360	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Ac	3856	0	3975	0	0
1	Ad	3856	0	3976	0	0
1	Ai	3856	0	3975	0	0
1	Aj	3856	0	3976	0	0
1	Ao	3856	0	3975	0	0
1	Ap	3856	0	3976	0	0
1	Au	3856	0	3975	0	0
1	Av	3856	0	3976	0	0
1	Ba	3856	0	3975	0	0
1	Bb	3856	0	3976	0	0
1	Bg	3856	0	3975	0	0
1	Bh	3856	0	3976	0	0
1	Bm	3856	0	3975	0	0
1	Bn	3856	0	3976	0	0
2	Af	728	0	761	0	0
2	Al	728	0	761	0	0
2	Ar	728	0	761	0	0
2	Ax	728	0	761	0	0
2	Bd	728	0	761	0	0
2	Bj	728	0	761	0	0
2	Bp	728	0	761	0	0
3	Ac	27	0	11	0	0
3	Ad	27	0	11	0	0
3	Ai	27	0	11	0	0
3	Aj	27	0	11	0	0
3	Ao	27	0	11	0	0
3	Ap	27	0	11	0	0
3	Au	27	0	11	0	0
3	Av	27	0	11	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	Ba	27	0	11	0	0
3	Bb	27	0	11	0	0
3	Bg	27	0	11	0	0
3	Bh	27	0	11	0	0
3	Bm	27	0	11	0	0
3	Bn	27	0	11	0	0
4	Ac	1	0	0	0	0
4	Ad	1	0	0	0	0
4	Ai	1	0	0	0	0
4	Aj	1	0	0	0	0
4	Ao	1	0	0	0	0
4	Ap	1	0	0	0	0
4	Au	1	0	0	0	0
4	Av	1	0	0	0	0
4	Ba	1	0	0	0	0
4	Bb	1	0	0	0	0
4	Bg	1	0	0	0	0
4	Bh	1	0	0	0	0
4	Bm	1	0	0	0	0
4	Bn	1	0	0	0	0
All	All	59472	0	61138	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 32.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	Ac	522/524 (100%)	494 (95%)	22 (4%)	6 (1%)	14 50
1	Ad	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15 52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Ai	522/524 (100%)	494 (95%)	22 (4%)	6 (1%)	14	50
1	Aj	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
1	Ao	522/524 (100%)	493 (94%)	23 (4%)	6 (1%)	14	50
1	Ap	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
1	Au	522/524 (100%)	494 (95%)	22 (4%)	6 (1%)	14	50
1	Av	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
1	Ba	522/524 (100%)	493 (94%)	23 (4%)	6 (1%)	14	50
1	Bb	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
1	Bg	522/524 (100%)	493 (94%)	23 (4%)	6 (1%)	14	50
1	Bh	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
1	Bm	522/524 (100%)	494 (95%)	22 (4%)	6 (1%)	14	50
1	Bn	522/524 (100%)	510 (98%)	7 (1%)	5 (1%)	15	52
2	Af	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Al	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Ar	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Ax	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Bd	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Bj	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
2	Bp	95/97 (98%)	82 (86%)	6 (6%)	7 (7%)	1	10
All	All	7973/8015 (100%)	7599 (95%)	248 (3%)	126 (2%)	13	42

5 of 126 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Ac	171	LYS
2	Af	4	ARG
2	Af	95	VAL
1	Ai	171	LYS
2	Al	4	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Ac	404/404 (100%)	372 (92%)	32 (8%)	12	41
1	Ad	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Ai	404/404 (100%)	373 (92%)	31 (8%)	13	42
1	Aj	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Ao	404/404 (100%)	373 (92%)	31 (8%)	13	42
1	Ap	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Au	404/404 (100%)	373 (92%)	31 (8%)	13	42
1	Av	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Ba	404/404 (100%)	372 (92%)	32 (8%)	12	41
1	Bb	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Bg	404/404 (100%)	373 (92%)	31 (8%)	13	42
1	Bh	404/404 (100%)	383 (95%)	21 (5%)	23	55
1	Bm	404/404 (100%)	372 (92%)	32 (8%)	12	41
1	Bn	404/404 (100%)	383 (95%)	21 (5%)	23	55
2	Af	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Al	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Ar	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Ax	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Bd	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Bj	80/80 (100%)	55 (69%)	25 (31%)	0	2
2	Bp	80/80 (100%)	55 (69%)	25 (31%)	0	2
All	All	6216/6216 (100%)	5674 (91%)	542 (9%)	14	37

5 of 542 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	Bj	94	ILE
1	Bm	199	TYR
2	Bj	81	GLU
2	Bp	4	ARG
1	Ap	281	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 70

such sidechains are listed below:

Mol	Chain	Res	Type
1	Bh	82	ASN
1	Bh	351	GLN
1	Bm	479	ASN
1	Ao	467	ASN
1	Ao	437	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 28 ligands modelled in this entry, 14 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	Ap	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Ba	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.40	2 (6%)
3	ADP	Bm	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.39	2 (6%)
3	ADP	Bb	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Ad	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Bh	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Ao	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.40	2 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	Au	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.40	2 (6%)
3	ADP	Ai	601	4	24,29,29	3.03	6 (25%)	29,45,45	1.39	2 (6%)
3	ADP	Av	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Ac	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.40	2 (6%)
3	ADP	Bg	601	4	24,29,29	3.02	6 (25%)	29,45,45	1.39	2 (6%)
3	ADP	Aj	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)
3	ADP	Bn	601	4	24,29,29	3.06	6 (25%)	29,45,45	1.50	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	Ap	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Ba	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Bm	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Bb	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Ad	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Bh	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Ao	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Au	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Av	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Ai	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Ac	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Bg	601	4	-	5/12/32/32	0/3/3/3
3	ADP	Aj	601	4	1/1/6/6	6/12/32/32	0/3/3/3
3	ADP	Bn	601	4	1/1/6/6	6/12/32/32	0/3/3/3

The worst 5 of 84 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	Bh	601	ADP	C3'-C4'	-7.99	1.32	1.53
3	Aj	601	ADP	C3'-C4'	-7.97	1.32	1.53
3	Av	601	ADP	C3'-C4'	-7.97	1.32	1.53
3	Ad	601	ADP	C3'-C4'	-7.97	1.32	1.53
3	Ai	601	ADP	O4'-C4'	7.97	1.62	1.45

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Ao	601	ADP	N3-C2-N1	-5.14	120.65	128.68
3	Au	601	ADP	N3-C2-N1	-5.13	120.65	128.68
3	Ac	601	ADP	N3-C2-N1	-5.12	120.67	128.68
3	Ba	601	ADP	N3-C2-N1	-5.12	120.68	128.68
3	Ai	601	ADP	N3-C2-N1	-5.11	120.69	128.68

5 of 7 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	Ad	601	ADP	C1'
3	Aj	601	ADP	C1'
3	Ap	601	ADP	C1'
3	Av	601	ADP	C1'
3	Bb	601	ADP	C1'


5 of 77 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
3	Ad	601	ADP	C5'-O5'-PA-O1A
3	Ad	601	ADP	C5'-O5'-PA-O2A
3	Ad	601	ADP	C5'-O5'-PA-O3A
3	Ad	601	ADP	O4'-C4'-C5'-O5'
3	Ad	601	ADP	C3'-C4'-C5'-O5'



There are no ring outliers.



No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-13308. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192



Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 215



Y Index: 159




Z Index: 175


6.3.2 Raw map



X Index: 222



Y Index: 159




Z Index: 175


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 1.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_13308_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 334 nm³; this corresponds to an approximate mass of 302 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.292 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.292 Å⁻¹

8.2 Resolution estimates


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.43	-	-
Author-provided FSC curve	3.43	4.00	3.45
Unmasked-calculated*	4.21	7.21	4.27

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.21 differs from the reported value 3.43 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-13308 and PDB model 7PBX. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 1.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (1.5).













































9.4 Atom inclusion [i](#)



At the recommended contour level, 76% of all backbone atoms, 61% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (1.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6120	 0.3810
Ac	 0.5040	 0.3360
Ad	 0.7350	 0.4450
Af	 0.5410	 0.2880
Ai	 0.5030	 0.3340
Aj	 0.7330	 0.4460
Al	 0.5510	 0.2900
Ao	 0.5010	 0.3350
Ap	 0.7340	 0.4460
Ar	 0.5540	 0.2910
Au	 0.5020	 0.3340
Av	 0.7350	 0.4450
Ax	 0.5470	 0.2890
Ba	 0.5030	 0.3350
Bb	 0.7330	 0.4450
Bd	 0.5510	 0.2900
Bg	 0.5030	 0.3330
Bh	 0.7340	 0.4460
Bj	 0.5470	 0.2880
Bm	 0.5020	 0.3340
Bn	 0.7330	 0.4450
Bp	 0.5510	 0.2900

