



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 30, 2022 – 03:18 pm GMT

PDB ID : 7QXO  
Title : The structure of T. forsythia NanH  
Authors : Rafferty, J.; Stafford, G.; Satur, M.  
Deposited on : 2022-01-26  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.3  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.3


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	519	 4% 88% 11% .
1	B	519	 10% 92% 8%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 16620 atoms, of which 8091 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called BNR/Asp-box repeat protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	519	8027	2504	4005	724	774	20	126	0	0
1	B	519	8038	2507	4010	725	776	20	127	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	237	ALA	ASP	engineered mutation	UNP G8UIQ1
B	237	ALA	ASP	engineered mutation	UNP G8UIQ1

- Molecule 2 is (2 {R},4 {S},5 {S},6 {S})-5-acetamido-4-oxidanyl-6-[(1 {R},2 {R})-1,2,3-tris(oxidanyl)propyl]-2-[[[(2 {R},3 {R},4 {S},5 {R},6 {S})-3,4,5-tris(oxidanyl)-6-[(2 {R},3 {S},4 {R},5 {R})-1,2,4,5-tetrakis(oxidanyl)-6-oxidanylidene-hexan-3-yl]oxy-oxan-2-yl]methoxy]oxane-2-carboxylic acid (three-letter code: GHD) (formula: C<sub>23</sub>H<sub>39</sub>NO<sub>19</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	11	0
			81	23	38	1	19		
2	B	1	Total	C	H	N	O	11	0
			81	23	38	1	19		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	211	Total	O	0	0
			211	211		
3	B	182	Total	O	0	0
			182	182		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: BNR/Asp-box repeat protein



- Molecule 1: BNR/Asp-box repeat protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.87Å 79.87Å 349.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.64 – 1.90 72.64 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (72.64-1.90) 99.9 (72.64-1.90)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.252 , 0.290 0.255 , 0.292	Depositor DCC
$R_{free}$ test set	4483 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	16620	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GHD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/4100	0.80	0/5568
1	B	0.67	0/4106	0.80	0/5576
All	All	0.66	0/8206	0.80	0/11144

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4022	4005	3990	41	0
1	B	4028	4010	3994	30	0
2	A	43	38	0	0	0
2	B	43	38	0	0	0
3	A	211	0	0	2	0
3	B	182	0	0	1	0
All	All	8529	8091	7984	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:322:GLN:HB3	1:B:324:MET:CE	1.98	0.92
1:A:263:MET:CE	1:A:324:MET:HB2	2.07	0.84
1:B:322:GLN:CB	1:B:324:MET:HE2	2.12	0.80
1:B:200:VAL:O	1:B:201:ARG:HD2	1.83	0.79
1:A:263:MET:HE1	1:A:324:MET:HB2	1.64	0.78
1:A:263:MET:HE3	1:A:324:MET:CB	2.14	0.78
1:B:322:GLN:CB	1:B:324:MET:CE	2.63	0.76
1:B:322:GLN:HB2	1:B:324:MET:HE2	1.65	0.76
1:A:263:MET:HE3	1:A:324:MET:HB3	1.69	0.75
1:A:160:MET:HG3	1:A:196:MET:HE1	1.69	0.75
1:B:322:GLN:HB3	1:B:324:MET:HE1	1.68	0.73
1:A:263:MET:CE	1:A:324:MET:CB	2.67	0.71
1:B:212:ARG:NH2	3:B:701:HOH:O	2.23	0.71
1:A:62:ALA:HB2	1:A:147:MET:HE3	1.76	0.67
1:A:160:MET:HG3	1:A:196:MET:CE	2.25	0.66
1:A:34:ASP:OD2	1:A:61:ASP:HB2	1.99	0.63
1:A:276:ASN:OD1	1:A:300:HIS:HD2	1.85	0.60
1:A:300:HIS:HE1	3:A:858:HOH:O	1.84	0.59
1:B:289:THR:OG1	1:B:291:THR:HG22	2.02	0.59
1:A:68:LEU:HD21	1:A:177:VAL:CG2	2.37	0.55
1:A:488:HIS:HD2	1:A:489:HIS:ND1	2.05	0.55
1:B:88:ARG:NH1	1:B:131:GLU:OE2	2.40	0.54
1:A:71:LEU:HD11	1:A:174:VAL:HG11	1.90	0.53
1:A:57:ILE:HD11	1:A:154:PHE:CD1	2.45	0.52
1:A:302:MET:CE	1:A:307:ALA:HA	2.40	0.52
1:A:213:ILE:HG21	1:A:280:ASP:HA	1.91	0.52
3:A:702:HOH:O	1:B:302:MET:HA	2.09	0.52
1:B:213:ILE:HG21	1:B:280:ASP:HA	1.91	0.50
1:B:71:LEU:C	1:B:71:LEU:HD12	2.31	0.50
1:B:286:ASP:HB3	1:B:291:THR:HG22	1.94	0.50
1:B:345:VAL:HB	1:B:397:ILE:HD12	1.93	0.50
1:A:217:VAL:HB	1:A:285:VAL:HG23	1.93	0.50
1:B:283:ILE:O	1:B:359:ARG:HA	2.12	0.49
1:A:283:ILE:O	1:A:359:ARG:HA	2.13	0.49
1:A:345:VAL:HB	1:A:397:ILE:HD12	1.94	0.49
1:B:302:MET:HE1	1:B:307:ALA:HB2	1.95	0.48
1:A:54:LEU:HD22	1:A:158:VAL:CG1	2.44	0.48
1:A:143:THR:CG2	1:A:145:GLN:H	2.27	0.48
1:A:263:MET:CE	1:A:324:MET:HB3	2.40	0.48
1:A:341:ILE:CG2	1:A:341:ILE:O	2.61	0.47
1:A:62:ALA:HB2	1:A:147:MET:CE	2.44	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:286:ASP:OD1	1:B:391:ARG:NH1	2.48	0.47
1:A:52:ASN:O	1:A:157:SER:HA	2.15	0.47
1:B:371:PRO:HG3	1:B:410:VAL:HG12	1.98	0.46
1:B:52:ASN:O	1:B:157:SER:HA	2.15	0.46
1:B:106:PRO:HD3	1:B:153:TYR:CD2	2.51	0.46
1:A:36:VAL:HG22	1:A:59:ILE:HD13	1.98	0.46
1:B:284:LEU:HD12	1:B:284:LEU:C	2.36	0.45
1:A:106:PRO:HD3	1:A:153:TYR:CD2	2.51	0.45
1:A:506:HIS:HA	1:A:550:LEU:O	2.16	0.45
1:B:302:MET:HE1	1:B:307:ALA:HA	2.00	0.44
1:A:69:ASN:HB2	1:A:178:VAL:CG2	2.47	0.44
1:A:302:MET:HB3	1:A:302:MET:HE2	1.65	0.44
1:B:506:HIS:HA	1:B:550:LEU:O	2.17	0.44
1:B:71:LEU:CD1	1:B:73:ILE:HD12	2.48	0.44
1:A:36:VAL:HG13	1:A:59:ILE:HD13	2.00	0.44
1:A:284:LEU:HD12	1:A:284:LEU:C	2.38	0.43
1:A:203:ALA:HB2	1:A:539:HIS:CE1	2.53	0.43
1:A:470:ILE:HG23	1:A:471:ILE:HD13	2.01	0.43
1:A:496:LEU:N	1:A:496:LEU:HD22	2.33	0.43
1:A:378:LEU:HD12	1:A:378:LEU:N	2.34	0.42
1:B:302:MET:HE1	1:B:307:ALA:CB	2.49	0.42
1:A:302:MET:HE3	1:A:307:ALA:HA	2.01	0.42
1:B:203:ALA:HB2	1:B:539:HIS:CE1	2.54	0.42
1:B:215:GLY:O	1:B:226:GLY:HA2	2.19	0.42
1:B:302:MET:CE	1:B:307:ALA:HA	2.49	0.42
1:B:201:ARG:NH2	1:B:206:ASP:OD2	2.53	0.42
1:A:215:GLY:O	1:A:226:GLY:HA2	2.20	0.42
1:A:72:THR:OG1	1:A:176:GLU:HB3	2.20	0.41
1:A:59:ILE:HD12	1:A:179:ILE:HD11	2.02	0.41
1:B:324:MET:HE3	1:B:340:ASN:ND2	2.35	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	517/519 (100%)	494 (96%)	21 (4%)	2 (0%)	34	24
1	B	518/519 (100%)	495 (96%)	21 (4%)	2 (0%)	34	24
All	All	1035/1038 (100%)	989 (96%)	42 (4%)	4 (0%)	34	24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	234	ASN
1	B	234	ASN
1	A	456	ILE
1	B	456	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	445/445 (100%)	438 (98%)	7 (2%)	62	60
1	B	446/445 (100%)	440 (99%)	6 (1%)	69	68
All	All	891/890 (100%)	878 (98%)	13 (2%)	65	62

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	ASP
1	A	68	LEU
1	A	142	LYS
1	A	178	VAL
1	A	240	GLU
1	A	352	PHE
1	A	471	ILE
1	B	134	THR
1	B	168	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	240	GLU
1	B	352	PHE
1	B	391	ARG
1	B	552	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	300	HIS
1	A	488	HIS
1	A	506	HIS
1	B	42	GLN
1	B	163	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GHD	B	601	-	42,44,44	0.55	1 (2%)	58,64,64	1.38	9 (15%)
2	GHD	A	601	-	42,44,44	0.79	1 (2%)	58,64,64	1.65	10 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GHD	B	601	-	-	20/44/84/84	0/2/2/2
2	GHD	A	601	-	-	25/44/84/84	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	GHD	O17-C21	-2.67	1.20	1.30
2	B	601	GHD	O17-C21	-2.13	1.22	1.30

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GHD	C6-C5-C4	5.23	122.94	113.03
2	B	601	GHD	C6-C5-C4	4.84	122.22	113.03
2	A	601	GHD	C22-C23-C3	4.79	117.34	109.98
2	A	601	GHD	O2-C5-C4	-4.58	99.61	109.50
2	B	601	GHD	C4-C3-N1	3.83	117.27	110.91
2	A	601	GHD	C8-C22-C23	3.48	117.71	110.66
2	A	601	GHD	C4-C3-N1	3.33	116.45	110.91
2	A	601	GHD	C3-N1-C2	2.87	130.17	123.18
2	B	601	GHD	O2-C5-C4	-2.87	103.29	109.50
2	A	601	GHD	C8-O5-C4	-2.79	107.73	114.20
2	B	601	GHD	O5-C4-C3	-2.68	107.16	109.78
2	B	601	GHD	C8-O5-C4	-2.67	108.00	114.20
2	B	601	GHD	C22-C8-C21	2.47	116.88	111.19
2	B	601	GHD	C8-C22-C23	2.41	115.54	110.66
2	B	601	GHD	C22-C23-C3	2.35	113.59	109.98
2	B	601	GHD	C3-N1-C2	2.22	128.58	123.18
2	A	601	GHD	O8-C12-C13	2.21	111.03	107.98
2	A	601	GHD	C19-C20-C10	2.19	114.14	110.24
2	A	601	GHD	O6-C8-O5	2.04	115.74	110.07

There are no chirality outliers.

All (45) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GHD	O1-C2-N1-C3
2	A	601	GHD	C4-C3-N1-C2
2	A	601	GHD	C3-C4-C5-C6
2	A	601	GHD	C3-C4-C5-O2
2	A	601	GHD	O5-C4-C5-C6
2	A	601	GHD	O5-C4-C5-O2
2	A	601	GHD	O17-C21-C8-O6
2	A	601	GHD	C21-C8-O6-C9
2	A	601	GHD	C22-C8-O6-C9
2	A	601	GHD	O5-C8-O6-C9
2	A	601	GHD	O8-C12-C13-O9
2	A	601	GHD	O8-C12-C13-C14
2	A	601	GHD	O9-C13-C14-O10
2	A	601	GHD	C13-C12-C15-O11
2	A	601	GHD	O8-C12-C15-O11
2	A	601	GHD	C13-C12-C15-C16
2	A	601	GHD	O8-C12-C15-C16
2	A	601	GHD	C13-C12-O8-C11
2	B	601	GHD	C4-C3-N1-C2
2	B	601	GHD	C3-C4-C5-C6
2	B	601	GHD	C3-C4-C5-O2
2	B	601	GHD	O5-C4-C5-C6
2	B	601	GHD	O5-C4-C5-O2
2	B	601	GHD	C5-C6-C7-O4
2	B	601	GHD	O3-C6-C7-O4
2	B	601	GHD	O17-C21-C8-O6
2	B	601	GHD	C21-C8-O6-C9
2	B	601	GHD	C22-C8-O6-C9
2	B	601	GHD	O5-C8-O6-C9
2	B	601	GHD	O8-C12-C15-O11
2	B	601	GHD	O8-C12-C15-C16
2	B	601	GHD	C10-C9-O6-C8
2	B	601	GHD	O7-C10-C9-O6
2	A	601	GHD	C12-C13-C14-O10
2	A	601	GHD	C1-C2-N1-C3
2	B	601	GHD	O1-C2-N1-C3
2	B	601	GHD	C1-C2-N1-C3
2	B	601	GHD	C20-C10-C9-O6
2	A	601	GHD	C10-C9-O6-C8
2	B	601	GHD	O18-C21-C8-O6

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
2	A	601	GHD	O3-C6-C7-O4
2	A	601	GHD	C5-C6-C7-O4
2	B	601	GHD	C15-C12-C13-O9
2	A	601	GHD	O7-C11-O8-C12
2	A	601	GHD	C18-C11-O8-C12

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	519/519 (100%)	0.57	19 (3%) 41 44	26, 43, 69, 92	0
1	B	519/519 (100%)	0.84	50 (9%) 8 9	30, 47, 62, 79	0
All	All	1038/1038 (100%)	0.70	69 (6%) 18 20	26, 45, 64, 92	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	34	ASP	4.9
1	A	101	ARG	4.9
1	A	148	VAL	4.1
1	B	274	GLY	3.8
1	A	135	VAL	3.8
1	B	167	LEU	3.4
1	A	34	ASP	3.3
1	A	132	VAL	3.2
1	B	288	ARG	3.1
1	B	261	ILE	3.0
1	B	137	ASN	3.0
1	B	366	GLY	2.9
1	B	353	LEU	2.9
1	B	471	ILE	2.9
1	B	369	VAL	2.9
1	B	271	LEU	2.9
1	A	413	VAL	2.8
1	B	246	LEU	2.8
1	A	125	TYR	2.8
1	B	548	LYS	2.7
1	B	74	ARG	2.7
1	B	170	LEU	2.7
1	B	225	LEU	2.6
1	B	401	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	266	GLY	2.5
1	B	552	ARG	2.5
1	B	293	TRP	2.5
1	B	395	TRP	2.5
1	B	389	LYS	2.5
1	A	36	VAL	2.5
1	B	352	PHE	2.5
1	A	57	ILE	2.5
1	B	296	ALA	2.5
1	B	307	ALA	2.4
1	B	503	LEU	2.4
1	B	547	ILE	2.4
1	A	96	ALA	2.4
1	B	283	ILE	2.4
1	B	524	ILE	2.3
1	B	420	LEU	2.3
1	B	238	LEU	2.3
1	B	442	TRP	2.3
1	B	177	VAL	2.3
1	B	432	VAL	2.3
1	B	470	ILE	2.3
1	B	89	LEU	2.3
1	B	73	ILE	2.2
1	A	139	LEU	2.2
1	A	73	ILE	2.2
1	A	89	LEU	2.2
1	B	101	ARG	2.2
1	B	368	LEU	2.1
1	B	279	GLY	2.1
1	A	75	PHE	2.1
1	A	87	VAL	2.1
1	B	138	THR	2.1
1	B	387	TYR	2.1
1	B	321	ALA	2.1
1	B	200	VAL	2.1
1	B	362	THR	2.1
1	B	242	ILE	2.1
1	B	354	LEU	2.1
1	A	154	PHE	2.1
1	B	292	VAL	2.1
1	A	136	ALA	2.0
1	B	255	THR	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	335	TRP	2.0
1	A	134	THR	2.0
1	B	291	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GHD	B	601	43/43	0.83	0.20	30,51,92,97	11
2	GHD	A	601	43/43	0.89	0.17	27,33,68,72	11




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around GHD B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around GHD A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.