



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 11, 2024 – 07:34 PM EDT

PDB ID : 7SHI
Title : Crystal Structure of Cytochrome P450 AmphL from *Streptomyces nodosus* and the Structural Basis for Substrate Selectivity in Macrolide Metabolizing P450s
Authors : Amaya, J.A.; Poulos, T.L.
Deposited on : 2021-10-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	 2% 83% 14% ..
1	B	396	 2% 85% 12% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6643 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AmphL.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	391	Total 3017	C 1903	N 527	O 573	S 14	0	3	0
1	B	391	Total 3017	C 1903	N 527	O 573	S 14	0	3	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 3 is Amphotericin B (three-letter code: 4UH) (formula: $C_{47}H_{73}NO_{17}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	N	O	0	0
			65	47	1	17		
3	B	1	Total	C	N	O	0	0
			65	47	1	17		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	180	Total	O	0	0
			180	180		
5	B	198	Total	O	0	0
			198	198		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: AmphL



- Molecule 1: AmphL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.46Å 68.15Å 147.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	36.88 – 2.00 36.88 – 2.00	Depositor EDS
% Data completeness (in resolution range)	95.0 (36.88-2.00) 95.0 (36.88-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.18_3845: ???)	Depositor
R, R_{free}	0.189 , 0.248 0.189 , 0.248	Depositor DCC
R_{free} test set	1999 reflections (4.14%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.371	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6643	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.32 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4026e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4UH, HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/3071	0.56	0/4175
1	B	0.38	0/3071	0.57	0/4175
All	All	0.38	0/6142	0.57	0/8350

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3017	0	3045	37	0
1	B	3017	0	3045	29	0
2	A	43	0	30	2	0
2	B	43	0	30	1	0
3	A	65	0	0	0	0
3	B	65	0	0	0	0
4	A	10	0	0	0	0
4	B	5	0	0	0	0
5	A	180	0	0	2	0
5	B	198	0	0	2	1
All	All	6643	0	6150	63	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:ARG:HH12	1:B:93:ARG:HH12	1.25	0.82
1:A:175:THR:HG22	1:A:177:ASN:H	1.46	0.81
1:A:17:VAL:HG12	1:A:18:LEU:HG	1.72	0.70
1:A:93:ARG:NH1	1:B:93:ARG:HH12	1.91	0.69
1:A:58:HIS:ND1	1:B:60:GLU:OE1	2.19	0.68
1:A:116:MET:HG2	1:A:353:MET:SD	2.38	0.62
1:B:263:LYS:HD3	1:B:266:ARG:HE	1.64	0.62
1:B:244:GLY:HA2	1:B:272:ILE:HD12	1.83	0.61
1:B:358:ALA:O	1:B:362:LEU:HB2	2.01	0.61
1:B:190:PHE:O	1:B:194:THR:HG23	2.03	0.59
1:A:366:LEU:HD22	1:A:396:TRP:CD1	2.38	0.59
1:A:93:ARG:NH2	5:A:415:HOH:O	2.36	0.58
1:B:170:LEU:HD13	1:B:234:LEU:HD22	1.85	0.58
1:A:112:VAL:HG21	1:A:349:PRO:HB2	1.86	0.57
1:B:149:LEU:HD22	1:B:167:LEU:HD13	1.88	0.55
1:B:17:VAL:HG12	1:B:18:LEU:HG	1.90	0.54
1:B:277:LYS:HG3	1:B:311:THR:HG23	1.91	0.53
1:A:229:LEU:CD2	2:A:397:HEM:HAC	2.40	0.52
1:A:149:LEU:HD22	1:A:167:LEU:HD13	1.91	0.52
1:A:263:LYS:HD3	1:A:266:ARG:HH21	1.75	0.51
1:A:374:SER:HB3	1:A:377:GLU:HG3	1.91	0.51
1:B:221:ARG:HB3	1:B:221:ARG:HH21	1.76	0.51
1:A:200:LYS:NZ	1:A:207:ASP:OD2	2.35	0.50
1:B:128:ASN:OD1	1:B:365:ARG:NH2	2.37	0.49
1:A:84:LEU:HD21	1:A:285:LEU:HD13	1.94	0.48
1:B:372:ALA:HB3	1:B:393:PRO:HB2	1.96	0.48
1:B:194:THR:HG22	1:B:223:GLY:HA3	1.95	0.48
1:B:263:LYS:HE3	1:B:266:ARG:HH21	1.79	0.47
1:A:358:ALA:O	1:A:362:LEU:HB2	2.14	0.47
1:A:7:PRO:HG2	1:A:43:GLU:HG3	1.96	0.47
1:B:112:VAL:HG21	1:B:349:PRO:HB2	1.96	0.46
1:A:373:SER:HB2	1:A:377:GLU:OE1	2.15	0.46
1:B:229:LEU:HD22	2:B:397:HEM:HAC	1.98	0.46
1:A:30[B]:ARG:NH1	1:A:30[B]:ARG:HA	2.32	0.45
1:B:268:GLY:O	1:B:272:ILE:HG12	2.17	0.45
1:A:19:ARG:HD3	1:A:19:ARG:O	2.17	0.45
1:B:381:THR:HG23	1:B:388:GLY:O	2.16	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:GLU:O	1:A:60:GLU:HG2	2.18	0.44
1:A:190:PHE:CE2	1:A:224:PRO:HB3	2.53	0.44
1:A:245:VAL:O	1:A:249:THR:HG23	2.17	0.44
1:B:119:VAL:HG11	1:B:353:MET:HE2	2.00	0.44
1:B:334:GLU:OE1	1:B:334:GLU:N	2.38	0.44
1:A:17:VAL:HG21	1:A:389:LEU:HD21	2.01	0.43
1:A:221:ARG:CZ	1:A:221:ARG:HB2	2.48	0.43
1:A:229:LEU:HD22	2:A:397:HEM:HAC	2.00	0.43
1:B:66:HIS:O	1:B:87:ASP:HA	2.18	0.43
1:A:19:ARG:NH2	5:A:414:HOH:O	2.48	0.43
1:A:194:THR:HG23	1:A:223:GLY:HA3	2.01	0.42
1:A:193:LEU:HD23	1:A:196:LEU:HD23	2.01	0.42
1:A:220:GLU:H	1:A:220:GLU:CD	2.24	0.41
1:B:111:ARG:NH2	5:B:415:HOH:O	2.53	0.41
1:A:83:LEU:HD23	1:A:83:LEU:HA	1.77	0.41
1:A:13:ALA:HB2	1:A:382:SER:HB3	2.02	0.41
1:B:119:VAL:HB	1:B:353:MET:HE1	2.02	0.41
1:A:6:PRO:HA	1:A:7:PRO:HD3	1.83	0.41
1:B:35:LYS:HE3	1:B:35:LYS:HB2	1.85	0.41
1:A:167:LEU:HD12	1:A:167:LEU:HA	1.93	0.41
1:B:30[B]:ARG:NH1	1:B:30[B]:ARG:HA	2.35	0.40
1:A:80:LEU:HD11	1:A:173:MET:HE3	2.02	0.40
1:B:211:ARG:NH2	5:B:426:HOH:O	2.55	0.40
1:A:184:ARG:HE	1:A:184:ARG:HB2	1.58	0.40
1:A:170:LEU:HD13	1:A:234:LEU:HD22	2.01	0.40
1:B:366:LEU:HD22	1:B:396:TRP:CD1	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:465:HOH:O	5:B:568:HOH:O[3_754]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/396 (99%)	381 (97%)	9 (2%)	2 (0%)	29	23
1	B	392/396 (99%)	382 (97%)	9 (2%)	1 (0%)	41	37
All	All	784/792 (99%)	763 (97%)	18 (2%)	3 (0%)	34	30

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	386	ASN
1	A	13	ALA
1	B	386	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/323 (99%)	302 (94%)	19 (6%)	19	15
1	B	321/323 (99%)	305 (95%)	16 (5%)	24	20
All	All	642/646 (99%)	607 (94%)	35 (6%)	22	17

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	GLU
1	A	19	ARG
1	A	30[A]	ARG
1	A	30[B]	ARG
1	A	53	LEU
1	A	80	LEU
1	A	139	LEU
1	A	140	ARG
1	A	167	LEU
1	A	215	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	221	ARG
1	A	259	LEU
1	A	298	VAL
1	A	340	HIS
1	A	362	LEU
1	A	371	LEU
1	A	380	VAL
1	A	381	THR
1	A	393	PRO
1	B	19	ARG
1	B	84	LEU
1	B	99	LEU
1	B	110	ARG
1	B	140	ARG
1	B	167	LEU
1	B	172	GLU
1	B	215	THR
1	B	259	LEU
1	B	298	VAL
1	B	334	GLU
1	B	340	HIS
1	B	353	MET
1	B	362	LEU
1	B	371	LEU
1	B	386	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	397	1	41,50,50	1.53	6 (14%)	45,82,82	1.59	10 (22%)
3	4UH	A	398	-	65,67,67	1.54	10 (15%)	76,92,92	2.82	14 (18%)
2	HEM	A	397	1	41,50,50	1.39	4 (9%)	45,82,82	1.45	6 (13%)
3	4UH	B	398	-	65,67,67	1.48	11 (16%)	76,92,92	2.76	19 (25%)
4	SO4	A	400	-	4,4,4	0.11	0	6,6,6	0.21	0
4	SO4	A	399	-	4,4,4	0.15	0	6,6,6	0.11	0
4	SO4	B	399	-	4,4,4	0.15	0	6,6,6	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	4UH	B	398	-	-	8/71/109/109	0/2/3/3
2	HEM	B	397	1	-	4/12/54/54	-
3	4UH	A	398	-	-	8/71/109/109	0/2/3/3
2	HEM	A	397	1	-	4/12/54/54	-

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	397	HEM	C3C-C2C	-4.38	1.34	1.40
3	B	398	4UH	O15-C49	-3.93	1.40	1.46
3	A	398	4UH	O15-C49	-3.78	1.40	1.46
3	A	398	4UH	C65-C63	3.63	1.54	1.44
2	A	397	HEM	C3C-C2C	-3.62	1.35	1.40
3	A	398	4UH	C61-C59	3.48	1.54	1.44
3	B	398	4UH	C65-C63	3.47	1.54	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	398	4UH	C61-C59	3.31	1.53	1.44
3	A	398	4UH	C64-C62	3.27	1.53	1.44
3	B	398	4UH	C57-C51	3.26	1.53	1.44
2	B	397	HEM	C3C-CAC	3.26	1.54	1.47
2	A	397	HEM	C3C-CAC	3.25	1.54	1.47
3	A	398	4UH	C60-C58	3.19	1.53	1.44
3	A	398	4UH	C45-C41	3.16	1.53	1.44
3	B	398	4UH	C64-C62	3.05	1.53	1.44
3	A	398	4UH	C57-C51	3.04	1.53	1.44
3	B	398	4UH	C45-C41	3.04	1.53	1.44
2	B	397	HEM	CAB-C3B	2.91	1.55	1.47
3	B	398	4UH	C60-C58	2.89	1.52	1.44
2	A	397	HEM	CAB-C3B	2.74	1.54	1.47
3	A	398	4UH	O15-C52	2.62	1.41	1.34
3	B	398	4UH	O15-C52	2.52	1.41	1.34
3	A	398	4UH	O04-C27	-2.44	1.41	1.44
2	B	397	HEM	CAA-C2A	2.39	1.55	1.52
3	B	398	4UH	O04-C27	-2.35	1.41	1.44
2	B	397	HEM	FE-NB	2.24	2.07	1.96
3	B	398	4UH	C24-C26	2.21	1.56	1.53
3	B	398	4UH	C46-C52	2.11	1.54	1.50
2	A	397	HEM	CMB-C2B	2.08	1.55	1.50
2	B	397	HEM	C1B-NB	-2.02	1.36	1.40
3	A	398	4UH	C21-C28	2.01	1.56	1.51

All (49) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	398	4UH	O02-C19-C24	-14.24	87.54	111.62
3	A	398	4UH	O02-C19-C24	-13.55	88.71	111.62
3	A	398	4UH	O01-C19-C23	11.87	123.12	108.92
3	B	398	4UH	O01-C19-C23	11.46	122.63	108.92
3	A	398	4UH	O02-C19-C23	-10.13	86.28	110.67
3	B	398	4UH	O02-C19-C23	-9.59	87.56	110.67
3	A	398	4UH	O15-C52-C46	4.08	118.96	111.46
2	B	397	HEM	CBA-CAA-C2A	-3.47	106.70	112.62
3	B	398	4UH	C39-C35-C34	-3.39	106.82	113.07
3	B	398	4UH	C64-C62-C60	-3.34	117.39	124.81
3	A	398	4UH	C53-C47-C49	-3.17	108.08	112.18
2	B	397	HEM	C3B-C2B-C1B	3.10	108.79	106.49
3	B	398	4UH	O15-C52-C46	3.01	117.00	111.46
2	B	397	HEM	C4B-CHC-C1C	2.98	126.49	122.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	398	4UH	C63-C65-C64	-2.93	118.29	124.81
2	A	397	HEM	CMA-C3A-C4A	-2.86	124.06	128.46
2	A	397	HEM	C4B-CHC-C1C	2.83	126.29	122.56
2	B	397	HEM	C1B-NB-C4B	2.82	107.98	105.07
3	B	398	4UH	C30-O04-C27	-2.81	110.99	115.63
3	A	398	4UH	C39-C35-C34	-2.80	107.89	113.07
2	A	397	HEM	C1B-NB-C4B	2.70	107.86	105.07
2	B	397	HEM	C4D-ND-C1D	2.62	107.78	105.07
3	A	398	4UH	O07-C28-O08	-2.59	118.22	124.09
2	A	397	HEM	CMC-C2C-C3C	2.58	129.50	124.68
2	B	397	HEM	CHC-C4B-NB	2.55	127.20	124.43
3	B	398	4UH	C57-C51-C45	-2.53	119.17	124.81
3	A	398	4UH	C62-C64-C65	-2.47	119.32	124.81
3	A	398	4UH	C64-C62-C60	-2.45	119.36	124.81
3	B	398	4UH	O07-C28-O08	-2.41	118.63	124.09
3	A	398	4UH	O07-C28-C21	2.40	121.13	114.01
3	A	398	4UH	C57-C51-C45	-2.34	119.61	124.81
2	A	397	HEM	CBA-CAA-C2A	-2.29	108.72	112.62
2	B	397	HEM	CHD-C1D-ND	2.26	126.89	124.43
3	B	398	4UH	C44-C43-C42	-2.26	110.75	114.18
3	B	398	4UH	C43-C44-C46	-2.25	108.71	113.19
3	B	398	4UH	O07-C28-C21	2.22	120.59	114.01
3	A	398	4UH	C30-O05-C35	-2.20	109.89	113.67
2	A	397	HEM	O2D-CGD-CBD	2.19	121.07	114.03
3	B	398	4UH	C65-C63-C61	-2.17	119.97	124.81
3	B	398	4UH	C51-C57-C59	-2.16	120.00	124.81
3	A	398	4UH	C59-C61-C63	-2.11	120.11	124.81
2	B	397	HEM	CMA-C3A-C4A	-2.10	125.24	128.46
2	B	397	HEM	C4A-C3A-C2A	2.08	108.44	107.00
3	B	398	4UH	C61-C59-C57	-2.04	120.28	124.81
3	A	398	4UH	C63-C65-C64	-2.03	120.29	124.81
2	B	397	HEM	CMC-C2C-C3C	2.02	128.46	124.68
3	B	398	4UH	C22-C21-C28	-2.02	105.81	109.78
3	B	398	4UH	C58-C60-C62	-2.01	120.34	124.81
3	B	398	4UH	C50-C48-C47	-2.01	110.87	114.96

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	398	4UH	O01-C19-C24-C26
3	A	398	4UH	C25-C27-C37-C41

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	398	4UH	O13-C42-C43-C44
3	B	398	4UH	O01-C19-C24-C26
3	B	398	4UH	C25-C27-C37-C41
3	B	398	4UH	C56-C58-C60-C62
3	B	398	4UH	C48-C50-C56-C58
3	B	398	4UH	O13-C42-C43-C44
3	A	398	4UH	O04-C27-C37-C41
3	A	398	4UH	C49-C47-C48-C50
3	A	398	4UH	C56-C58-C60-C62
2	A	397	HEM	CAA-CBA-CGA-O1A
2	A	397	HEM	CAA-CBA-CGA-O2A
2	B	397	HEM	CAA-CBA-CGA-O1A
3	A	398	4UH	C53-C47-C48-C50
2	B	397	HEM	CAD-CBD-CGD-O2D
2	B	397	HEM	CAA-CBA-CGA-O2A
3	B	398	4UH	O04-C27-C37-C41
2	A	397	HEM	CAD-CBD-CGD-O2D
2	A	397	HEM	CAD-CBD-CGD-O1D
2	B	397	HEM	CAD-CBD-CGD-O1D
3	B	398	4UH	C53-C47-C49-O15
3	B	398	4UH	C47-C49-O15-C52
3	A	398	4UH	C47-C48-C50-C56


There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	397	HEM	1	0
2	A	397	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/396 (98%)	-0.04	6 (1%) 73 72	13, 27, 46, 66	0
1	B	391/396 (98%)	-0.09	9 (2%) 60 59	12, 24, 42, 82	0
All	All	782/792 (98%)	-0.06	15 (1%) 66 65	12, 25, 45, 82	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	14	ALA	11.1
1	B	13	ALA	7.7
1	B	12	ASP	4.1
1	B	177	ASN	3.8
1	B	15	PRO	3.7
1	A	30[A]	ARG	3.4
1	B	11	GLU	3.2
1	A	12	ASP	2.7
1	A	7	PRO	2.6
1	A	29	MET	2.5
1	A	26	GLU	2.4
1	B	26	GLU	2.3
1	A	70	ALA	2.3
1	B	30[A]	ARG	2.3
1	B	71	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	4UH	A	398	65/65	0.91	0.15	17,27,35,41	0
3	4UH	B	398	65/65	0.93	0.15	16,24,34,40	0
4	SO4	A	400	5/5	0.97	0.10	34,39,40,44	5
2	HEM	B	397	43/43	0.99	0.13	8,12,19,23	0
4	SO4	A	399	5/5	0.99	0.08	24,24,28,36	0
2	HEM	A	397	43/43	0.99	0.13	9,13,21,25	0
4	SO4	B	399	5/5	0.99	0.11	24,25,27,27	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM B 397:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.