



## wwPDB EM Validation Summary Report ⓘ

Dec 8, 2022 – 06:27 PM EST

PDB ID : 7V05  
EMDB ID : EMD-26936  
Title : Complex of Plasmodium falciparum circumsporozoite protein with 850 Fab  
Authors : Kucharska, I.; Prieto, K.; Rubinstein, J.L.; Julien, J.P.  
Deposited on : 2022-05-09  
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev43  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	35% 99% .
1	B	226	8% 99% .
1	C	226	. 99% .
1	D	226	99% .
1	E	226	99% .
1	F	226	99% .
1	G	226	99% .
1	H	226	99% .
1	I	226	99% .

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	J	226	 19% 98%
1	K	226	 99%
1	L	226	 99%
1	M	226	 99%
1	N	226	 6% 99%
2	a	213	 36% 100%
2	b	213	 9% 100%
2	c	213	 100%
2	d	213	 100%
2	e	213	 100%
2	f	213	 100%
2	g	213	 100%
2	h	213	 100%
2	i	213	 100%
2	j	213	 20% 100%
2	k	213	 100%
2	l	213	 100%
2	m	213	 100%
2	n	213	 100%
3	X	372	 29% 70%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 47382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 850 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	224	1689	1068	283	332	6	0	0
1	B	224	1689	1068	283	332	6	0	0
1	C	224	1689	1068	283	332	6	0	0
1	D	224	1689	1068	283	332	6	0	0
1	E	224	1689	1068	283	332	6	0	0
1	F	224	1689	1068	283	332	6	0	0
1	G	224	1689	1068	283	332	6	0	0
1	H	224	1689	1068	283	332	6	0	0
1	I	224	1689	1068	283	332	6	0	0
1	J	224	1689	1068	283	332	6	0	0
1	K	224	1689	1068	283	332	6	0	0
1	L	224	1689	1068	283	332	6	0	0
1	M	224	1689	1068	283	332	6	0	0
1	N	224	1689	1068	283	332	6	0	0

- Molecule 2 is a protein called 850 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	a	212	1640	1029	272	334	5	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	b	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	c	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	d	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	e	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	f	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	g	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	h	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	i	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	j	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	k	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	l	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	m	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		
2	n	212	Total	C	N	O	S	0	0
			1640	1029	272	334	5		

- Molecule 3 is a protein called Circumsporozoite protein.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	X	111	Total	C	N	O	0	0
			776	444	166	166		

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
X	74	ALA	VAL	conflict	UNP Q7K740
X	75	ASN	ASP	conflict	UNP Q7K740
X	258	GLN	ASN	conflict	UNP Q7K740
X	261	PRO	-	expression tag	UNP Q7K740
X	262	HIS	-	expression tag	UNP Q7K740
X	263	HIS	-	expression tag	UNP Q7K740

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
X	264	HIS	-	expression tag	UNP Q7K740
X	265	HIS	-	expression tag	UNP Q7K740
X	266	HIS	-	expression tag	UNP Q7K740
X	267	HIS	-	expression tag	UNP Q7K740







• Molecule 1: 850 Fab Heavy Chain




• Molecule 1: 850 Fab Heavy Chain




• Molecule 1: 850 Fab Heavy Chain




• Molecule 1: 850 Fab Heavy Chain



• Molecule 1: 850 Fab Heavy Chain




• Molecule 1: 850 Fab Heavy Chain







- Molecule 1: 850 Fab Heavy Chain




- Molecule 1: 850 Fab Heavy Chain




- Molecule 1: 850 Fab Heavy Chain




- Molecule 2: 850 Fab Light Chain



- Molecule 2: 850 Fab Light Chain



- Molecule 2: 850 Fab Light Chain



- Molecule 2: 850 Fab Light Chain

Chain d:  100%



• Molecule 2: 850 Fab Light Chain

Chain e:  100%



• Molecule 2: 850 Fab Light Chain

Chain f:  100%



• Molecule 2: 850 Fab Light Chain

Chain g:  100%



• Molecule 2: 850 Fab Light Chain

Chain h:  100%




• Molecule 2: 850 Fab Light Chain

Chain i:  100%



• Molecule 2: 850 Fab Light Chain

Chain j:   100%





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	165747	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	45.24	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	2.494	Depositor
Minimum map value	-1.458	Depositor
Average map value	-0.004	Depositor
Map value standard deviation	0.071	Depositor
Recommended contour level	0.2	Depositor
Map size (Å)	350.19998, 350.19998, 350.19998	wwPDB
Map dimensions	340, 340, 340	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.03, 1.03, 1.03	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.28	0/1732	0.49	0/2361
1	B	0.33	0/1732	0.50	0/2361
1	C	0.42	0/1732	0.51	0/2361
1	D	0.54	0/1732	0.53	0/2361
1	E	0.58	0/1732	0.52	0/2361
1	F	0.56	0/1732	0.53	0/2361
1	G	0.59	0/1732	0.53	0/2361
1	H	0.55	0/1732	0.53	0/2361
1	I	0.57	0/1732	0.54	0/2361
1	J	0.33	0/1732	0.50	0/2361
1	K	0.50	0/1732	0.53	0/2361
1	L	0.55	0/1732	0.53	0/2361
1	M	0.54	0/1732	0.53	0/2361
1	N	0.41	0/1732	0.52	0/2361
2	a	0.28	0/1677	0.49	0/2279
2	b	0.34	0/1677	0.50	0/2279
2	c	0.39	0/1677	0.51	0/2279
2	d	0.46	0/1677	0.52	0/2279
2	e	0.50	0/1677	0.52	0/2279
2	f	0.48	0/1677	0.52	0/2279
2	g	0.50	0/1677	0.53	0/2279
2	h	0.48	0/1677	0.53	0/2279
2	i	0.48	0/1677	0.52	0/2279
2	j	0.27	0/1677	0.49	0/2279
2	k	0.38	0/1677	0.51	0/2279
2	l	0.50	0/1677	0.53	0/2279
2	m	0.45	0/1677	0.52	0/2279
2	n	0.31	0/1677	0.50	0/2279
3	X	0.45	0/803	0.53	0/1133
All	All	0.46	0/48529	0.52	0/66093

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/226 (98%)	217 (98%)	5 (2%)	0	100	100
1	B	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	C	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	D	222/226 (98%)	219 (99%)	3 (1%)	0	100	100
1	E	222/226 (98%)	219 (99%)	3 (1%)	0	100	100
1	F	222/226 (98%)	217 (98%)	5 (2%)	0	100	100
1	G	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	H	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	I	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	J	222/226 (98%)	217 (98%)	5 (2%)	0	100	100
1	K	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	L	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	M	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
1	N	222/226 (98%)	218 (98%)	4 (2%)	0	100	100
2	a	210/213 (99%)	204 (97%)	6 (3%)	0	100	100
2	b	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	c	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	d	210/213 (99%)	205 (98%)	5 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	e	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	f	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	g	210/213 (99%)	204 (97%)	6 (3%)	0	100	100
2	h	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	i	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	j	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	k	210/213 (99%)	204 (97%)	6 (3%)	0	100	100
2	l	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	m	210/213 (99%)	205 (98%)	5 (2%)	0	100	100
2	n	210/213 (99%)	204 (97%)	6 (3%)	0	100	100
3	X	109/372 (29%)	89 (82%)	20 (18%)	0	100	100
All	All	6157/6518 (94%)	6006 (98%)	151 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	B	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	C	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	D	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	E	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	F	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	G	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	H	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	I	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	J	189/192 (98%)	187 (99%)	2 (1%)	73	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	K	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	L	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	M	189/192 (98%)	188 (100%)	1 (0%)	88	94
1	N	189/192 (98%)	188 (100%)	1 (0%)	88	94
2	a	187/189 (99%)	187 (100%)	0	100	100
2	b	187/189 (99%)	187 (100%)	0	100	100
2	c	187/189 (99%)	187 (100%)	0	100	100
2	d	187/189 (99%)	187 (100%)	0	100	100
2	e	187/189 (99%)	187 (100%)	0	100	100
2	f	187/189 (99%)	187 (100%)	0	100	100
2	g	187/189 (99%)	187 (100%)	0	100	100
2	h	187/189 (99%)	187 (100%)	0	100	100
2	i	187/189 (99%)	187 (100%)	0	100	100
2	j	187/189 (99%)	187 (100%)	0	100	100
2	k	187/189 (99%)	187 (100%)	0	100	100
2	l	187/189 (99%)	187 (100%)	0	100	100
2	m	187/189 (99%)	187 (100%)	0	100	100
2	n	187/189 (99%)	187 (100%)	0	100	100
3	X	83/314 (26%)	79 (95%)	4 (5%)	25	56
All	All	5347/5648 (95%)	5328 (100%)	19 (0%)	91	95

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	100(C)	ASN
1	M	52	TRP
1	N	52	TRP
1	L	52	TRP
3	X	37	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	m	37	GLN



### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-26936. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 170




Y Index: 170




Z Index: 170


### 6.2.2 Raw map



X Index: 170



Y Index: 170




Z Index: 170


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 174




Y Index: 171




Z Index: 168


### 6.3.2 Raw map



X Index: 173



Y Index: 171




Z Index: 168

The images above show the largest variance slices of the map in three orthogonal directions.


## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.2. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.5 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 525 nm<sup>3</sup>; this corresponds to an approximate mass of 474 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.294 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.294  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.17	6.82	4.27

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.17 differs from the reported value 3.4 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-26936 and PDB model 7V05. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.2 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.2).





























































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8522	 0.3810
A	 0.5322	 0.2360
B	 0.7736	 0.3090
C	 0.8742	 0.3750
D	 0.9091	 0.3960
E	 0.9500	 0.4550
F	 0.9524	 0.4560
G	 0.9404	 0.4500
H	 0.9374	 0.4440
I	 0.9278	 0.4220
J	 0.6755	 0.3050
K	 0.8808	 0.4050
L	 0.9422	 0.4520
M	 0.9163	 0.4220
N	 0.8176	 0.3640
X	 0.9691	 0.4860
a	 0.5229	 0.2510
b	 0.7546	 0.3050
c	 0.8389	 0.3490
d	 0.8885	 0.3830
e	 0.9263	 0.4340
f	 0.9275	 0.4330
g	 0.9244	 0.4230
h	 0.9170	 0.4170
i	 0.9133	 0.4140
j	 0.6357	 0.2310
k	 0.8798	 0.3610
l	 0.9337	 0.4330
m	 0.9040	 0.3890
n	 0.8067	 0.3060

