



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 7, 2024 – 02:54 pm BST

PDB ID : 8RLT
Title : TCR in complex with HLA-E*01:03 bound to HBV envelope 371-379 index peptide
Authors : Pengelly, R.J.; Godinho, L.F.
Deposited on : 2024-01-03
Resolution : 2.25 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	 2% 89% 10%
1	F	276	 2% 87% 11%
2	B	100	 2% 92% 8%
2	G	100	 2% 89% 11%
3	C	9	 78% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	9	 89% 11%
4	D	199	 5% 78% 15% 8%
4	I	199	 8% 79% 10% 10%
5	E	243	 5% 86% 10%
5	J	243	 9% 77% 16% 6%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 13060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, alpha chain E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	276	Total	C	N	O	S	0	1	0
			2253	1408	403	435	7			
1	F	271	Total	C	N	O	S	0	0	0
			2208	1383	394	424	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	G	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
G	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Large envelope protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	Total	C	N	O	0	0	0
			72	52	9	11			
3	H	9	Total	C	N	O	0	0	0
			72	52	9	11			

- Molecule 4 is a protein called T cell receptor alpha variable 12-2,T cell receptor alpha chain MC.7.G5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	184	Total	C	N	O	S	0	0	0
			1421	884	242	285	10			
4	I	179	Total	C	N	O	S	0	0	0
			1385	861	237	277	10			

There are 52 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	MET	-	initiating methionine	UNP A0A075B6T6
D	1	ALA	-	expression tag	UNP A0A075B6T6
D	31	VAL	GLN	variant	UNP A0A075B6T6
D	49	SER	PHE	variant	UNP A0A075B6T6
D	52	LEU	SER	variant	UNP A0A075B6T6
D	55	LEU	ASP	variant	UNP A0A075B6T6
D	92	GLY	-	variant	UNP A0A075B6T6
D	93	ASN	-	variant	UNP A0A075B6T6
D	94	HIS	-	variant	UNP A0A075B6T6
D	95	ASN	-	variant	UNP A0A075B6T6
D	96	THR	-	variant	UNP A0A075B6T6
D	97	GLY	-	variant	UNP A0A075B6T6
D	98	ASN	-	variant	UNP A0A075B6T6
D	99	MET	-	variant	UNP A0A075B6T6
D	100	LEU	-	variant	UNP A0A075B6T6
D	101	THR	-	variant	UNP A0A075B6T6
D	102	PHE	-	variant	UNP A0A075B6T6
D	103	GLY	-	variant	UNP A0A075B6T6
D	104	GLY	-	variant	UNP A0A075B6T6
D	105	GLY	-	variant	UNP A0A075B6T6
D	106	THR	-	variant	UNP A0A075B6T6
D	107	ARG	-	variant	UNP A0A075B6T6
D	108	LEU	-	variant	UNP A0A075B6T6
D	109	MET	-	variant	UNP A0A075B6T6
D	113	HIS	ASN	variant	UNP P0DTU3
D	160	CYS	THR	variant	UNP P0DTU3
I	0	MET	-	initiating methionine	UNP A0A075B6T6
I	1	ALA	-	expression tag	UNP A0A075B6T6
I	31	VAL	GLN	variant	UNP A0A075B6T6
I	49	SER	PHE	variant	UNP A0A075B6T6
I	52	LEU	SER	variant	UNP A0A075B6T6
I	55	LEU	ASP	variant	UNP A0A075B6T6
I	92	GLY	-	variant	UNP A0A075B6T6
I	93	ASN	-	variant	UNP A0A075B6T6
I	94	HIS	-	variant	UNP A0A075B6T6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	95	ASN	-	variant	UNP A0A075B6T6
I	96	THR	-	variant	UNP A0A075B6T6
I	97	GLY	-	variant	UNP A0A075B6T6
I	98	ASN	-	variant	UNP A0A075B6T6
I	99	MET	-	variant	UNP A0A075B6T6
I	100	LEU	-	variant	UNP A0A075B6T6
I	101	THR	-	variant	UNP A0A075B6T6
I	102	PHE	-	variant	UNP A0A075B6T6
I	103	GLY	-	variant	UNP A0A075B6T6
I	104	GLY	-	variant	UNP A0A075B6T6
I	105	GLY	-	variant	UNP A0A075B6T6
I	106	THR	-	variant	UNP A0A075B6T6
I	107	ARG	-	variant	UNP A0A075B6T6
I	108	LEU	-	variant	UNP A0A075B6T6
I	109	MET	-	variant	UNP A0A075B6T6
I	113	HIS	ASN	variant	UNP P0DTU3
I	160	CYS	THR	variant	UNP P0DTU3

- Molecule 5 is a protein called T cell receptor beta variable 6-5,T cell receptor beta chain MC.7.G5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	240	1910	1200	333	368	9	0	0	0
5	J	229	1819	1147	313	350	9	0	0	0

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	0	MET	-	initiating methionine	UNP A0A0K0K1A5
E	29	TYR	HIS	variant	UNP A0A0K0K1A5
E	51	SER	GLY	variant	UNP A0A0K0K1A5
E	54	LEU	ILE	variant	UNP A0A0K0K1A5
E	94	HIS	-	variant	UNP A0A0K0K1A5
E	95	ARG	-	variant	UNP A0A0K0K1A5
E	96	ASN	-	variant	UNP A0A0K0K1A5
E	97	ARG	-	variant	UNP A0A0K0K1A5
E	98	LEU	-	variant	UNP A0A0K0K1A5
E	99	THR	-	variant	UNP A0A0K0K1A5
E	100	GLU	-	variant	UNP A0A0K0K1A5
E	101	ALA	-	variant	UNP A0A0K0K1A5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	102	PHE	-	variant	UNP A0A0K0K1A5
E	105	GLN	PRO	variant	UNP P0DTU4
E	112	VAL	LEU	variant	UNP P0DTU4
E	169	CYS	SER	variant	UNP P0DTU4
E	187	ALA	CYS	variant	UNP P0DTU4
E	201	ASP	ASN	variant	UNP P0DTU4
J	0	MET	-	initiating methionine	UNP A0A0K0K1A5
J	29	TYR	HIS	variant	UNP A0A0K0K1A5
J	51	SER	GLY	variant	UNP A0A0K0K1A5
J	54	LEU	ILE	variant	UNP A0A0K0K1A5
J	94	HIS	-	variant	UNP A0A0K0K1A5
J	95	ARG	-	variant	UNP A0A0K0K1A5
J	96	ASN	-	variant	UNP A0A0K0K1A5
J	97	ARG	-	variant	UNP A0A0K0K1A5
J	98	LEU	-	variant	UNP A0A0K0K1A5
J	99	THR	-	variant	UNP A0A0K0K1A5
J	100	GLU	-	variant	UNP A0A0K0K1A5
J	101	ALA	-	variant	UNP A0A0K0K1A5
J	102	PHE	-	variant	UNP A0A0K0K1A5
J	105	GLN	PRO	variant	UNP P0DTU4
J	112	VAL	LEU	variant	UNP P0DTU4
J	169	CYS	SER	variant	UNP P0DTU4
J	187	ALA	CYS	variant	UNP P0DTU4
J	201	ASP	ASN	variant	UNP P0DTU4

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	63	Total O 63 63	0	0
6	B	32	Total O 32 32	0	0
6	C	4	Total O 4 4	0	0
6	D	18	Total O 18 18	0	0
6	E	17	Total O 17 17	0	0
6	F	67	Total O 67 67	0	0
6	G	21	Total O 21 21	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	2	Total O 2 2	0	0
6	I	14	Total O 14 14	0	0
6	J	8	Total O 8 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA class I histocompatibility antigen, alpha chain E




- Molecule 1: HLA class I histocompatibility antigen, alpha chain E




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Large envelope protein







- Molecule 3: Large envelope protein

Chain H: 89% 11%





- Molecule 4: T cell receptor alpha variable 12-2,T cell receptor alpha chain MC.7.G5

Chain D: 5% 78% 15% 8%




- Molecule 4: T cell receptor alpha variable 12-2,T cell receptor alpha chain MC.7.G5

Chain I: 8% 79% 10% 10%




- Molecule 5: T cell receptor beta variable 6-5,T cell receptor beta chain MC.7.G5


Chain E: 5% 86% 10% 2%



- Molecule 5: T cell receptor beta variable 6-5,T cell receptor beta chain MC.7.G5

Chain J: 9% 77% 16% 6%





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.70Å 155.06Å 93.41Å 90.00° 97.04° 90.00°	Depositor
Resolution (Å)	79.57 – 2.25 79.57 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.5 (79.57-2.25) 98.5 (79.57-2.25)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.25Å)	Xtrriage
Refinement program	REFMAC 5.8.0403	Depositor
R, R_{free}	0.209 , 0.252 0.222 , 0.263	Depositor DCC
R_{free} test set	5136 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	51.8	Xtrriage
Anisotropy	0.348	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13060	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/2320	0.84	0/3154
1	F	0.50	0/2273	0.84	2/3089 (0.1%)
2	B	0.47	0/860	0.85	0/1162
2	G	0.49	0/860	0.80	0/1162
3	C	0.54	0/74	0.82	0/100
3	H	0.56	0/74	0.84	0/100
4	D	0.45	0/1448	0.81	0/1957
4	I	0.45	0/1410	0.83	0/1904
5	E	0.41	0/1961	0.85	0/2670
5	J	0.42	0/1866	0.86	1/2540 (0.0%)
All	All	0.46	0/13146	0.84	3/17838 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	F	0	4
2	G	0	2
4	I	0	1
5	E	0	5
5	J	0	3
All	All	0	16

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	157	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	F	157	ARG	NE-CZ-NH2	-5.71	117.44	120.30
5	J	95	ARG	CB-CA-C	-5.04	100.33	110.40

There are no chirality outliers.

5 of 16 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	111	ARG	Sidechain
5	E	191	ARG	Sidechain
5	E	193	ARG	Sidechain
5	E	203	ARG	Sidechain
5	E	225	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2253	0	2092	20	0
1	F	2208	0	2056	16	0
2	B	837	0	803	8	0
2	G	837	0	803	6	0
3	C	72	0	85	2	0
3	H	72	0	85	1	0
4	D	1421	0	1352	20	0
4	I	1385	0	1318	16	0
5	E	1910	0	1811	27	0
5	J	1819	0	1732	40	0
6	A	63	0	0	0	0
6	B	32	0	0	2	0
6	C	4	0	0	0	0
6	D	18	0	0	0	0
6	E	17	0	0	1	0
6	F	67	0	0	2	0
6	G	21	0	0	0	0
6	H	2	0	0	1	0
6	I	14	0	0	0	0
6	J	8	0	0	0	0
All	All	13060	0	12137	127	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:I:124:LEU:HD22	5:J:127:GLU:O	1.28	1.31
4:I:124:LEU:HD21	5:J:129:SER:N	1.83	0.94
4:I:124:LEU:CD2	5:J:127:GLU:O	2.22	0.86
5:E:108:ARG:HG3	5:E:108:ARG:HH11	1.47	0.79
4:I:123:GLN:O	4:I:124:LEU:HD23	1.86	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/276 (100%)	265 (96%)	10 (4%)	0	100	100
1	F	267/276 (97%)	257 (96%)	10 (4%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	G	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	H	7/9 (78%)	7 (100%)	0	0	100	100
4	D	178/199 (89%)	167 (94%)	11 (6%)	0	100	100
4	I	171/199 (86%)	162 (95%)	9 (5%)	0	100	100
5	E	238/243 (98%)	221 (93%)	17 (7%)	0	100	100
5	J	225/243 (93%)	209 (93%)	16 (7%)	0	100	100
All	All	1564/1654 (95%)	1485 (95%)	79 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/237 (100%)	234 (98%)	4 (2%)	56	66
1	F	233/237 (98%)	228 (98%)	5 (2%)	48	57
2	B	95/95 (100%)	94 (99%)	1 (1%)	70	79
2	G	95/95 (100%)	93 (98%)	2 (2%)	48	57
3	C	9/9 (100%)	9 (100%)	0	100	100
3	H	9/9 (100%)	9 (100%)	0	100	100
4	D	161/175 (92%)	154 (96%)	7 (4%)	25	28
4	I	156/175 (89%)	149 (96%)	7 (4%)	23	26
5	E	209/211 (99%)	204 (98%)	5 (2%)	44	53
5	J	201/211 (95%)	194 (96%)	7 (4%)	31	38
All	All	1406/1454 (97%)	1368 (97%)	38 (3%)	40	48

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	I	173	SER
5	J	141	LEU
4	I	175	VAL
5	J	55	THR
5	J	191	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	197	HIS
1	A	226	GLN
5	E	11	GLN
5	E	65	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	276/276 (100%)	-0.01	5 (1%) 67 68	32, 57, 95, 136	1 (0%)
1	F	271/276 (98%)	-0.06	5 (1%) 67 68	38, 55, 97, 125	0
2	B	100/100 (100%)	-0.07	2 (2%) 64 64	43, 54, 80, 98	0
2	G	100/100 (100%)	0.17	2 (2%) 64 64	47, 64, 88, 107	0
3	C	9/9 (100%)	-0.33	0 100 100	44, 46, 51, 51	0
3	H	9/9 (100%)	-0.11	0 100 100	40, 42, 48, 52	0
4	D	184/199 (92%)	0.31	9 (4%) 36 35	42, 60, 122, 144	0
4	I	179/199 (89%)	0.41	15 (8%) 18 18	40, 61, 123, 135	0
5	E	240/243 (98%)	0.55	11 (4%) 38 37	46, 79, 121, 136	0
5	J	229/243 (94%)	0.74	21 (9%) 16 15	42, 80, 123, 145	0
All	All	1597/1654 (96%)	0.26	70 (4%) 39 39	32, 63, 115, 145	1 (0%)

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	221	GLY	4.9
4	D	154	VAL	3.8
5	J	181	LEU	3.7
4	I	184	ALA	3.6
5	J	206	PHE	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.