

wwPDB X-ray Structure Validation Summary Report (i)

Jun 27, 2024 – 06:08 PM EDT

PDB ID : 8TD3

Title : Structure of PYCR1 complexed with NADH and (S)-Tetrahydro-2H-pyran-2-

carboxylic acid

Authors : Tanner, J.J.; Meeks, K.R.

Deposited on : 2023-07-02

Resolution : 1.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.37.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	316	85%	•	11%
1	В	316	84%		11%
1	С	316	84%		12%
1	D	316	84%	5%	11%
1	E	316	83%	5%	12%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 11225 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Pyrroline-5-carboxylate reductase 1, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	281	Total	С	N	Ο	S	0	8	0
1 A	201	2044	1292	358	380	14	0	8		
1	В	280	Total	С	N	О	S	0	5	0
1	Ъ	280	2030	1288	351	377	14	U	0	
1	С	277	Total	С	N	О	S	0	3	0
1		211	2004	1263	350	377	14	U		
1	D	281	Total	С	N	О	S	0	4	0
1	ש	201	2049	1290	362	383	14	U	4	
1	Е	277	Total	С	N	О	S	0	5	0
1	ш	211	1999	1261	354	370	14	U	9	

There are 110 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP P32322
A	-20	HIS	-	expression tag	UNP P32322
A	-19	HIS	-	expression tag	UNP P32322
A	-18	HIS	-	expression tag	UNP P32322
A	-17	HIS	-	expression tag	UNP P32322
A	-16	HIS	_	expression tag	UNP P32322
A	-15	HIS	-	expression tag	UNP P32322
A	-14	SER	-	expression tag	UNP P32322
A	-13	SER	-	expression tag	UNP P32322
A	-12	GLY	-	expression tag	UNP P32322
A	-11	VAL	_	expression tag	UNP P32322
A	-10	ASP	-	expression tag	UNP P32322
A	-9	LEU	-	expression tag	UNP P32322
A	-8	GLY	-	expression tag	UNP P32322
A	-7	THR	-	expression tag	UNP P32322
A	-6	GLU	-	expression tag	UNP P32322
A	-5	ASN	-	expression tag	UNP P32322
A	-4	LEU	-	expression tag	UNP P32322
A	-3	TYR	-	expression tag	UNP P32322



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-2	PHE	-	expression tag	UNP P32322
A	-1	GLN	-	expression tag	UNP P32322
A	0	SER	_	expression tag	UNP P32322
В	-21	MET	-	initiating methionine	UNP P32322
В	-20	HIS	-	expression tag	UNP P32322
В	-19	HIS	-	expression tag	UNP P32322
В	-18	HIS	-	expression tag	UNP P32322
В	-17	HIS	-	expression tag	UNP P32322
В	-16	HIS	-	expression tag	UNP P32322
В	-15	HIS	-	expression tag	UNP P32322
В	-14	SER	-	expression tag	UNP P32322
В	-13	SER	-	expression tag	UNP P32322
В	-12	GLY	-	expression tag	UNP P32322
В	-11	VAL	-	expression tag	UNP P32322
В	-10	ASP	-	expression tag	UNP P32322
В	-9	LEU	-	expression tag	UNP P32322
В	-8	GLY	-	expression tag	UNP P32322
В	-7	THR	-	expression tag	UNP P32322
В	-6	GLU	-	expression tag	UNP P32322
В	-5	ASN	-	expression tag	UNP P32322
В	-4	LEU	-	expression tag	UNP P32322
В	-3	TYR	-	expression tag	UNP P32322
В	-2	PHE	_	expression tag	UNP P32322
В	-1	GLN	-	expression tag	UNP P32322
В	0	SER	-	expression tag	UNP P32322
С	-21	MET	_	initiating methionine	UNP P32322
С	-20	HIS	-	expression tag	UNP P32322
С	-19	HIS	-	expression tag	UNP P32322
С	-18	HIS	_	expression tag	UNP P32322
С	-17	HIS	-	expression tag	UNP P32322
С	-16	HIS	-	expression tag	UNP P32322
С	-15	HIS	_	expression tag	UNP P32322
С	-14	SER	-	expression tag	UNP P32322
С	-13	SER	_	expression tag	UNP P32322
С	-12	GLY	-	expression tag	UNP P32322
С	-11	VAL	-	expression tag	UNP P32322
С	-10	ASP	-	expression tag	UNP P32322
С	-9	LEU	-	expression tag	UNP P32322
С	-8	GLY		expression tag	UNP P32322
С	-7	THR	-	expression tag	UNP P32322
С	-6	GLU	-	expression tag	UNP P32322
С	-5	ASN	-	expression tag	UNP P32322



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
С	-4	LEU	-	expression tag	UNP P32322
С	-3	TYR	-	expression tag	UNP P32322
С	-2	PHE	_	expression tag	UNP P32322
С	-1	GLN	-	expression tag	UNP P32322
С	0	SER	-	expression tag	UNP P32322
D	-21	MET	_	initiating methionine	UNP P32322
D	-20	HIS	-	expression tag	UNP P32322
D	-19	HIS	-	expression tag	UNP P32322
D	-18	HIS	-	expression tag	UNP P32322
D	-17	HIS	-	expression tag	UNP P32322
D	-16	HIS	-	expression tag	UNP P32322
D	-15	HIS	-	expression tag	UNP P32322
D	-14	SER	-	expression tag	UNP P32322
D	-13	SER	_	expression tag	UNP P32322
D	-12	GLY	-	expression tag	UNP P32322
D	-11	VAL	-	expression tag	UNP P32322
D	-10	ASP	-	expression tag	UNP P32322
D	-9	LEU	-	expression tag	UNP P32322
D	-8	GLY	-	expression tag	UNP P32322
D	-7	THR	_	expression tag	UNP P32322
D	-6	GLU	-	expression tag	UNP P32322
D	-5	ASN	-	expression tag	UNP P32322
D	-4	LEU	-	expression tag	UNP P32322
D	-3	TYR	-	expression tag	UNP P32322
D	-2	PHE	-	expression tag	UNP P32322
D	-1	GLN	-	expression tag	UNP P32322
D	0	SER	-	expression tag	UNP P32322
E	-21	MET	-	initiating methionine	UNP P32322
Е	-20	HIS	-	expression tag	UNP P32322
Е	-19	HIS	-	expression tag	UNP P32322
Е	-18	HIS	-	expression tag	UNP P32322
Е	-17	HIS	-	expression tag	UNP P32322
Е	-16	HIS	-	expression tag	UNP P32322
Е	-15	HIS	-	expression tag	UNP P32322
Е	-14	SER	-	expression tag	UNP P32322
Е	-13	SER	-	expression tag	UNP P32322
Е	-12	GLY	-	expression tag	UNP P32322
Е	-11	VAL	-	expression tag	UNP P32322
Е	-10	ASP	-	expression tag	UNP P32322
Е	-9	LEU	-	expression tag	UNP P32322
Е	-8	GLY	-	expression tag	UNP P32322
E	-7	THR	_	expression tag	UNP P32322



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-6	GLU	-	expression tag	UNP P32322
Е	-5	ASN	-	expression tag	UNP P32322
E	-4	LEU	-	expression tag	UNP P32322
E	-3	TYR	-	expression tag	UNP P32322
E	-2	PHE	-	expression tag	UNP P32322
Е	-1	GLN	-	expression tag	UNP P32322
Е	0	SER	-	expression tag	UNP P32322


• Molecule 2 is 1,4-DIHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE (three-letter code: NAI) (formula: $C_{21}H_{29}N_7O_{14}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	Λ	1	Total	С	N	О	Р	0	0
2	Λ		44	21	7	14	2	U	
2	В	1	Total	С	N	О	Р	0	0
2	Б	1	44	21	7	14	2		U
2	С	1	Total	С	N	О	Р	0	0
2		1	44	21	7	14	2	U	
2	D	1	Total	С	N	О	Р	0	0
2		$D \mid 1 \mid$	44	21	7	14	2	U	
2	Е	1	Total	С	N	О	Р	0	0
	157	1	44	21	7	14	2	U	U


 \bullet Molecule 3 is (2S)-oxane-2-carboxylic acid (three-letter code: GJM) (formula: $C_6H_{10}O_3)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Δ	1	Total C O	0	0
	Λ	1	9 6 3	U	0
3	В	1	Total C O	0	0
	D	1	9 6 3		0
3	\mathbf{C}	1	Total C O	0	0
		1	9 6 3	U	
3	D	1	Total C O	0	0
	D	1	9 6 3	U	
3	Е	1	Total C O	0	0
	L	1	9 6 3	U	


• Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	Е	1	Total C O 7 4 3	0	0
4	E	1	Total C O 7 4 3	0	0

 \bullet Molecule 5 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $\mathrm{C_6H_{14}O_4}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total C O 10 6 4	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	187	Total O 187 187	0	0
6	В	160	Total O 160 160	0	0
6	С	155	Total O 155 155	0	0
6	D	149	Total O 149 149	0	0
6	Е	152	Total O 152 152	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial



• Molecule 1: Pyrroline-5-carboxylate reductase 1, mitochondrial





4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants	111.62Å 179.96Å 87.72Å	Depositor
a, b, c, α , β , γ	90.00° 106.86° 90.00°	Depositor
Resolution (Å)	91.86 - 1.67	Depositor
Resolution (A)	91.86 - 1.67	EDS
% Data completeness	98.2 (91.86-1.67)	Depositor
(in resolution range)	98.2 (91.86-1.67)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.63 (at 1.67Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
P. P.	0.169 , 0.191	Depositor
R, R_{free}	0.165 , 0.187	DCC
R_{free} test set	9320 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.245	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 43.2	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11225	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GJM, PGE, NAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles	
IVIOI	Mol Chain		# Z > 5	RMSZ	# Z >5
1	A	0.34	0/2098	0.58	0/2849
1	В	0.34	0/2077	0.56	0/2819
1	С	0.34	0/2044	0.55	0/2775
1	D	0.35	1/2093~(0.0%)	0.55	0/2840
1	Е	0.34	0/2044	0.56	0/2775
All	All	0.34	$1/10356 \ (0.0\%)$	0.56	0/14058

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
1	D	120	CYS	CB-SG	-5.32	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2044	0	2074	8	0
1	В	2030	0	2044	9	0
1	С	2004	0	2016	9	0
1	D	2049	0	2064	9	0
1	Е	1999	0	2022	9	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	44	0	27	0	0
2	В	44	0	27	0	0
2	С	44	0	27	0	0
2	D	44	0	27	1	0
2	Ε	44	0	27	1	0
3	A	9	0	0	0	0
3	В	9	0	0	0	0
3	С	9	0	0	0	0
3	D	9	0	0	0	0
3	Ε	9	0	0	0	0
4	A	7	0	10	0	0
4	Ε	14	0	20	2	0
5	D	10	0	14	0	0
6	A	187	0	0	0	0
6	В	160	0	0	0	0
6	С	155	0	0	3	0
6	D	149	0	0	0	0
6	Ε	152	0	0	0	0
All	All	11225	0	10399	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 40 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ (ext{\AA})$	overlap (Å)
1:E:133:THR:HG21	1:E:153[A]:LEU:HD13	1.66	0.77
1:E:87:GLU:OE1	1:E:89:ARG:NH1	2.25	0.69
1:A:133:THR:HG21	1:A:153[A]:LEU:HD13	1.75	0.68
1:C:210:LEU:HD13	1:D:210:LEU:HD11	1.76	0.67
1:C:190:ASP:OD1	1:C:199[B]:ARG:NH1	2.27	0.65

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the r	number of	residues for	which	the	backbone	conformation	was
analysed, and the total number of	residues.						

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	A	$287/316 \ (91\%)$	284 (99%)	3 (1%)	0	100	100
1	В	283/316~(90%)	280 (99%)	3 (1%)	0	100	100
1	\mathbf{C}	$278/316\ (88\%)$	276 (99%)	2 (1%)	0	100	100
1	D	$283/316 \ (90\%)$	278 (98%)	5 (2%)	0	100	100
1	E	280/316 (89%)	276 (99%)	4 (1%)	0	100	100
All	All	1411/1580 (89%)	1394 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	210/251 (84%)	210 (100%)	0	100 100
1	В	206/251 (82%)	205 (100%)	1 (0%)	88 83
1	С	206/251 (82%)	206 (100%)	0	100 100
1	D	211/251 (84%)	210 (100%)	1 (0%)	88 83
1	E	203/251 (81%)	201 (99%)	2 (1%)	76 65
All	All	1036/1255 (82%)	1032 (100%)	4 (0%)	91 86

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	232	SER
1	D	150	GLU
1	Е	210	LEU
1	Е	232	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	В	62	HIS
1	С	10	GLN
1	Е	140	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	gles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
4	PEG	Е	403	-	6,6,6	0.20	0	5,5,5	0.09	0
2	NAI	С	401	_	42,48,48	1.15	3 (7%)	47,73,73	1.50	8 (17%)
5	PGE	D	303	-	9,9,9	0.32	0	8,8,8	0.36	0
2	NAI	A	401	-	42,48,48	1.19	5 (11%)	47,73,73	1.49	8 (17%)
2	NAI	D	302	-	42,48,48	1.23	5 (11%)	47,73,73	1.46	9 (19%)
3	GJM	С	402	-	9,9,9	1.11	0	10,11,11	1.76	2 (20%)
3	GJM	Е	402	-	9,9,9	1.00	0	10,11,11	1.35	1 (10%)
4	PEG	A	403	-	6,6,6	0.16	0	5,5,5	0.07	0
3	GJM	A	402	_	9,9,9	1.11	0	10,11,11	1.18	1 (10%)
2	NAI	Е	401	-	42,48,48	1.21	3 (7%)	47,73,73	1.48	10 (21%)



Mol	Trino	Chain	Res	Link	Bond lengths			Bond angles		
Moi Tyl	Type		nes	tes Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GJM	D	301	-	9,9,9	1.09	0	10,11,11	1.23	1 (10%)
3	GJM	В	301	-	9,9,9	1.01	0	10,11,11	1.59	2 (20%)
2	NAI	В	302	-	42,48,48	1.22	4 (9%)	47,73,73	1.42	10 (21%)
4	PEG	Е	404	-	6,6,6	0.17	0	5,5,5	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	Е	403	-	-	3/4/4/4	-
2	NAI	С	401	-	-	3/25/72/72	0/5/5/5
5	PGE	D	303	-	-	2/7/7/7	-
2	NAI	A	401	-	-	3/25/72/72	0/5/5/5
2	NAI	D	302	-	-	3/25/72/72	0/5/5/5
3	GJM	С	402	-	-	0/4/12/12	0/1/1/1
3	GJM	E	402	_	-	2/4/12/12	0/1/1/1
4	PEG	A	403	-	-	3/4/4/4	-
3	GJM	A	402	-	-	1/4/12/12	0/1/1/1
2	NAI	Е	401	-	-	2/25/72/72	0/5/5/5
3	GJM	D	301	-	-	0/4/12/12	0/1/1/1
3	GJM	В	301	-	-	1/4/12/12	0/1/1/1
2	NAI	В	302	_	-	2/25/72/72	0/5/5/5
4	PEG	Е	404	-	-	2/4/4/4	-

The worst 5 of 20 bond length outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(ext{\AA})$
2	D	302	NAI	PA-O5B	3.28	1.72	1.59
2	Ε	401	NAI	PA-O5B	3.20	1.72	1.59
2	В	302	NAI	PN-O5D	2.97	1.71	1.59
2	С	401	NAI	PN-O5D	2.82	1.70	1.59
2	E	401	NAI	PN-O5D	2.79	1.70	1.59

The worst 5 of 52 bond angle outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	${f Atoms}$	\mathbf{Z}	$Observed(^{o})$	$\operatorname{Ideal}(^{o})$
2	D	302	NAI	PN-O3-PA	-4.61	117.00	132.83



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^{o})$
2	Ε	401	NAI	PN-O3-PA	-4.36	117.85	132.83
2	С	401	NAI	PN-O3-PA	-4.36	117.86	132.83
2	В	302	NAI	PN-O3-PA	-4.30	118.06	132.83
2	A	401	NAI	PN-O3-PA	-4.11	118.71	132.83

There are no chirality outliers.

5 of 27 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	PEG	O2-C3-C4-O4
5	D	303	PGE	O1-C1-C2-O2
4	Е	403	PEG	C1-C2-O2-C3
4	Е	403	PEG	O1-C1-C2-O2
4	Е	403	PEG	O2-C3-C4-O4


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	403	PEG	2	0
2	D	302	NAI	1	0
2	Е	401	NAI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	281/316 (88%)	-0.21	6 (2%) 63 67	19, 27, 49, 72	0
1	В	280/316 (88%)	-0.21	4 (1%) 75 79	19, 30, 52, 72	0
1	С	277/316 (87%)	-0.23	1 (0%) 92 93	20, 31, 53, 71	0
1	D	281/316 (88%)	-0.09	5 (1%) 68 72	19, 32, 55, 75	0
1	Е	277/316 (87%)	-0.24	2 (0%) 87 89	19, 31, 52, 77	0
All	All	1396/1580 (88%)	-0.20	18 (1%) 77 80	19, 30, 53, 77	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-3	TYR	9.2
1	В	-3	TYR	4.1
1	В	-2	PHE	3.9
1	В	-1	GLN	3.4
1	A	-2	PHE	3.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	PEG	Е	404	7/7	0.79	0.21	42,45,53,56	0
4	PEG	A	403	7/7	0.81	0.23	40,45,48,51	0
4	PEG	Е	403	7/7	0.82	0.26	40,45,48,53	0
3	GJM	В	301	9/9	0.91	0.09	22,28,31,37	0
3	GJM	С	402	9/9	0.92	0.09	22,25,29,33	9
5	PGE	D	303	10/10	0.92	0.20	42,47,55,55	0
3	GJM	Е	402	9/9	0.93	0.08	22,25,29,32	9
3	GJM	A	402	9/9	0.94	0.08	25,28,32,32	0
3	GJM	D	301	9/9	0.95	0.06	26,29,33,34	0
2	NAI	Е	401	44/44	0.96	0.08	24,31,37,41	0
2	NAI	С	401	44/44	0.97	0.07	25,30,35,38	0
2	NAI	D	302	44/44	0.97	0.08	25,31,38,40	0
2	NAI	A	401	44/44	0.97	0.08	23,26,31,34	0
2	NAI	В	302	44/44	0.97	0.07	23,28,31,33	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GJM B 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GJM C 402:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around GJM D 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around NAI E 401:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAI C 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAI D 302:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAI A 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAI B 302:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

