

wwPDB X-ray Structure Validation Summary Report (i)

Oct 2, 2024 – 02:08 PM EDT

PDB ID : 9DSZ

Title : Crystal structure of ADP-ribose diphosphatase from Klebsiella pneumoniae

(DTP bound)

Authors : Seattle Structural Genomics Center for Infectious Disease; Seattle Structural

Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2024-09-30

Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS: 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

 $Density-Fitness \quad : \quad 1.0.11$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\#\text{Entries, resolution range}(\mathring{\mathbf{A}}))$
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	218	81%	7%	12%
1	В	218	81%	8%	11%
1	С	218	86%	69	% 9%
1	D	218	78%	11%	11%



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	E	218	12%	• 9%
1	F	218	8%	6% 12%
1	G	218	6% 87%	6% 7%
1	Н	218	85%	• 11%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13459 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ADP-ribose pyrophosphatase.

Mol	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace	
1	A	192	Total	С	N	О	S	0	5	0	
1	A	192	1547	984	268	292	3	0	9	0	
1	В	195	Total	С	N	О	S	0	2	0	
1	Б	199	1562	989	273	297	3	0	2		
1	С	199	Total	С	N	О	S	0	2	0	
1		199	1585	1004	277	301	3	0	<u> </u>		
1	D	195	Total	С	N	О	S	0	1	0	
1	D	190	1559	985	275	296	3	U			
1	E	199	Total	С	N	О	S	0	2	0	
1	ш	199	1603	1016	285	298	4	0	2		
1	F	191	Total	С	N	О	S	0	3	0	
1	I.	191	1532	972	267	290	3	U	0	U	
1	G	203	Total	С	N	О	\mathbf{S}	0	2	0	
1	G	200	1607	1018	281	305	3	U	2	U	
1	Н	194	Total	С	N	О	S	0	4	0	
1	11	134	1551	985	270	293	3		1		

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A0A0H3GVQ7
A	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
A	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
A	-4	HIS	_	expression tag	UNP A0A0H3GVQ7
A	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
A	-2	HIS	_	expression tag	UNP A0A0H3GVQ7
A	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
A	0	HIS	_	expression tag	UNP A0A0H3GVQ7
В	-7	MET	-	expression tag	UNP A0A0H3GVQ7
В	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
В	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
В	-4	HIS	_	expression tag	UNP A0A0H3GVQ7
В	-3	HIS	-	expression tag	UNP A0A0H3GVQ7



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
В	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
В	0	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-7	MET	_	expression tag	UNP A0A0H3GVQ7
С	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
С	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
С	0	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-7	MET	-	expression tag	UNP A0A0H3GVQ7
D	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
D	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
D	0	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-7	MET	-	expression tag	UNP A0A0H3GVQ7
Е	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
Е	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-1	HIS	_	expression tag	UNP A0A0H3GVQ7
Е	0	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-7	MET	-	expression tag	UNP A0A0H3GVQ7
F	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
F	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
F	0	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-7	MET	-	expression tag	UNP A0A0H3GVQ7
G	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
G	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-1	HIS	_	expression tag	UNP A0A0H3GVQ7



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-7	MET	-	expression tag	UNP A0A0H3GVQ7
Н	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
Н	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	0	HIS	-	expression tag	UNP A0A0H3GVQ7

• Molecule 2 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula: $C_{10}H_{16}N_5O_{12}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
	Λ	1	30	10	5	12	3	U	U
2	В	1	Total	С	N	Ο	Р	0	0
	Ъ	1	30	10	5	12	3	U	0
2	\mathbf{C}	1	Total	С	N	Ο	Р	0	0
2		1	30	10	5	12	3	U	
2	D	1	Total	С	N	Ο	Р	0	0
	D	1	30	10	5	12	3	U	0
2	E	1	Total	С	N	О	Р	0	0
	<u> 1</u> 2	1	30	10	5	12	3	U	0
2	F	1	Total	С	N	O	Р	0	0
	I.	1	30	10	5	12	3	U	U



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Н	1	Total C N O 18 10 5 3	0	0
2	Н	1	Total C N O P 26 10 5 9 2	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	Е	1	Total Mg 1 1	0	0
3	F	2	Total Mg 2 2	0	0
3	G	2	Total Mg 2 2	0	0
3	Н	1	Total Mg 1 1	0	0

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	88	Total O 88 88	0	0
4	В	86	Total O 86 86	0	0
4	С	81	Total O 81 81	0	1
4	D	71	Total O 71 71	0	0
4	E	75	Total O 75 75	0	0
4	F	79	Total O 79 79	0	0
4	G	110	Total O 110 110	0	0



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	Н	89	Total O 89 89	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ADP-ribose pyrophosphatase








• Molecule 1: ADP-ribose pyrophosphatase








• Molecule 1: ADP-ribose pyrophosphatase







• Molecule 1: ADP-ribose pyrophosphatase





K210

 \bullet Molecule 1: ADP-ribose pyrophosphatase

Chain H: 85% • 11%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	92.84Å 79.31Å 115.83Å	Depositor
a, b, c, α , β , γ	90.00° 90.30° 90.00°	Depositor
Resolution (Å)	72.63 - 1.90	Depositor
Resolution (A)	72.63 - 1.90	EDS
% Data completeness	99.6 (72.63-1.90)	Depositor
(in resolution range)	99.7 (72.63-1.90)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.01 (at 1.90Å)	Xtriage
Refinement program	PHENIX (dev_5396: ???)	Depositor
D.D.	0.196 , 0.233	Depositor
R, R_{free}	0.203 , 0.238	DCC
R_{free} test set	6561 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 45.2	EDS
L-test for twinning ²	$< L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	0.055 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13459	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 63.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0038e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.38	0/1588	0.62	0/2154	
1	В	0.36	0/1597	0.61	0/2162	
1	С	0.37	0/1621	0.63	0/2194	
1	D	0.38	0/1591	0.62	0/2155	
1	Е	0.38	0/1642	0.63	0/2222	
1	F	0.37	0/1570	0.62	0/2126	
1	G	0.42	0/1644	0.65	0/2229	
1	Н	0.38	0/1592	0.66	0/2158	
All	All	0.38	0/12845	0.63	0/17400	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1547	0	1524	15	0
1	В	1562	0	1530	13	0
1	С	1585	0	1552	9	0
1	D	1559	0	1525	18	0
1	Е	1603	0	1571	8	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1532	0	1507	10	0
1	G	1607	0	1568	16	0
1	Н	1551	0	1516	6	0
2	A	30	0	12	1	0
2	В	30	0	12	0	0
2	С	30	0	12	0	0
2	D	30	0	12	0	0
2	Ε	30	0	12	0	0
2	F	30	0	12	0	0
2	Н	44	0	24	4	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0
3	Ε	1	0	0	0	0
3	F	2	0	0	0	0
3	G	2	0	0	0	0
3	Н	1	0	0	0	0
4	A	88	0	0	0	0
4	В	86	0	0	1	0
4	С	81	0	0	0	0
4	D	71	0	0	0	0
4	Е	75	0	0	1	0
4	F	79	0	0	0	0
4	G	110	0	0	1	0
4	Н	89	0	0	1	0
All	All	13459	0	12389	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 78 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	$ \text{overlap } (\mathring{\mathrm{A}})$
1:B:75:VAL:HG13	1:B:166:ILE:HG23	1.56	0.86
1:F:75:VAL:HG13	1:F:166:ILE:HG23	1.64	0.78
1:H:75:VAL:HG13	1:H:166:ILE:HG23	1.66	0.78
1:A:75[A]:VAL:HG13	1:A:166:ILE:HG23	1.67	0.77
1:E:75:VAL:HG13	1:E:166:ILE:HG23	1.69	0.74

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	193/218 (88%)	187 (97%)	6 (3%)	0	100	100
1	В	193/218 (88%)	187 (97%)	6 (3%)	0	100	100
1	С	197/218 (90%)	191 (97%)	6 (3%)	0	100	100
1	D	192/218 (88%)	185 (96%)	7 (4%)	0	100	100
1	E	195/218 (89%)	189 (97%)	6 (3%)	0	100	100
1	F	190/218 (87%)	184 (97%)	6 (3%)	0	100	100
1	G	203/218 (93%)	195 (96%)	8 (4%)	0	100	100
1	Н	194/218 (89%)	188 (97%)	6 (3%)	0	100	100
All	All	1557/1744 (89%)	1506 (97%)	51 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	163/183~(89%)	163 (100%)	0	100 100
1	В	163/183~(89%)	163 (100%)	0	100 100
1	С	165/183~(90%)	165 (100%)	0	100 100
1	D	163/183~(89%)	162 (99%)	1 (1%)	84 86
1	E	169/183~(92%)	169 (100%)	0	100 100
1	F	161/183~(88%)	161 (100%)	0	100 100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	G	166/183 (91%)	166 (100%)	0	100	100	
1	Н	161/183 (88%)	160 (99%)	1 (1%)	84	86	
All	All	1311/1464 (90%)	1309 (100%)	2 (0%)	92	93	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	22	GLU
1	Н	26	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 18 ligands modelled in this entry, 10 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Вс	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DTP	D	301	3	28,32,32	0.63	0	35,50,50	0.92	2 (5%)
2	DTP	С	301	3	28,32,32	0.64	0	35,50,50	0.86	1 (2%)
2	DTP	В	301	3	28,32,32	0.74	0	35,50,50	0.91	1 (2%)
2	DTP	Е	301	-	28,32,32	0.62	0	35,50,50	0.88	1 (2%)
2	DTP	Н	301	-	17,20,32	0.70	0	18,29,50	1.19	2 (11%)
2	DTP	Н	302	3	24,28,32	0.71	0	30,43,50	1.01	2 (6%)
2	DTP	F	301	-	28,32,32	0.69	0	35,50,50	0.95	2 (5%)
2	DTP	A	301	3	28,32,32	0.64	0	35,50,50	0.92	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DTP	D	301	3	-	2/18/34/34	0/3/3/3
2	DTP	С	301	3	-	9/18/34/34	0/3/3/3
2	DTP	В	301	3	-	8/18/34/34	0/3/3/3
2	DTP	Е	301	-	-	5/18/34/34	0/3/3/3
2	DTP	Н	301	-	-	1/2/18/34	0/3/3/3
2	DTP	Н	302	3	-	2/12/28/34	0/3/3/3
2	DTP	F	301	-	-	4/18/34/34	0/3/3/3
2	DTP	A	301	3	-	5/18/34/34	0/3/3/3

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
2	A	301	DTP	C5-C6-N6	2.40	123.96	120.31
2	Н	301	DTP	C5-C6-N6	2.27	123.77	120.31
2	F	301	DTP	O4'-C1'-N9	-2.27	105.74	108.75
2	С	301	DTP	C5-C6-N6	2.26	123.75	120.31
2	Н	302	DTP	O4'-C1'-N9	-2.24	105.78	108.75

There are no chirality outliers.

5 of 36 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	301	DTP	C5'-O5'-PA-O1A
2	В	301	DTP	C5'-O5'-PA-O1A
2	В	301	DTP	C5'-O5'-PA-O2A
2	В	301	DTP	C5'-O5'-PA-O3A
2	С	301	DTP	C5'-O5'-PA-O1A


There are no ring outliers.

2 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Н	301	DTP	4	0
2	A	301	DTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$	$OWAB(\AA^2)$	Q<0.9
1	A	192/218 (88%)	0.63	11 (5%) 30 32	19, 41, 70, 97	5 (2%)
1	В	195/218 (89%)	0.73	24 (12%) 9 10	23, 40, 86, 116	2 (1%)
1	С	199/218 (91%)	0.48	12 (6%) 29 30	18, 38, 61, 85	2 (1%)
1	D	195/218 (89%)	0.75	25 (12%) 9 9	22, 42, 79, 116	1 (0%)
1	E	199/218 (91%)	0.88	26 (13%) 8 8	19, 43, 82, 105	2 (1%)
1	F	191/218 (87%)	0.76	17 (8%) 17 17	22, 41, 82, 120	3 (1%)
1	G	203/218 (93%)	0.34	12 (5%) 29 30	15, 31, 65, 90	2 (0%)
1	Н	194/218 (88%)	0.48	9 (4%) 38 39	18, 36, 65, 80	4 (2%)
All	All	1568/1744 (89%)	0.63	136 (8%) 17 18	15, 39, 76, 120	21 (1%)

The worst 5 of 136 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	156	ILE	6.1
1	Н	209	THR	5.5
1	D	114	LEU	5.1
1	A	10	THR	5.1
1	Е	0	HIS	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B ext{-}factors}({f \AA}^2)$	Q<0.9
2	DTP	Е	301	30/30	0.63	0.16	53,98,118,122	0
2	DTP	С	301	30/30	0.66	0.14	47,84,121,130	0
2	DTP	A	301	30/30	0.66	0.14	38,82,122,130	0
3	MG	С	302	1/1	0.72	0.15	61,61,61,61	0
3	MG	Н	303	1/1	0.74	0.19	79,79,79,79	0
2	DTP	В	301	30/30	0.77	0.13	30,70,102,108	0
2	DTP	D	301	30/30	0.80	0.12	40,70,95,111	0
2	DTP	F	301	30/30	0.80	0.13	33,71,94,103	0
2	DTP	Н	302	26/30	0.82	0.13	29,60,97,107	0
3	MG	A	302	1/1	0.82	0.13	53,53,53,53	0
3	MG	F	303	1/1	0.84	0.11	59,59,59,59	0
2	DTP	Н	301	18/30	0.85	0.13	30,42,69,72	0
3	MG	Е	302	1/1	0.85	0.17	60,60,60,60	0
3	MG	В	302	1/1	0.86	0.10	71,71,71,71	0
3	MG	D	302	1/1	0.90	0.10	66,66,66,66	0
3	MG	G	301	1/1	0.93	0.07	37,37,37,37	0
3	MG	F	302	1/1	0.95	0.14	46,46,46,46	0
3	MG	G	302	1/1	0.98	0.05	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.










Electron density around DTP C 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around DTP A 301:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




















Electron density around DTP B 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around DTP D 301:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around DTP F 301:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




















Electron density around MG F 303: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around DTP H 301:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)


















Electron density around MG D 302: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




Electron density around MG G 301:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















6.5 Other polymers (i)

There are no such residues in this entry.

