

Full wwPDB X-ray Structure Validation Report (i)

Jun 12, 2024 – 10:58 PM EDT

PDB ID : 1DNW

Title: HUMAN MYELOPEROXIDASE-CYANIDE-THIOCYANATE COMPLEX

Authors: Blair-Johnson, M.; Fiedler, T.J.; Fenna, R.E.

Deposited on : 1999-12-16

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned} ext{Similar resolution} \ (\# ext{Entries, resolution range}(ext{Å})) \end{aligned}$		
R_{free}	130704	6207 (1.90-1.90)		
Clashscore	141614	6847 (1.90-1.90)		
Ramachandran outliers	138981	6760 (1.90-1.90)		
Sidechain outliers	138945	6760 (1.90-1.90)		
RSRZ outliers	127900	6082 (1.90-1.90)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	104	93%	7%				
1	В	104	94%	6%				
2	С	466	89%	11%				
2	D	466	6% 89%	10%				
3	Е	6	100%					



Continued from previous page...

Mol	Chain	Length	Quality of chain						
3	F	6	83%	17%					

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SCN	В	107	-	-	X	-



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 10349 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called MYELOPEROXIDASE.

\mathbf{Mol}	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	104	Total 838	C 529	N 148	O 156	S 5	0	0	0
1	В	104	Total 838	C 529	N 148	O 156	S 5	0	0	0

• Molecule 2 is a protein called MYELOPEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	С	466	Total 3733	C 2351	N 687	O 668	S 27	0	0	0
2	D	466	Total 3733	C 2351	N 687	O 668	S 27	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	150	CSO	CYS	modified residue	UNP P05164
D	150	CSO	CYS	modified residue	UNP P05164


• Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace	
3	E	6	Total 71	C 40		0	0	0
3	F	6	Total 71	C 40	O 29	0	0	0




 \bullet Molecule 4 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
1	Δ	1	Total	С	N	S	0	0	
4	4 /	1	3	1	1	1		0	
4	С	1	Total	С	N	S	0	0	
4		1	3	1	1	1			
4	В	1	Total	С	N	S	0	0	
4	Б	1	3	1	1	1	0	U	
4	D	1	Total	С	N	S	0	0	
4	В	1	3	1	1	1	0		


 \bullet Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	С	1	Total O S 5 4 1	0	0
5	В	1	Total O S 5 4 1	0	0


• Molecule 6 is CYANIDE ION (three-letter code: CYN) (formula: CN).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N 2 1 1	0	0
6	С	1	Total C N 2 1 1	0	0
6	В	1	Total C N 2 1 1	0	0
6	D	1	Total C N 2 1 1	0	0

• Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
7	C	1	Total C N O	0	0	
'		1	14 8 1 5	O		
7	C	1	Total C N O		0	
'	O	1	14 8 1 5	0		
7	D	1	Total C N O	0	0	
'	D	1	14 8 1 5	0	U	
7	D	1	Total C N O	0	0	
'	D	$D \mid I \mid$	14 8 1 5	0		

• Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	С	1	Total Ca 1 1	0	0
8	D	1	Total Ca 1 1	0	0


 \bullet Molecule 9 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $\rm C_{34}H_{32}FeN_4O_4).$





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
0	С	1	Total	С	Fe	N	О	0	0
	1	43	34	1	4	4	0	0	
0	D	1	Total	С	Fe	N	О	0	0
9 B	1	43	34	1	4	4	0	U	

 \bullet Molecule 10 is ACETIC ACID (three-letter code: ACY) (formula: $\mathrm{C_2H_4O_2}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	С	1	Total C O 4 2 2	0	0
10	С	1	Total C O 4 2 2	0	0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	С	1	Total C O 4 2 2	0	0
10	D	1	Total C O 4 2 2	0	0
10	D	1	Total C O 4 2 2	0	0
10	D	1	Total C O 4 2 2	0	0


• Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	95	Total O 95 95	0	0
11	С	338	Total O 338 338	0	0
11	В	100	Total O 100 100	0	0
11	D	329	Total O 329 329	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



Chain E:	100%

 $\bullet \ \, \text{Molecule 3: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]} \\ \text{beta-D-mannopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]} \\ 2\text{-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]} \\ 2\text{-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)} \\$

Chain F: 83% 17%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	111.26Å 63.77Å 92.60Å	Depositor
a, b, c, α , β , γ	90.00° 97.51° 90.00°	Depositor
Resolution (Å)	10.00 - 1.90	Depositor
resolution (A)	29.38 - 1.90	EDS
% Data completeness	93.4 (10.00-1.90)	Depositor
(in resolution range)	92.3 (29.38-1.90)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.74 (at 1.91Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
P.P.	0.197 , 0.224	Depositor
R, R_{free}	0.188 , 0.216	DCC
R_{free} test set	4686 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 62.9	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10349	wwPDB-VP
Average B, all atoms $(Å^2)$	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CSO, HEM, FUC, SCN, CYN, ACY, BMA, MAN, CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.42	0/863	0.74	0/1174
1	В	0.41	0/863	0.72	0/1174
2	С	0.39	0/3811	0.66	0/5168
2	D	0.41	0/3811	0.64	0/5168
All	All	0.40	0/9348	0.67	0/12684

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	838	0	798	5	0
1	В	838	0	798	6	0
2	С	3733	0	3724	32	0
2	D	3733	0	3724	32	0
3	Е	71	0	61	0	0
3	F	71	0	61	0	0
4	A	3	0	0	0	0
4	В	6	0	0	2	0
4	С	3	0	0	1	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	5	0	0	0	0
5	В	5	0	0	0	0
5	С	5	0	0	0	0
6	A	2	0	0	1	0
6	В	2	0	0	1	0
6	С	2	0	0	0	0
6	D	2	0	0	0	0
7	С	28	0	26	0	0
7	D	28	0	26	0	0
8	С	1	0	0	0	0
8	D	1	0	0	0	0
9	В	43	0	30	1	0
9	С	43	0	30	1	0
10	С	12	0	9	0	0
10	D	12	0	9	1	0
11	A	95	0	0	1	0
11	В	100	0	0	1	0
11	С	338	0	0	3	0
11	D	329	0	0	1	0
All	All	10349	0	9296	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance } (\text{\AA}) \end{array}$	Clash overlap (Å)
2:C:333:ARG:HH21	2:C:421:ASN:HD22	1.31	0.75
2:C:333:ARG:HH21	2:C:421:ASN:ND2	1.85	0.74
11:A:193:HOH:O	2:C:129:LYS:HD3	1.92	0.69
2:C:200:ASN:HD22	2:C:203:PHE:H	1.41	0.67
4:C:9:SCN:S	11:C:866:HOH:O	2.54	0.66
2:D:244:PRO:HB2	2:D:343:MET:SD	2.39	0.63
2:D:349:ARG:HG3	2:D:351:GLN:HG2	1.82	0.62
2:C:211:LEU:HD23	2:C:254:LEU:HD13	1.82	0.61
2:C:200:ASN:ND2	2:C:203:PHE:H	2.01	0.58
2:D:241:SER:O	2:D:366:PHE:HA	2.04	0.58
1:A:4:GLN:HG2	1:A:5:ASP:N	2.18	0.57
2:C:200:ASN:ND2	2:C:202:ARG:H	2.02	0.57
2:C:354:GLU:HB3	2:C:355:PRO:HA	1.87	0.56
2:C:135:PRO:HG2	11:C:905:HOH:O	2.07	0.54



 $Continued\ from\ previous\ page...$

Continuea from previ		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap(Å)
2:C:271:TRP:CZ3	2:C:279:GLU:HG3	2.44	0.53
2:D:244:PRO:HD3	2:D:364:VAL:O	2.07	0.53
2:D:113:VAL:HG21	2:D:122:GLN:HB2	1.89	0.53
1:B:83:SER:HB3	2:D:554:SER:O	2.10	0.52
1:A:83:SER:HB3	2:C:554:SER:O	2.09	0.52
2:C:244:PRO:HD3	2:C:364:VAL:O	2.08	0.52
2:D:393:ARG:HB2	2:D:396:GLN:HB2	1.91	0.52
2:C:200:ASN:HD22	2:C:202:ARG:H	1.56	0.51
2:D:333:ARG:HH21	2:D:421:ASN:ND2	2.09	0.51
4:B:107:SCN:C	2:D:242:GLU:HG3	2.41	0.50
2:C:417:LEU:HB3	2:C:418:PRO:HD3	1.92	0.50
2:D:417:LEU:HB3	2:D:418:PRO:HD3	1.94	0.50
4:B:107:SCN:N	6:B:108:CYN:N	2.60	0.50
2:D:349:ARG:HE	2:D:351:GLN:HG3	1.76	0.50
1:B:101:PRO:HD2	2:D:164:ILE:O	2.12	0.49
1:A:101:PRO:HD2	2:C:164:ILE:O	2.13	0.49
2:D:572:LEU:O	2:D:575:TRP:HB2	2.13	0.49
2:C:465:MET:HE1	2:C:471:PRO:HG3	1.94	0.49
2:C:333:ARG:HD3	2:C:421:ASN:ND2	2.28	0.49
1:A:29:PHE:CE2	2:C:165:ASN:HB2	2.47	0.48
2:D:411:MET:CE	2:D:415:LEU:HD21	2.43	0.48
6:A:107:CYN:C	9:C:14:HEM:ND	2.77	0.48
2:D:252:LEU:HD11	2:D:537:ILE:HA	1.96	0.48
9:B:109:HEM:HBB2	2:D:242:GLU:OE1	2.14	0.47
1:B:29:PHE:CZ	2:D:329:THR:HG21	2.50	0.47
2:C:128:LEU:HB2	2:C:144:ILE:HB	1.96	0.47
2:D:116:GLU:OE1	2:D:411:MET:HE3	2.14	0.47
2:D:535:ARG:NH2	2:D:567:LEU:O	2.49	0.45
1:A:11:THR:O	2:C:168:THR:HG22	2.16	0.45
2:D:394:GLN:HB3	2:D:460:LEU:HD22	1.99	0.45
2:D:523:GLN:CD	2:D:523:GLN:H	2.21	0.45
2:C:462:ARG:O	2:C:466:GLU:HG2	2.18	0.44
2:D:522:MET:HG3	10:D:17:ACY:H1	2.00	0.44
2:D:411:MET:HE2	2:D:415:LEU:HD21	1.99	0.44
2:D:378:ASP:HB2	2:D:379:PRO:HD3	2.00	0.44
2:C:378:ASP:OD1	2:C:541:THR:HB	2.18	0.44
1:B:32:TRP:CE2	2:D:325:ALA:HB2	2.53	0.43
2:C:282:LYS:HG2	2:C:520:PHE:CZ	2.54	0.43
2:D:361:LEU:O	2:D:364:VAL:HG22	2.19	0.42
2:C:526:GLN:NE2	2:C:530:GLN:HE21	2.17	0.42
2:C:548:LYS:HD2	2:C:560:ASP:HA	2.00	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å)
2:C:139:ASN:HB2	2:C:142:ASP:OD1	2.19	0.42
2:D:221:CYS:HA	2:D:366:PHE:O	2.19	0.42
2:D:568:PRO:HG2	11:D:913:HOH:O	2.19	0.42
2:C:114:ASN:HA	11:C:879:HOH:O	2.20	0.42
1:B:1:CYS:HB2	11:B:759:HOH:O	2.20	0.41
2:C:242:GLU:O	2:C:365:PHE:HA	2.20	0.41
2:C:333:ARG:NH2	2:C:421:ASN:HD22	2.08	0.41
2:D:242:GLU:O	2:D:365:PHE:HA	2.20	0.41
2:D:516:ASN:HB3	2:D:519:VAL:CG2	2.50	0.41
2:C:163:GLN:HG2	2:C:428:HIS:NE2	2.35	0.41
2:C:181:GLU:HB2	2:C:182:PRO:HD3	2.02	0.41
2:C:411:MET:HE3	2:C:415:LEU:HD21	2.02	0.41
1:B:29:PHE:CE2	2:D:165:ASN:HB2	2.56	0.41
2:D:378:ASP:OD1	2:D:541:THR:HB	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	102/104 (98%)	100 (98%)	2 (2%)	0	100	100
1	В	102/104 (98%)	100 (98%)	2 (2%)	0	100	100
2	С	463/466 (99%)	451 (97%)	11 (2%)	1 (0%)	47	38
2	D	463/466 (99%)	450 (97%)	13 (3%)	0	100	100
All	All	1130/1140 (99%)	1101 (97%)	28 (2%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	С	114	ASN



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Rotameric Outliers		Percentiles		
1	A	90/90~(100%)	89 (99%)	1 (1%)	73	73		
1	В	90/90 (100%)	89 (99%)	1 (1%)	73	73		
2	С	410/410 (100%)	404 (98%)	6 (2%)	65	62		
2	D	410/410 (100%)	403 (98%)	7 (2%)	60	57		
All	All	1000/1000 (100%)	985 (98%)	15 (2%)	65	62		

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	ASN
2	С	123	PRO
2	С	175	MET
2	С	217	HIS
2	С	254	LEU
2	С	267	LEU
2	С	523	GLN
1	В	54	ASN
2	D	138	LYS
2	D	175	MET
2	D	254	LEU
2	D	267	LEU
2	D	487	ARG
2	D	497	CYS
2	D	523	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	ASN
2	С	122	GLN
2	С	200	ASN
2	С	421	ASN



Continued from previous page...

Mol	Chain	Res	Type	
2	С	526	GLN	
1	В	54	ASN	
2	D	140	GLN	
2	D	549	ASN	

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mo	Type	Type Chain F		Link	В	${ m gths}$	Bond angles			
IVIO	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CSO	С	150	2	3,6,7	0.72	0	0,6,8	-	-
2	CSO	D	150	2	3,6,7	0.65	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

	Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
	2	CSO	С	150	2	-	0/1/5/7	_
Ī	2	CSO	D	150	2	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



No monomer is involved in short contacts.

5.5 Carbohydrates (i)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Dag	Link	Bo	ond leng	ths	Bond angles			
Mol	Type	Chain	Res	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	NAG	Е	1	2,3	14,14,15	0.57	0	17,19,21	0.87	0	
3	NAG	Е	2	3	14,14,15	0.41	0	17,19,21	0.73	0	
3	BMA	Е	3	3	11,11,12	0.36	0	15,15,17	0.45	0	
3	MAN	Е	4	3	11,11,12	0.51	0	15,15,17	0.61	0	
3	MAN	Е	5	3	11,11,12	0.44	0	15,15,17	0.56	0	
3	FUC	Е	6	3	10,10,11	0.43	0	14,14,16	0.52	0	
3	NAG	F	1	2,3	14,14,15	0.49	0	17,19,21	0.84	0	
3	NAG	F	2	3	14,14,15	0.45	0	17,19,21	0.97	2 (11%)	
3	BMA	F	3	3	11,11,12	0.46	0	15,15,17	0.58	0	
3	MAN	F	4	3	11,11,12	0.44	0	15,15,17	0.68	0	
3	MAN	F	5	3	11,11,12	0.48	0	15,15,17	0.57	0	
3	FUC	F	6	3	10,10,11	0.48	0	14,14,16	0.67	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	Ε	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	Е	2	3	-	0/6/23/26	0/1/1/1
3	BMA	Е	3	3	-	0/2/19/22	0/1/1/1
3	MAN	Ε	4	3	-	0/2/19/22	0/1/1/1
3	MAN	Е	5	3	-	0/2/19/22	0/1/1/1
3	FUC	Е	6	3	-	-	0/1/1/1
3	NAG	F	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	0/2/19/22	0/1/1/1



Continued from previous page...

Mo	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MAN	F	4	3	-	0/2/19/22	0/1/1/1
3	MAN	F	5	3	-	0/2/19/22	0/1/1/1
3	FUC	F	6	3	-	-	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	F	2	NAG	C4-C3-C2	-2.63	107.16	111.02
3	F	2	NAG	C2-N2-C7	-2.24	119.71	122.90

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 2 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	NAG	С	630	2	14,14,15	0.44	0	17,19,21	0.74	0
5	SO4	A	106	-	4,4,4	0.30	0	6,6,6	0.09	0
10	ACY	С	20	-	3,3,3	1.40	1 (33%)	3,3,3	1.59	1 (33%)
6	CYN	С	6	-	0,1,1	-	-	=		
10	ACY	С	19	-	3,3,3	1.30	1 (33%)	3,3,3	1.65	1 (33%)
10	ACY	D	18	-	3,3,3	1.27	0	3,3,3	1.67	1 (33%)



Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
6	CYN	A	107	9	0,1,1	-	-	-		
9	HEM	С	14	6,1,2	41,50,50	1.29	5 (12%)	45,82,82	1.25	3 (6%)
6	CYN	В	108	9	0,1,1	-	-	-		
9	HEM	В	109	6,1,2	41,50,50	1.35	7 (17%)	45,82,82	1.49	7 (15%)
10	ACY	С	21	-	3,3,3	1.18	0	3,3,3	1.71	1 (33%)
10	ACY	D	16	-	3,3,3	1.31	0	3,3,3	1.60	1 (33%)
4	SCN	В	107	-	1,2,2	1.77	0	0,1,1	-	-
5	SO4	В	106	-	4,4,4	0.68	0	6,6,6	0.12	0
7	NAG	С	620	2	14,14,15	0.46	0	17,19,21	0.89	1 (5%)
5	SO4	С	13	-	4,4,4	0.34	0	6,6,6	0.18	0
7	NAG	D	630	2	14,14,15	0.52	0	17,19,21	0.64	0
4	SCN	С	9	-	1,2,2	1.81	0	0,1,1	_	-
4	SCN	В	105	-	1,2,2	1.59	0	0,1,1	-	-
10	ACY	D	17	-	3,3,3	1.53	1 (33%)	3,3,3	1.74	1 (33%)
6	CYN	D	5	-	0,1,1	-	-	-		
7	NAG	D	620	2	14,14,15	0.50	0	17,19,21	0.77	0
4	SCN	A	105	-	1,2,2	1.33	0	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	D	630	2	-	2/6/23/26	0/1/1/1
7	NAG	С	630	2	-	0/6/23/26	0/1/1/1
9	HEM	В	109	6,1,2	-	4/12/54/54	-
7	NAG	С	620	2	-	0/6/23/26	0/1/1/1
7	NAG	D	620	2	-	0/6/23/26	0/1/1/1
9	HEM	С	14	6,1,2	-	4/12/54/54	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\rm Observed(\mathring{A})$	$\operatorname{Ideal}(\text{\AA})$
9	С	14	HEM	C3C-CAC	-3.82	1.40	1.47
9	В	109	HEM	C3C-CAC	-3.62	1.40	1.47
9	В	109	HEM	CBB-CAB	3.17	1.46	1.30
9	С	14	HEM	CBB-CAB	2.72	1.43	1.30
10	D	17	ACY	O-C	2.41	1.33	1.22
9	С	14	HEM	CAB-C3B	-2.35	1.41	1.47



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
9	В	109	HEM	CAB-C3B	-2.33	1.41	1.47
9	В	109	HEM	CBC-CAC	2.20	1.43	1.29
10	С	20	ACY	O-C	2.19	1.32	1.22
10	С	19	ACY	O-C	2.10	1.32	1.22
9	В	109	HEM	CHA-C4D	2.09	1.40	1.35
9	С	14	HEM	CBC-CAC	2.07	1.43	1.29
9	В	109	HEM	CHB-C1B	2.06	1.40	1.35
9	В	109	HEM	C1B-NB	-2.02	1.36	1.40
9	С	14	HEM	C4A-NA	2.02	1.40	1.36

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\mathrm{Ideal}(^{o})$
9	В	109	HEM	C4B-C3B-C2B	-4.73	103.36	107.11
9	В	109	HEM	CMC-C2C-C3C	4.02	132.20	124.68
9	С	14	HEM	CMC-C2C-C3C	3.89	131.96	124.68
9	С	14	HEM	C4A-C3A-C2A	-3.13	104.82	107.00
9	В	109	HEM	CMB-C2B-C1B	-3.09	120.33	125.04
9	В	109	HEM	C3B-C2B-C1B	2.82	108.58	106.49
7	С	620	NAG	C2-N2-C7	-2.49	119.35	122.90
9	В	109	HEM	CBA-CAA-C2A	-2.48	108.39	112.62
9	С	14	HEM	C4C-CHD-C1D	2.48	125.83	122.56
9	В	109	HEM	C4B-CHC-C1C	2.40	125.73	122.56
10	D	17	ACY	O-C-CH3	-2.39	113.04	122.33
10	С	21	ACY	O-C-CH3	-2.37	113.12	122.33
10	D	18	ACY	O-C-CH3	-2.30	113.36	122.33
10	С	19	ACY	O-C-CH3	-2.25	113.56	122.33
9	В	109	HEM	C4C-CHD-C1D	2.25	125.52	122.56
10	D	16	ACY	O-C-CH3	-2.20	113.77	122.33
10	С	20	ACY	O-C-CH3	-2.17	113.87	122.33

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	D	630	NAG	C4-C5-C6-O6
7	D	630	NAG	O5-C5-C6-O6
9	С	14	HEM	CAD-CBD-CGD-O2D
9	В	109	HEM	CAD-CBD-CGD-O1D
9	В	109	HEM	CAD-CBD-CGD-O2D
9	С	14	HEM	CAD-CBD-CGD-O1D
9	С	14	HEM	CAA-CBA-CGA-O1A



Continued from previous page...

Mol	Chain	Res	Type	Atoms
9	С	14	HEM	CAA-CBA-CGA-O2A
9	В	109	HEM	CAA-CBA-CGA-O2A
9	В	109	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.

7 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	107	CYN	1	0
9	С	14	HEM	1	0
6	В	108	CYN	1	0
9	В	109	HEM	1	0
4	В	107	SCN	2	0
4	С	9	SCN	1	0
10	D	17	ACY	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	$104/104\ (100\%)$	0.11	6 (5%) 23 25	6, 11, 23, 38	0
1	В	104/104 (100%)	0.00	0 100 100	7, 12, 21, 26	0
2	С	465/466~(99%)	0.15	15 (3%) 47 50	5, 13, 24, 33	0
2	D	465/466~(99%)	0.36	26 (5%) 24 27	6, 15, 27, 35	0
All	All	1138/1140 (99%)	0.22	47 (4%) 37 40	5, 13, 26, 38	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	С	113	VAL	9.4
2	С	355	PRO	7.1
2	D	568	PRO	6.1
2	С	217	HIS	5.3
2	D	355	PRO	5.3
2	D	578	ALA	5.2
2	D	217	HIS	5.1
1	A	3	GLU	5.0
2	D	349	ARG	4.5
2	D	113	VAL	3.6
2	D	226	ARG	3.5
1	A	4	GLN	3.5
2	D	576	ARG	3.4
1	A	1	CYS	3.2
2	С	356	ASN	3.2
2	D	190	MET	3.1
2	С	349	ARG	3.1
2	D	567	LEU	3.1
2	D	270	ARG	3.1
2	D	253	LEU	3.1
2	С	348	ASN	3.0



Continued from previous page...

Mol	Chain	Res	Type	RSRZ	
2	С	354	GLU	2.9	
2	D	356	ASN	2.9	
2	С	253	LEU	2.8	
2	D	354	GLU	2.8	
2	D	565	SER	2.8	
2	D	269	PRO	2.8	
2	С	353	MET	2.8	
2	D	563	ASN	2.7	
2	D	566	THR	2.7	
2	С	190	MET	2.6	
1	A	104	ALA	2.5	
2	D	202	ARG	2.5	
2	D	267	LEU	2.5	
2	D	157	ASN	2.5	
2	D	155	GLY	2.4	
2	С	157	ASN	2.4	
2	D	348	ASN	2.3	
2	D	138	LYS	2.2	
1	A	92	LEU	2.1	
1	A	89	TRP	2.1	
2	С	246	LEU	2.1	
2	С	218	ASP	2.1	
2	С	351	GLN	2.1	
2	D	557	TYR	2.0	
2	С	557	TYR	2.0	
2	D	476	ILE	2.0	

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	CSO	С	150	7/8	0.91	0.10	8,9,10,12	0
2	CSO	D	150	7/8	0.94	0.11	7,10,10,14	0




6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
3	MAN	F	4	11/12	0.90	0.19	20,23,25,26	0
3	MAN	E	4	11/12	0.92	0.15	21,23,24,24	0
3	MAN	F	5	11/12	0.92	0.11	14,15,16,16	0
3	NAG	E	1	14/15	0.95	0.08	9,11,16,17	0
3	MAN	Ε	5	11/12	0.95	0.09	13,14,15,15	0
3	FUC	Ε	6	10/11	0.95	0.08	13,15,16,16	0
3	NAG	F	1	14/15	0.95	0.10	11,13,18,20	0
3	NAG	Ε	2	14/15	0.95	0.09	9,10,12,12	0
3	BMA	Ε	3	11/12	0.95	0.10	12,13,15,19	0
3	FUC	F	6	10/11	0.95	0.13	14,15,16,18	0
3	BMA	F	3	11/12	0.96	0.12	12,13,14,17	0
3	NAG	F	2	14/15	0.96	0.09	9,10,13,13	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.









6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ m \AA}^2)$	Q<0.9
6	CYN	D	5	2/2	0.71	0.27	11,11,11,14	0
10	ACY	D	17	4/4	0.73	0.38	33,36,36,38	0
5	SO4	В	106	5/5	0.81	0.48	55,55,56,57	0
7	NAG	D	630	14/15	0.84	0.24	26,29,30,33	0
10	ACY	С	19	4/4	0.86	0.13	28,29,29,29	0
7	NAG	С	630	14/15	0.89	0.22	22,25,31,31	0
7	NAG	D	620	14/15	0.90	0.16	17,21,25,26	0
6	CYN	С	6	2/2	0.90	0.10	21,21,21,24	0
10	ACY	D	18	4/4	0.91	0.13	25,25,25,28	0




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
7	NAG	С	620	14/15	0.91	0.17	14,19,22,23	0
10	ACY	С	21	4/4	0.92	0.15	26,26,27,27	0
10	ACY	С	20	4/4	0.92	0.17	26,26,26,26	0
10	ACY	D	16	4/4	0.92	0.18	29,29,29,31	0
5	SO4	С	13	5/5	0.94	0.13	27,27,29,29	0
6	CYN	A	107	2/2	0.95	0.17	12,12,12,15	0
6	CYN	В	108	2/2	0.96	0.15	18,18,18,18	0
4	SCN	С	9	3/3	0.96	0.11	19,19,20,21	0
9	HEM	В	109	43/43	0.96	0.12	6,10,14,14	0
4	SCN	В	107	3/3	0.97	0.09	26,26,26,28	0
9	HEM	С	14	43/43	0.97	0.12	4,9,10,12	0
4	SCN	A	105	3/3	0.97	0.09	3,3,4,12	0
8	CA	D	2	1/1	0.99	0.06	8,8,8,8	0
4	SCN	В	105	3/3	0.99	0.12	5,5,5,10	0
5	SO4	A	106	5/5	0.99	0.09	25,26,27,28	0
8	CA	С	1	1/1	1.00	0.05	7,7,7,7	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









6.5 Other polymers (i)

There are no such residues in this entry.

