



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 05:27 PM EDT

PDB ID : 1ETH
Title : TRIACYLGLYCEROL LIPASE/COLIPASE COMPLEX
Authors : Hermoso, J.; Pignol, D.; Kerfelec, B.; Crenon, I.; Chapus, C.; Fontecilla-Camps, J.C.
Deposited on : 1995-09-13
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	448	31% 52% 15% .
1	C	448	30% 52% 17% .
2	B	95	36% 41% 15% 8%
2	D	95	34% 44% 14% 8%
3	E	5	40% 60%
3	F	5	20% 80%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	E	1	-	-	X	-
3	NAG	E	2	-	-	X	-
3	BMA	E	4	X	-	-	-
3	BMA	E	5	X	-	-	-
3	BMA	F	4	X	-	-	-
3	BMA	F	5	X	-	-	-
6	BME	C	457	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11761 atoms, of which 2850 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIACYLGLYCEROL ACYL-HYDROLASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	448	4301	2219	788	606	670	18	0	0	0
1	C	448	4301	2219	788	606	670	18	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASN	deletion	UNP P00591
A	?	-	ASN	deletion	UNP P00591
C	?	-	ASN	deletion	UNP P00591
C	?	-	ASN	deletion	UNP P00591

- Molecule 2 is a protein called COLIPASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	87	811	400	155	116	130	10	0	0	0
2	D	87	811	400	155	116	130	10	0	0	0

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	E	5	118	34	57	2	25	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
3	F	5	118	34	57	2	25	0	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	C	1	Total	Ca	0	0
			1	1		

- Molecule 5 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula: C₁₆H₃₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	0	0
			55	16	34	5		
5	A	1	Total	C	H	O	0	0
			55	16	34	5		
5	C	1	Total	C	H	O	0	0
			55	16	34	5		
5	C	1	Total	C	H	O	0	0
			55	16	34	5		

- Molecule 6 is BETA-MERCAPTOETHANOL (three-letter code: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			4	2	1	1		
6	C	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	128	Total	H	O	0	0
			384	256	128		
7	B	31	Total	H	O	0	0
			93	62	31		
7	C	166	Total	H	O	0	0
			498	332	166		
7	D	32	Total	H	O	0	0
			96	64	32		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.


- Molecule 1: TRIACYLGLYCEROL ACYL-HYDROLASE


Chain A: 



- Molecule 1: TRIACYLGLYCEROL ACYL-HYDROLASE

Chain C: 






- Molecule 2: COLIPASE



- Molecule 2: COLIPASE



- Molecule 3: beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	F 2 3	Depositor
Cell constants a, b, c, α , β , γ	289.10Å 289.10Å 289.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.80)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.210 , 0.290	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	11761	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CA, C8E, BME, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3602	0.71	1/4892 (0.0%)
1	C	0.55	0/3602	0.70	0/4892
2	B	0.56	0/664	0.72	0/894
2	D	0.56	0/664	0.73	0/894
All	All	0.55	0/8532	0.71	1/11572 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	17
1	C	0	18
All	All	0	35

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	123	ARG	NE-CZ-NH2	-5.07	117.76	120.30

There are no chirality outliers.

5 of 35 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11	PHE	Sidechain
1	A	115	TYR	Sidechain
1	A	2	GLU	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	208	ALA	Mainchain
1	A	42	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3513	788	3370	304	0
1	C	3513	788	3370	319	0
2	B	656	155	631	56	0
2	D	656	155	631	57	0
3	E	61	57	51	12	0
3	F	61	57	51	9	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	42	68	50	1	0
5	C	42	68	49	1	0
6	A	4	0	6	2	0
6	C	4	0	6	5	0
7	A	128	256	0	20	0
7	B	31	62	0	5	0
7	C	166	332	0	30	0
7	D	32	64	0	7	0
All	All	8911	2850	8215	738	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

The worst 5 of 738 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:182:CYS:SG	6:C:457:BME:S2	2.35	1.13
1:A:182:CYS:SG	6:A:457:BME:S2	2.39	1.05
1:C:108:LYS:HB2	1:C:112:ARG:HH12	1.23	1.00
1:A:108:LYS:HB2	1:A:112:ARG:HH12	1.25	1.00
1:C:356:ILE:HD11	1:C:373:TYR:HB3	1.43	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	446/448 (100%)	361 (81%)	66 (15%)	19 (4%)	2	8
1	C	446/448 (100%)	363 (81%)	63 (14%)	20 (4%)	2	8
2	B	85/95 (90%)	75 (88%)	8 (9%)	2 (2%)	6	20
2	D	85/95 (90%)	76 (89%)	7 (8%)	2 (2%)	6	20
All	All	1062/1086 (98%)	875 (82%)	144 (14%)	43 (4%)	3	9

5 of 43 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	6	PRO
1	A	23	ARG
1	A	186	THR
1	C	23	ARG
1	C	186	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/388 (100%)	311 (80%)	77 (20%)	1	4
1	C	388/388 (100%)	311 (80%)	77 (20%)	1	4
2	B	76/83 (92%)	62 (82%)	14 (18%)	1	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	76/83 (92%)	62 (82%)	14 (18%)	1	5
All	All	928/942 (98%)	746 (80%)	182 (20%)	1	4

5 of 182 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	165	ARG
1	C	350	LYS
1	C	193	ASP
1	C	249	ILE
1	C	367	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	46	ASN
1	C	93	ASN
2	D	81	ASN
1	C	64	ASN
1	C	117	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	E	1	1,3	14,14,15	3.18	7 (50%)	17,19,21	4.84	11 (64%)
3	NAG	E	2	3	14,14,15	3.30	8 (57%)	17,19,21	3.41	10 (58%)
3	BMA	E	3	3	11,11,12	2.40	3 (27%)	15,15,17	2.84	8 (53%)
3	BMA	E	4	3	11,11,12	2.30	4 (36%)	15,15,17	3.69	9 (60%)
3	BMA	E	5	3	11,11,12	3.35	5 (45%)	15,15,17	2.35	7 (46%)
3	NAG	F	1	1,3	14,14,15	1.72	2 (14%)	17,19,21	3.42	8 (47%)
3	NAG	F	2	3	14,14,15	2.81	5 (35%)	17,19,21	4.54	12 (70%)
3	BMA	F	3	3	11,11,12	2.82	6 (54%)	15,15,17	3.22	6 (40%)
3	BMA	F	4	3	11,11,12	2.50	6 (54%)	15,15,17	4.30	11 (73%)
3	BMA	F	5	3	11,11,12	2.75	5 (45%)	15,15,17	2.18	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	E	1	1,3	-	3/6/23/26	0/1/1/1
3	NAG	E	2	3	-	3/6/23/26	0/1/1/1
3	BMA	E	3	3	-	2/2/19/22	0/1/1/1
3	BMA	E	4	3	1/1/5/5	2/2/19/22	0/1/1/1
3	BMA	E	5	3	1/1/5/5	2/2/19/22	1/1/1/1
3	NAG	F	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1
3	BMA	F	3	3	-	2/2/19/22	0/1/1/1
3	BMA	F	4	3	1/1/5/5	0/2/19/22	0/1/1/1
3	BMA	F	5	3	1/1/5/5	0/2/19/22	1/1/1/1

The worst 5 of 51 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	C4-C3	8.03	1.72	1.52
3	E	5	BMA	C2-C3	7.03	1.62	1.52
3	E	1	NAG	O5-C5	6.12	1.55	1.43
3	E	3	BMA	C2-C3	5.97	1.61	1.52
3	E	5	BMA	C4-C5	5.96	1.65	1.53

The worst 5 of 86 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	2	NAG	C6-C5-C4	12.16	141.49	113.00
3	E	1	NAG	C2-N2-C7	-11.36	106.73	122.90
3	E	4	BMA	C6-C5-C4	9.61	135.52	113.00
3	F	4	BMA	C2-C3-C4	-7.74	97.50	110.89
3	F	2	NAG	C3-C4-C5	-7.65	96.60	110.24

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	E	4	BMA	C1
3	E	5	BMA	C1
3	F	4	BMA	C1
3	F	5	BMA	C1

5 of 16 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	2	NAG	C3-C2-N2-C7
3	E	4	BMA	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6
3	E	4	BMA	O5-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6


All (2) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	F	5	BMA	C1-C2-C3-C4-C5-O5
3	E	5	BMA	C1-C2-C3-C4-C5-O5

7 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	1	NAG	9	0
3	F	4	BMA	2	0
3	F	1	NAG	5	0
3	E	4	BMA	2	0
3	F	2	NAG	6	0
3	E	2	NAG	7	0
3	F	3	BMA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	C8E	C	456	-	20,20,20	1.27	1 (5%)	19,19,19	1.07	1 (5%)
6	BME	A	457	-	3,3,3	0.33	0	1,2,2	0.16	0
5	C8E	C	455	-	20,20,20	0.96	1 (5%)	19,19,19	0.98	1 (5%)
5	C8E	A	456	-	20,20,20	1.13	1 (5%)	19,19,19	1.03	1 (5%)
6	BME	C	457	-	3,3,3	0.60	0	1,2,2	0.28	0
5	C8E	A	455	-	20,20,20	1.06	1 (5%)	19,19,19	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C8E	C	456	-	-	6/18/18/18	-
6	BME	A	457	-	-	1/1/1/1	-
5	C8E	C	455	-	-	5/18/18/18	-
5	C8E	A	456	-	-	7/18/18/18	-
6	BME	C	457	-	-	0/1/1/1	-
5	C8E	A	455	-	-	9/18/18/18	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	456	C8E	O21-C20	-2.36	1.29	1.42
5	A	455	C8E	O21-C20	-2.25	1.30	1.42
5	C	456	C8E	O21-C20	-2.21	1.30	1.42
5	C	455	C8E	O21-C20	-2.19	1.30	1.42

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	456	C8E	O21-C20-C19	2.30	125.18	111.81
5	C	456	C8E	C19-O18-C17	2.20	122.80	113.29
5	C	455	C8E	O18-C17-C16	-2.08	101.00	110.39

There are no chirality outliers.

5 of 28 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	456	C8E	C3-C4-C5-C6
5	C	456	C8E	C2-C3-C4-C5
5	C	456	C8E	C6-C7-C8-O9
5	A	456	C8E	C3-C4-C5-C6
5	A	455	C8E	O15-C16-C17-O18


There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	456	C8E	1	0
6	A	457	BME	2	0
6	C	457	BME	5	0
5	A	455	C8E	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.