



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 7, 2023 – 05:16 am GMT

PDB ID : 1GZG
Title : Complex of a Mg²-dependent porphobilinogen synthase from *Pseudomonas aeruginosa* (mutant D139N) with 5-fluorolevulinic acid
Authors : Frere, F.; Schubert, W.-D.; Stauffer, F.; Frankenberg, N.; Neier, R.; Jahn, D.; Heinz, D.W.
Deposited on : 2002-05-21
Resolution : 1.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	337	
1	B	337	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	LAF	B	1337	-	-	X	-
2	LAF	B	1338[B]	-	-	X	-
2	LAF	B	1338[C]	-	-	X	-
5	SO4	B	1343	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DELTA-AMINOLEVULINIC ACID DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	329	2631	1652	469	499	11	0	11	0
1	B	323	2743	1720	490	520	13	0	27	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	139	ASN	ASP	engineered mutation	UNP Q59643
B	139	ASN	ASP	engineered mutation	UNP Q59643
A	199	VAL	ILE	SEE REMARK 999	UNP Q59643
B	199	VAL	ILE	SEE REMARK 999	UNP Q59643

- Molecule 2 is 5-FLUOROLEVULINIC ACID (three-letter code: LAF) (formula: C₅H₇FO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	F	O	0	0
			8	5	1	2		
2	A	1	Total	C	F	O	0	0
			8	5	1	2		
2	B	1	Total	C	F	O	0	1
			26	15	3	8		
2	B	1	Total	C	F	O	0	0
			8	5	1	2		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		
4	B	1	Total	Na	0	0
			1	1		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	2	Total K 2 2	0	0

- Molecule 7 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	354	Total O 354 354	0	0
7	B	317	Total O 317 317	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



- Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.13Å 127.13Å 86.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.29 – 1.66	Depositor
% Data completeness (in resolution range)	99.1 (91.29-1.66)	Depositor
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC 5.0.32	Depositor
R, R_{free}	0.175 , 0.198	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	6111	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, LAF, NA, SO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.48	1/2682 (0.0%)	0.85	8/3637 (0.2%)
1	B	0.46	0/2793	0.95	15/3784 (0.4%)
All	All	0.47	1/5475 (0.0%)	0.90	23/7421 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	260	LYS	C-N	5.15	1.44	1.34

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131	ASP	CB-CG-OD2	6.97	124.58	118.30
1	B	186	ARG	NE-CZ-NH2	-6.32	117.14	120.30
1	B	204	ALA	C-N-CA	6.06	136.86	121.70
1	B	142	LEU	CA-CB-CG	6.02	129.15	115.30
1	A	260	LYS	CA-C-O	-5.62	108.29	120.10

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	205[B]	LYS	Mainchain
1	B	94	LYS	Peptide,Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2631	0	2598	32	0
1	B	2743	0	2701	134	0
2	A	16	0	12	1	0
2	B	34	0	23	24	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	5	0	0	0	0
5	B	5	0	0	3	0
6	B	2	0	0	0	0
7	A	354	0	0	7	0
7	B	317	0	0	19	0
All	All	6111	0	5334	172	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 172 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:ASP:O	1:B:159[C]:ARG:CZ	1.85	1.24
2:B:1338[B]:LAF:C1	7:B:2316:HOH:O	1.90	1.18
1:B:155:ASP:HB3	1:B:159[C]:ARG:NH2	1.56	1.17
1:B:205[B]:LYS:NZ	2:B:1338[B]:LAF:H3C2	1.61	1.15
2:B:1338[C]:LAF:O1	7:B:2316:HOH:O	1.62	1.15

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/337 (100%)	328 (97%)	10 (3%)	0	100	100
1	B	347/337 (103%)	321 (92%)	18 (5%)	8 (2%)	6	0
All	All	685/674 (102%)	649 (95%)	28 (4%)	8 (1%)	19	2

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	142	LEU
1	B	95[A]	SER
1	B	95[B]	SER
1	B	97[A]	ASP
1	B	97[B]	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/269 (102%)	267 (97%)	7 (3%)	46	21
1	B	286/269 (106%)	277 (97%)	9 (3%)	40	14
All	All	560/538 (104%)	544 (97%)	16 (3%)	44	16

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	283	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	256[B]	MET
1	B	92[B]	GLU
1	B	256[A]	MET
1	B	92[A]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	138	GLN
1	B	298	ASN
1	B	332	GLN
1	A	138	GLN
1	A	136	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 6 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LAF	B	1338[B]	4	7,8,8	2.23	1 (14%)	7,9,9	2.65	3 (42%)
2	LAF	B	1337	1	7,7,8	0.92	0	6,7,9	2.18	4 (66%)
2	LAF	B	1338[C]	-	7,7,8	0.91	0	6,7,9	1.09	0
5	SO4	B	1343	-	4,4,4	0.17	0	6,6,6	0.12	0
5	SO4	A	1340	-	4,4,4	0.16	0	6,6,6	0.08	0
2	LAF	B	1338[A]	4	7,8,8	2.23	1 (14%)	7,9,9	2.75	3 (42%)
2	LAF	A	1336	4,1	7,7,8	1.00	0	6,7,9	1.15	1 (16%)
2	LAF	A	1337	1	7,7,8	0.88	0	6,7,9	1.59	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LAF	B	1338[B]	4	-	4/5/7/7	-
2	LAF	B	1337	1	-	3/5/5/7	-
2	LAF	B	1338[C]	-	-	4/5/5/7	-
2	LAF	B	1338[A]	4	-	3/5/7/7	-
2	LAF	A	1336	4,1	-	2/5/5/7	-
2	LAF	A	1337	1	-	1/5/5/7	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1338[A]	LAF	O4-C4	-5.25	1.12	1.21
2	B	1338[B]	LAF	O4-C4	-5.25	1.12	1.21

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1338[A]	LAF	O4-C4-C5	5.11	130.50	121.67
2	B	1338[B]	LAF	O4-C4-C5	5.01	130.34	121.67
2	B	1338[A]	LAF	O4-C4-C3	-4.20	110.32	121.44
2	B	1338[B]	LAF	O4-C4-C3	-4.02	110.82	121.44
2	B	1337	LAF	O1-C1-C2	-3.20	112.80	123.08

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1337	LAF	C3-C4-C5-F
2	A	1336	LAF	C2-C3-C4-C5
2	B	1338[C]	LAF	C2-C3-C4-C5
2	B	1338[A]	LAF	C1-C2-C3-C4
2	A	1337	LAF	C2-C3-C4-C5

There are no ring outliers.

6 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1338[B]	LAF	14	0
2	B	1337	LAF	5	0
2	B	1338[C]	LAF	7	0
5	B	1343	SO4	3	0
2	B	1338[A]	LAF	3	0
2	A	1336	LAF	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.