



wwPDB X-ray Structure Validation Summary Report

Mar 23, 2024 – 04:05 PM EDT

PDB ID : 1MT5
Title : CRYSTAL STRUCTURE OF FATTY ACID AMIDE HYDROLASE
Authors : Bracey, M.H.; Hanson, M.A.; Masuda, K.R.; Stevens, R.C.; Cravatt, B.F.
Deposited on : 2002-09-20
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	537	76% 22% .
1	B	537	76% 20% .
1	C	537	76% 19% .
1	D	537	77% 19% .
1	E	537	75% 21% .
1	F	537	75% 20% 5%
1	G	537	76% 21% . .
1	H	537	77% 20% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	I	537	 77% 21% .
1	J	537	 75% 22% .
1	K	537	 74% 23% .
1	L	537	 75% 22% .
1	M	537	 69% 28% .
1	N	537	 74% 22% .
1	O	537	 75% 21% .
1	P	537	 73% 24% .

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 64524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fatty-acid amide hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	537	4010	2568	669	744	29	0	0	0
1	B	537	4008	2568	669	742	29	0	0	0
1	C	537	4008	2568	669	742	29	0	0	0
1	D	537	4008	2568	669	742	29	0	0	0
1	E	537	4008	2568	669	742	29	0	0	0
1	F	537	4008	2568	669	742	29	0	0	0
1	G	537	4010	2568	669	744	29	0	0	0
1	H	537	4008	2568	669	742	29	0	0	0
1	I	537	4010	2568	669	744	29	0	0	0
1	J	537	4008	2568	669	742	29	0	0	0
1	K	537	4008	2568	669	742	29	0	0	0
1	L	537	4010	2568	669	744	29	0	0	0
1	M	537	4010	2568	669	744	29	0	0	0
1	N	537	4008	2568	669	742	29	0	0	0
1	O	537	4010	2568	669	744	29	0	0	0
1	P	537	4008	2568	669	742	29	0	0	0

- Molecule 2 is METHYL ARACHIDONYL FLUOROPHOSPHONATE (three-letter code: MAY) (formula: C₂₁H₃₆FO₂P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	24	21	2	1	0	0
2	B	1	24	21	2	1	0	0
2	C	1	24	21	2	1	0	0
2	D	1	24	21	2	1	0	0
2	E	1	24	21	2	1	0	0
2	F	1	24	21	2	1	0	0
2	G	1	24	21	2	1	0	0
2	H	1	24	21	2	1	0	0
2	I	1	24	21	2	1	0	0
2	J	1	24	21	2	1	0	0
2	K	1	24	21	2	1	0	0
2	L	1	24	21	2	1	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	M	1	Total 24	C 21	O 2	P 1	0	0
2	N	1	Total 24	C 21	O 2	P 1	0	0
2	O	1	Total 24	C 21	O 2	P 1	0	0
2	P	1	Total 24	C 21	O 2	P 1	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: Fatty-acid amide hydrolase




- Molecule 1: Fatty-acid amide hydrolase




- Molecule 1: Fatty-acid amide hydrolase






- Molecule 1: Fatty-acid amide hydrolase

Chain D: 77% 19%




- Molecule 1: Fatty-acid amide hydrolase

Chain E: 75% 21%




- Molecule 1: Fatty-acid amide hydrolase

Chain F: 75% 20% 5%




- Molecule 1: Fatty-acid amide hydrolase

Chain G: 76% 21%




- Molecule 1: Fatty-acid amide hydrolase

Chain H: 77% 20%




- Molecule 1: Fatty-acid amide hydrolase

Chain I: 77% 21%




- Molecule 1: Fatty-acid amide hydrolase

Chain J: 75% 22%




- Molecule 1: Fatty-acid amide hydrolase

Chain K: 74% 23%




- Molecule 1: Fatty-acid amide hydrolase

Chain L:  75% 22%




• Molecule 1: Fatty-acid amide hydrolase

Chain M:  69% 28%




• Molecule 1: Fatty-acid amide hydrolase


Chain N:  74% 22%



L562
R563
T573


● Molecule 1: Fatty-acid amide hydrolase

Chain O:  75% 21%



● Molecule 1: Fatty-acid amide hydrolase

Chain P:  73% 24%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	147.11Å 272.02Å 147.22Å 90.00° 115.21° 90.00°	Depositor
Resolution (Å)	141.42 – 2.80	Depositor
% Data completeness (in resolution range)	71.4 (141.42-2.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.218 , 0.262	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	64524	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MAY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.95	2/4102 (0.0%)	0.88	16/5589 (0.3%)
1	B	0.97	2/4100 (0.0%)	0.89	14/5586 (0.3%)
1	C	0.95	2/4100 (0.0%)	0.88	16/5586 (0.3%)
1	D	0.93	2/4100 (0.0%)	0.87	17/5586 (0.3%)
1	E	1.01	3/4100 (0.1%)	0.88	16/5586 (0.3%)
1	F	0.97	3/4100 (0.1%)	0.89	16/5586 (0.3%)
1	G	0.97	2/4102 (0.0%)	0.88	16/5589 (0.3%)
1	H	0.93	0/4100	0.88	14/5586 (0.3%)
1	I	0.96	2/4102 (0.0%)	0.86	14/5589 (0.3%)
1	J	0.97	2/4100 (0.0%)	0.87	15/5586 (0.3%)
1	K	0.98	3/4100 (0.1%)	0.89	16/5586 (0.3%)
1	L	1.02	2/4102 (0.0%)	0.88	16/5589 (0.3%)
1	M	1.13	2/4102 (0.0%)	0.93	15/5589 (0.3%)
1	N	1.09	2/4100 (0.0%)	0.89	14/5586 (0.3%)
1	O	0.96	1/4102 (0.0%)	0.88	17/5589 (0.3%)
1	P	1.00	3/4100 (0.1%)	0.88	16/5586 (0.3%)
All	All	0.99	33/65612 (0.1%)	0.88	248/89394 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	3
1	C	0	2
1	D	0	4
1	E	0	4
1	F	0	6
1	G	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	6
1	I	0	5
1	J	0	4
1	K	0	3
1	L	0	1
1	M	0	4
1	N	0	4
1	O	0	4
1	P	0	2
All	All	0	57

The worst 5 of 33 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	469	VAL	CB-CG1	-6.53	1.39	1.52
1	A	549	VAL	CA-CB	6.08	1.67	1.54
1	A	287	VAL	CA-CB	5.88	1.67	1.54
1	D	507	VAL	CA-CB	5.77	1.66	1.54
1	F	83	VAL	CA-CB	5.72	1.66	1.54

The worst 5 of 248 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	237	ASP	CB-CG-OD2	8.76	126.18	118.30
1	H	333	ASP	CB-CG-OD2	8.51	125.96	118.30
1	A	529	ASP	CB-CG-OD2	8.42	125.88	118.30
1	D	333	ASP	CB-CG-OD2	8.38	125.84	118.30
1	B	529	ASP	CB-CG-OD2	8.37	125.83	118.30

There are no chirality outliers.

5 of 57 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	120	ASP	Peptide
1	A	197	SER	Peptide
1	B	120	ASP	Peptide
1	B	40	ALA	Peptide
1	B	475	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4010	0	3953	29	0
1	B	4008	0	3953	28	0
1	C	4008	0	3953	27	0
1	D	4008	0	3953	30	0
1	E	4008	0	3953	29	0
1	F	4008	0	3953	29	0
1	G	4010	0	3953	29	0
1	H	4008	0	3953	25	0
1	I	4010	0	3953	21	0
1	J	4008	0	3953	27	0
1	K	4008	0	3953	29	0
1	L	4010	0	3953	24	0
1	M	4010	0	3953	30	0
1	N	4008	0	3953	37	0
1	O	4010	0	3953	26	0
1	P	4008	0	3953	29	0
2	A	24	0	36	5	0
2	B	24	0	36	3	0
2	C	24	0	36	3	0
2	D	24	0	36	4	0
2	E	24	0	36	3	0
2	F	24	0	36	4	0
2	G	24	0	36	1	0
2	H	24	0	36	4	0
2	I	24	0	36	2	0
2	J	24	0	36	2	0
2	K	24	0	36	3	0
2	L	24	0	36	1	0
2	M	24	0	36	2	0
2	N	24	0	36	4	0
2	O	24	0	36	3	0
2	P	24	0	36	2	0
All	All	64524	0	63824	447	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 447 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:453:MET:SD	1:L:453:MET:CE	2.02	1.48
1:D:325:LEU:H	1:D:358:HIS:HD2	1.18	0.90
1:A:155:GLY:HA3	1:A:198:ASN:HD21	1.38	0.87
1:C:325:LEU:H	1:C:358:HIS:HD2	1.27	0.81
1:D:86:LEU:HG	1:D:136:GLY:HA3	1.64	0.80

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	535/537 (100%)	498 (93%)	31 (6%)	6 (1%)	14	41
1	B	535/537 (100%)	503 (94%)	24 (4%)	8 (2%)	10	33
1	C	535/537 (100%)	507 (95%)	23 (4%)	5 (1%)	17	46
1	D	535/537 (100%)	499 (93%)	31 (6%)	5 (1%)	17	46
1	E	535/537 (100%)	491 (92%)	36 (7%)	8 (2%)	10	33
1	F	535/537 (100%)	502 (94%)	27 (5%)	6 (1%)	14	41
1	G	535/537 (100%)	506 (95%)	26 (5%)	3 (1%)	25	56
1	H	535/537 (100%)	498 (93%)	31 (6%)	6 (1%)	14	41
1	I	535/537 (100%)	496 (93%)	37 (7%)	2 (0%)	34	66
1	J	535/537 (100%)	506 (95%)	26 (5%)	3 (1%)	25	56
1	K	535/537 (100%)	497 (93%)	32 (6%)	6 (1%)	14	41
1	L	535/537 (100%)	494 (92%)	36 (7%)	5 (1%)	17	46
1	M	535/537 (100%)	466 (87%)	57 (11%)	12 (2%)	6	22
1	N	535/537 (100%)	494 (92%)	33 (6%)	8 (2%)	10	33
1	O	535/537 (100%)	499 (93%)	30 (6%)	6 (1%)	14	41

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	535/537 (100%)	497 (93%)	34 (6%)	4 (1%)	22	53
All	All	8560/8592 (100%)	7953 (93%)	514 (6%)	93 (1%)	14	41

5 of 93 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	ASP
1	A	480	ASP
1	E	189	GLN
1	E	480	ASP
1	E	488	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/455 (94%)	354 (83%)	72 (17%)	2	6
1	B	425/455 (93%)	351 (83%)	74 (17%)	2	6
1	C	425/455 (93%)	341 (80%)	84 (20%)	1	4
1	D	425/455 (93%)	354 (83%)	71 (17%)	2	6
1	E	425/455 (93%)	345 (81%)	80 (19%)	1	5
1	F	425/455 (93%)	339 (80%)	86 (20%)	1	4
1	G	426/455 (94%)	345 (81%)	81 (19%)	1	4
1	H	425/455 (93%)	350 (82%)	75 (18%)	2	5
1	I	426/455 (94%)	344 (81%)	82 (19%)	1	4
1	J	425/455 (93%)	344 (81%)	81 (19%)	1	4
1	K	425/455 (93%)	341 (80%)	84 (20%)	1	4
1	L	426/455 (94%)	338 (79%)	88 (21%)	1	3
1	M	426/455 (94%)	323 (76%)	103 (24%)	0	2
1	N	425/455 (93%)	345 (81%)	80 (19%)	1	5
1	O	426/455 (94%)	340 (80%)	86 (20%)	1	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	P	425/455 (93%)	339 (80%)	86 (20%)	1 4
All	All	6806/7280 (94%)	5493 (81%)	1313 (19%)	1 4

5 of 1313 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	387	SER
1	O	86	LEU
1	L	547	VAL
1	L	383	ASP
1	M	444	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 122 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	273	GLN
1	O	273	GLN
1	J	189	GLN
1	O	259	ASN
1	P	358	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MAY	P	600	1	20,23,24	0.39	0	18,23,26	0.63	1 (5%)
2	MAY	F	600	1	20,23,24	0.43	0	18,23,26	0.53	0
2	MAY	J	600	1	20,23,24	0.45	0	18,23,26	0.68	1 (5%)
2	MAY	O	600	1	20,23,24	0.45	0	18,23,26	0.62	1 (5%)
2	MAY	G	600	1	20,23,24	0.34	0	18,23,26	0.77	1 (5%)
2	MAY	A	600	1	20,23,24	0.38	0	18,23,26	0.77	1 (5%)
2	MAY	E	600	1	20,23,24	0.43	0	18,23,26	0.72	1 (5%)
2	MAY	M	600	1	20,23,24	0.34	0	18,23,26	0.72	1 (5%)
2	MAY	B	600	1	20,23,24	0.45	0	18,23,26	0.69	1 (5%)
2	MAY	L	600	1	20,23,24	0.43	0	18,23,26	0.77	1 (5%)
2	MAY	K	600	1	20,23,24	0.43	0	18,23,26	0.64	1 (5%)
2	MAY	N	600	1	20,23,24	0.44	0	18,23,26	0.66	1 (5%)
2	MAY	H	600	1	20,23,24	0.43	0	18,23,26	0.73	1 (5%)
2	MAY	C	600	1	20,23,24	0.33	0	18,23,26	0.70	1 (5%)
2	MAY	I	600	1	20,23,24	0.33	0	18,23,26	0.71	1 (5%)
2	MAY	D	600	1	20,23,24	0.36	0	18,23,26	0.78	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAY	P	600	1	-	7/18/22/24	-
2	MAY	F	600	1	-	8/18/22/24	-
2	MAY	J	600	1	-	7/18/22/24	-
2	MAY	O	600	1	-	7/18/22/24	-
2	MAY	G	600	1	-	7/18/22/24	-
2	MAY	A	600	1	-	7/18/22/24	-
2	MAY	E	600	1	-	7/18/22/24	-
2	MAY	M	600	1	-	7/18/22/24	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAY	B	600	1	-	9/18/22/24	-
2	MAY	L	600	1	-	9/18/22/24	-
2	MAY	K	600	1	-	8/18/22/24	-
2	MAY	N	600	1	-	8/18/22/24	-
2	MAY	H	600	1	-	8/18/22/24	-
2	MAY	C	600	1	-	8/18/22/24	-
2	MAY	I	600	1	-	7/18/22/24	-
2	MAY	D	600	1	-	9/18/22/24	-

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	MAY	C3-C2-C1	-2.74	107.04	113.85
2	D	600	MAY	C3-C2-C1	-2.67	107.22	113.85
2	E	600	MAY	C3-C2-C1	-2.60	107.41	113.85
2	L	600	MAY	C3-C2-C1	-2.57	107.47	113.85
2	G	600	MAY	C3-C2-C1	-2.52	107.59	113.85

There are no chirality outliers.

5 of 123 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	600	MAY	P1-C1-C2-C3
2	K	600	MAY	P1-C1-C2-C3
2	M	600	MAY	P1-C1-C2-C3
2	O	600	MAY	C1-C2-C3-C4
2	D	600	MAY	C1-C2-C3-C4

There are no ring outliers.

16 monomers are involved in 46 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	P	600	MAY	2	0
2	F	600	MAY	4	0
2	J	600	MAY	2	0
2	O	600	MAY	3	0
2	G	600	MAY	1	0
2	A	600	MAY	5	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	600	MAY	3	0
2	M	600	MAY	2	0
2	B	600	MAY	3	0
2	L	600	MAY	1	0
2	K	600	MAY	3	0
2	N	600	MAY	4	0
2	H	600	MAY	4	0
2	C	600	MAY	3	0
2	I	600	MAY	2	0
2	D	600	MAY	4	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












































5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.