



Full wwPDB NMR Structure Validation Report ⓘ

Feb 27, 2022 – 09:43 PM EST

PDB ID : 2DNK
Title : Solution structure of RNA binding domain in Bruno-like 4 RNA binding protein
Authors : Tsuda, K.; Muto, Y.; Inoue, M.; Kigawa, T.; Terada, T.; Shirouzu, M.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on : 2006-04-26

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.27
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.27


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	105	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 15 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:152-A:183, A:190-A:228 (71)	0.20	15

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 5, 8, 10, 12, 13, 15, 16, 18
2	4, 7, 9, 11, 14, 17, 19
3	2, 6
Single-model clusters	20

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1514 atoms, of which 743 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Bruno-like 4, RNA binding protein.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
1	A	105	1514	469	743	141	156	5	0

There are 13 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	136	GLY	-	cloning artifact	UNP Q9BQ96
A	137	SER	-	cloning artifact	UNP Q9BQ96
A	138	SER	-	cloning artifact	UNP Q9BQ96
A	139	GLY	-	cloning artifact	UNP Q9BQ96
A	140	SER	-	cloning artifact	UNP Q9BQ96
A	141	SER	-	cloning artifact	UNP Q9BQ96
A	142	GLY	-	cloning artifact	UNP Q9BQ96
A	235	SER	-	cloning artifact	UNP Q9BQ96
A	236	GLY	-	cloning artifact	UNP Q9BQ96
A	237	PRO	-	cloning artifact	UNP Q9BQ96
A	238	SER	-	cloning artifact	UNP Q9BQ96
A	239	SER	-	cloning artifact	UNP Q9BQ96
A	240	GLY	-	cloning artifact	UNP Q9BQ96

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Bruno-like 4, RNA binding protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: Bruno-like 4, RNA binding protein



4.2.2 Score per residue for model 2

- Molecule 1: Bruno-like 4, RNA binding protein






4.2.3 Score per residue for model 3

- Molecule 1: Bruno-like 4, RNA binding protein


Chain A: 42% 23% 32%



4.2.4 Score per residue for model 4

- Molecule 1: Bruno-like 4, RNA binding protein


Chain A: 45% 21% 32%



4.2.5 Score per residue for model 5


- Molecule 1: Bruno-like 4, RNA binding protein

Chain A: 52% 13% 32%




4.2.6 Score per residue for model 6

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.7 Score per residue for model 7

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.8 Score per residue for model 8

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.9 Score per residue for model 9

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.10 Score per residue for model 10

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.11 Score per residue for model 11

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.12 Score per residue for model 12

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.13 Score per residue for model 13

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.14 Score per residue for model 14

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.15 Score per residue for model 15 (medoid)

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.16 Score per residue for model 16

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.17 Score per residue for model 17

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.18 Score per residue for model 18

- Molecule 1: Bruno-like 4, RNA binding protein




4.2.19 Score per residue for model 19

- Molecule 1: Bruno-like 4, RNA binding protein



4.2.20 Score per residue for model 20

- Molecule 1: Bruno-like 4, RNA binding protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics, restrained molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *target function, structures with the lowest energy, structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	2.0.17
CYANA	refinement	2.0.17

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	539	535	535	16±2
All	All	10780	10700	10700	319

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:183:LEU:HD11	1:A:195:PHE:CE2	1.07	1.84	11	15
1:A:167:VAL:HG13	1:A:171:PHE:CE1	0.78	2.13	15	20
1:A:183:LEU:HD11	1:A:195:PHE:CD2	0.72	2.20	13	3
1:A:170:LEU:O	1:A:173:ALA:HB3	0.71	1.85	20	20
1:A:167:VAL:HG13	1:A:171:PHE:CD1	0.68	2.23	9	15
1:A:171:PHE:CB	1:A:177:ILE:HD11	0.67	2.19	4	3
1:A:205:GLN:HA	1:A:208:ILE:HD12	0.66	1.65	7	5
1:A:155:VAL:HG13	1:A:226:VAL:HG22	0.64	1.68	1	5
1:A:217:MET:SD	1:A:224:LEU:HD13	0.59	2.37	2	1
1:A:182:ILE:HD13	1:A:194:ALA:CB	0.59	2.28	11	1
1:A:171:PHE:HB3	1:A:177:ILE:HD11	0.59	1.74	4	1
1:A:155:VAL:HG21	1:A:171:PHE:CZ	0.58	2.32	12	2
1:A:216:THR:HG23	1:A:222:SER:O	0.57	1.99	16	18
1:A:153:LEU:HD21	1:A:228:PHE:CE1	0.57	2.33	12	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:177:ILE:HG23	1:A:197:LYS:O	0.57	2.00	10	3
1:A:155:VAL:HG22	1:A:226:VAL:HG13	0.56	1.76	8	4
1:A:183:LEU:CD1	1:A:195:PHE:CE2	0.55	2.77	11	1
1:A:211:LEU:HB3	1:A:224:LEU:HD23	0.55	1.78	1	2
1:A:153:LEU:HD12	1:A:198:TYR:CD1	0.55	2.36	4	4
1:A:183:LEU:HD21	1:A:195:PHE:CE2	0.54	2.38	17	1
1:A:157:MET:SD	1:A:225:VAL:HG21	0.53	2.43	2	2
1:A:171:PHE:CD2	1:A:196:VAL:HG11	0.53	2.39	8	4
1:A:164:GLU:HA	1:A:182:ILE:HD11	0.52	1.81	1	1
1:A:170:LEU:O	1:A:173:ALA:CB	0.52	2.58	20	20
1:A:205:GLN:C	1:A:205:GLN:CD	0.51	2.68	19	1
1:A:153:LEU:CD2	1:A:228:PHE:CE1	0.51	2.93	12	1
1:A:172:GLU:CD	1:A:172:GLU:C	0.51	2.70	12	2
1:A:155:VAL:HG22	1:A:226:VAL:HG22	0.51	1.83	8	1
1:A:180:CYS:SG	1:A:196:VAL:HG13	0.50	2.46	8	1
1:A:208:ILE:O	1:A:212:HIS:N	0.50	2.45	14	20
1:A:182:ILE:HD13	1:A:194:ALA:HB2	0.50	1.82	11	1
1:A:159:ASN:N	1:A:159:ASN:OD1	0.50	2.45	7	2
1:A:165:ASP:CG	1:A:166:ASP:N	0.50	2.65	9	1
1:A:159:ASN:OD1	1:A:160:LYS:N	0.49	2.45	18	1
1:A:164:GLU:O	1:A:168:ARG:CB	0.49	2.60	4	4
1:A:171:PHE:HB2	1:A:177:ILE:HD11	0.49	1.85	4	1
1:A:168:ARG:CD	1:A:168:ARG:C	0.49	2.80	7	1
1:A:159:ASN:ND2	1:A:218:PRO:O	0.49	2.46	8	3
1:A:171:PHE:CE2	1:A:196:VAL:HG21	0.48	2.43	19	6
1:A:153:LEU:HD21	1:A:208:ILE:HD11	0.48	1.86	1	2
1:A:156:GLY:O	1:A:158:LEU:N	0.47	2.48	20	2
1:A:165:ASP:OD1	1:A:166:ASP:N	0.47	2.48	9	1
1:A:212:HIS:CE1	1:A:225:VAL:HA	0.47	2.45	20	1
1:A:210:ALA:O	1:A:214:SER:CB	0.47	2.63	3	1
1:A:201:HIS:O	1:A:204:ALA:HB3	0.46	2.10	13	3
1:A:172:GLU:O	1:A:174:PHE:N	0.46	2.49	8	5
1:A:178:GLU:N	1:A:178:GLU:OE1	0.46	2.48	3	1
1:A:152:LYS:HE2	1:A:195:PHE:CD1	0.46	2.46	13	1
1:A:158:LEU:HD23	1:A:217:MET:HG3	0.46	1.86	6	1
1:A:152:LYS:HD3	1:A:195:PHE:CD1	0.45	2.46	17	3
1:A:153:LEU:CD1	1:A:198:TYR:CD1	0.45	2.99	4	2
1:A:171:PHE:O	1:A:173:ALA:N	0.45	2.49	2	3
1:A:176:ASN:O	1:A:198:TYR:CD2	0.45	2.69	13	1
1:A:201:HIS:O	1:A:202:ALA:C	0.45	2.55	13	13
1:A:212:HIS:ND1	1:A:224:LEU:O	0.45	2.49	20	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:212:HIS:HD1	1:A:225:VAL:HA	0.45	1.72	15	5
1:A:223:SER:O	1:A:224:LEU:C	0.45	2.55	3	3
1:A:198:TYR:OH	1:A:207:ALA:CB	0.45	2.65	2	3
1:A:167:VAL:O	1:A:168:ARG:C	0.45	2.55	4	6
1:A:158:LEU:HD23	1:A:217:MET:SD	0.45	2.52	1	1
1:A:216:THR:HG23	1:A:222:SER:C	0.44	2.32	16	1
1:A:216:THR:O	1:A:217:MET:C	0.44	2.55	10	9
1:A:224:LEU:HD12	1:A:225:VAL:H	0.44	1.72	4	3
1:A:156:GLY:O	1:A:157:MET:C	0.44	2.55	20	1
1:A:153:LEU:HD23	1:A:228:PHE:HA	0.44	1.89	2	1
1:A:153:LEU:HD23	1:A:228:PHE:CD1	0.44	2.47	12	1
1:A:153:LEU:CD2	1:A:208:ILE:HD11	0.44	2.43	1	2
1:A:153:LEU:HD11	1:A:204:ALA:HB1	0.44	1.89	6	1
1:A:159:ASN:HD21	1:A:218:PRO:N	0.44	2.10	8	1
1:A:152:LYS:HE3	1:A:195:PHE:CD1	0.44	2.48	15	1
1:A:206:ALA:O	1:A:210:ALA:CB	0.44	2.66	6	1
1:A:153:LEU:CD2	1:A:228:PHE:CD1	0.44	3.00	12	1
1:A:224:LEU:HG	1:A:226:VAL:HG23	0.43	1.89	14	1
1:A:183:LEU:HD12	1:A:193:CYS:HB3	0.43	1.89	14	1
1:A:175:GLY:CA	1:A:203:GLU:OE1	0.43	2.67	3	1
1:A:172:GLU:C	1:A:174:PHE:N	0.43	2.72	18	5
1:A:168:ARG:NH1	1:A:172:GLU:OE1	0.43	2.52	18	1
1:A:201:HIS:O	1:A:204:ALA:N	0.42	2.52	18	4
1:A:213:GLY:CA	1:A:223:SER:HB3	0.42	2.45	3	1
1:A:183:LEU:HD21	1:A:195:PHE:HE2	0.42	1.73	11	1
1:A:208:ILE:HG12	1:A:226:VAL:HG12	0.42	1.91	16	1
1:A:211:LEU:O	1:A:214:SER:CB	0.42	2.68	5	2
1:A:167:VAL:CG1	1:A:171:PHE:CD1	0.42	3.01	11	1
1:A:165:ASP:O	1:A:168:ARG:N	0.42	2.53	1	1
1:A:171:PHE:C	1:A:173:ALA:N	0.42	2.73	6	2
1:A:177:ILE:CG2	1:A:178:GLU:N	0.42	2.83	16	3
1:A:155:VAL:HG21	1:A:171:PHE:HZ	0.41	1.75	17	2
1:A:159:ASN:ND2	1:A:162:GLN:HG3	0.41	2.30	7	1
1:A:205:GLN:O	1:A:206:ALA:C	0.41	2.59	19	1
1:A:212:HIS:CE1	1:A:224:LEU:O	0.41	2.73	20	1
1:A:215:GLN:HG3	1:A:216:THR:N	0.41	2.31	9	1
1:A:191:LYS:O	1:A:192:GLY:C	0.40	2.59	14	1
1:A:183:LEU:HD11	1:A:195:PHE:HE2	0.40	1.74	6	1
1:A:211:LEU:HB3	1:A:224:LEU:CD2	0.40	2.46	4	1
1:A:205:GLN:O	1:A:208:ILE:N	0.40	2.55	19	1
1:A:159:ASN:ND2	1:A:218:PRO:HD2	0.40	2.32	8	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:208:ILE:HG22	1:A:209:ASN:N	0.40	2.32	3	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	71/105 (68%)	54±3 (76±4%)	15±3 (22±4%)	2±1 (3±2%)	8	44
All	All	1420/2100 (68%)	1075 (76%)	308 (22%)	37 (3%)	8	44

All 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	208	ILE	12
1	A	224	LEU	6
1	A	173	ALA	5
1	A	217	MET	4
1	A	172	GLU	3
1	A	157	MET	2
1	A	176	ASN	2
1	A	190	SER	1
1	A	223	SER	1
1	A	162	GLN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	58/84 (69%)	56±1 (97±2%)	2±1 (3±2%)	40	87

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1160/1680 (69%)	1121 (97%)	39 (3%)	40 87

All 18 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	172	GLU	7
1	A	165	ASP	5
1	A	157	MET	3
1	A	214	SER	3
1	A	159	ASN	3
1	A	227	LYS	3
1	A	197	LYS	2
1	A	217	MET	2
1	A	178	GLU	2
1	A	193	CYS	1
1	A	223	SER	1
1	A	203	GLU	1
1	A	160	LYS	1
1	A	168	ARG	1
1	A	215	GLN	1
1	A	179	GLU	1
1	A	216	THR	1
1	A	205	GLN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided