



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 11:45 AM EDT

PDB ID : 2QMJ
Title : Crystal Structure of the N-terminal Subunit of Human Maltase-Glucoamylase
in Complex with Acarbose
Authors : Sim, L.; Rose, D.R.
Deposited on : 2007-07-16
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	870	
2	X	2	
3	B	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	3004	-	-	X	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7739 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltase-glucoamylase, intestinal.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	863	6940	4426	1176	1311	27	0	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	869	ALA	-	expression tag	UNP O43451
A	870	HIS	-	expression tag	UNP O43451

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	X	2	28	16	2	10	0	0	0

- Molecule 3 is an oligosaccharide called 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	B	3	44	25	1	18	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
4	A	1	28	16	2	10	0	1
4	A	1	14	8	1	5	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	6	3	3	0	0
5	A	1	6	3	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	626	Total O 626 626	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltase-glucoamylase, intestinal



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 4,6-dideoxy-4-[(1S,4R,5S,6S)-4,5,6-trihydroxy-3-(hydroxymethyl)cyclohex-2-en-1-yl]amino}-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.97Å 109.37Å 109.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.78 – 1.90 19.77 – 1.89	Depositor EDS
% Data completeness (in resolution range)	98.2 (19.78-1.90) 98.2 (19.77-1.89)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.177 , 0.215 0.177 , 0.212	Depositor DCC
R_{free} test set	4082 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtrriage
Anisotropy	0.200	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for -h,l,k	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7739	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, AC1, SO4, GOL, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.77	0/7154	0.79	2/9756 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	624	ARG	NE-CZ-NH2	-6.59	117.00	120.30
1	A	624	ARG	NE-CZ-NH1	5.71	123.15	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	300	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6940	0	6618	95	0
2	X	28	0	25	0	0
3	B	44	0	30	1	0
4	A	42	0	39	3	0
5	A	54	0	72	14	0
6	A	5	0	0	0	0
7	A	626	0	0	11	0
All	All	7739	0	6784	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:204:THR:HG21	1:A:473:LEU:CD2	1.71	1.19
1:A:204:THR:HG21	1:A:473:LEU:HD23	1.23	1.15
1:A:779:VAL:HB	5:A:3004:GOL:H32	1.52	0.91
1:A:26:CYS:HG	1:A:44:CYS:HG	1.21	0.85
1:A:814:ASN:HB3	7:A:4168:HOH:O	1.77	0.83
5:A:3004:GOL:H12	7:A:4169:HOH:O	1.80	0.82
1:A:645:HIS:NE2	5:A:3004:GOL:H31	1.94	0.82
1:A:98:HIS:HE1	1:A:261:ASP:OD1	1.62	0.81
1:A:135:ARG:HB3	1:A:136:GLN:HE21	1.47	0.80
1:A:777:ASP:HA	5:A:3004:GOL:H2	1.63	0.79
1:A:712:ARG:H	1:A:714:GLN:HE21	1.32	0.78
1:A:777:ASP:CA	5:A:3004:GOL:H2	2.13	0.78
1:A:727:LEU:H	5:A:3008:GOL:H11	1.51	0.76
1:A:493:GLN:HE21	1:A:497:HIS:HD2	1.37	0.71
1:A:741[B]:ASN:OD1	4:A:2001[B]:NAG:O5	2.00	0.70
1:A:135:ARG:HB3	1:A:136:GLN:NE2	2.07	0.69
1:A:793:GLN:H	1:A:793:GLN:CD	1.96	0.68
1:A:544:THR:CG2	1:A:577:LEU:HD22	2.24	0.67
1:A:204:THR:HG21	1:A:473:LEU:HD21	1.75	0.67
1:A:717:GLU:HG3	1:A:717:GLU:O	1.95	0.66
1:A:493:GLN:NE2	1:A:497:HIS:HD2	1.94	0.65
5:A:3004:GOL:C1	7:A:4169:HOH:O	2.40	0.65
1:A:777:ASP:N	5:A:3004:GOL:H2	2.12	0.64
1:A:201:ASN:O	1:A:497:HIS:HE1	1.81	0.64
1:A:90:GLU:OE1	1:A:98:HIS:HD2	1.82	0.63
1:A:204:THR:CG2	1:A:473:LEU:HD23	2.15	0.62
1:A:835:PRO:HA	1:A:836:SER:HB3	1.82	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:711:TRP:CG	1:A:716:VAL:HG11	2.37	0.60
1:A:183:HIS:HE1	1:A:198:PRO:O	1.84	0.60
1:A:576:ALA:O	1:A:577:LEU:HB2	2.02	0.59
1:A:183:HIS:HD2	1:A:185:HIS:NE2	2.01	0.59
1:A:446:GLU:N	1:A:447:VAL:HA	2.18	0.59
1:A:691:VAL:CG2	7:A:4343:HOH:O	2.50	0.58
1:A:741[A]:ASN:ND2	4:A:2001[A]:NAG:H82	2.19	0.58
1:A:179:GLY:O	1:A:190:HIS:HE1	1.87	0.57
1:A:631:TYR:HA	1:A:634:LEU:HG	1.85	0.57
1:A:424:TRP:CE2	1:A:428:PHE:HE2	2.22	0.56
1:A:779:VAL:HB	5:A:3004:GOL:C3	2.30	0.56
1:A:793:GLN:CD	1:A:793:GLN:N	2.59	0.56
1:A:493:GLN:NE2	1:A:497:HIS:CD2	2.74	0.55
1:A:464:ASN:ND2	1:A:484:MET:H	2.04	0.55
1:A:68[B]:ARG:HG3	1:A:104:GLN:OE1	2.05	0.55
1:A:98:HIS:CE1	1:A:261:ASP:OD1	2.52	0.54
1:A:591:GLY:HA2	1:A:594:TYR:CD2	2.42	0.54
1:A:201:ASN:H	1:A:498:ASN:ND2	2.05	0.54
1:A:777:ASP:H	5:A:3004:GOL:H2	1.72	0.53
1:A:164:GLN:NE2	1:A:462:ASN:HD22	2.07	0.53
1:A:291:ALA:O	1:A:295:HIS:HE1	1.91	0.53
1:A:681:THR:HG22	1:A:691:VAL:HG11	1.90	0.53
1:A:779:VAL:CB	5:A:3004:GOL:H32	2.31	0.53
1:A:618:LEU:HD11	1:A:723:ASP:HB3	1.91	0.52
1:A:183:HIS:CD2	1:A:185:HIS:NE2	2.78	0.52
1:A:201:ASN:H	1:A:498:ASN:HD21	1.57	0.52
1:A:295:HIS:HD2	1:A:325:HIS:NE2	2.08	0.52
1:A:53:HIS:HD2	1:A:72:LEU:HG	1.75	0.52
1:A:148:ASN:ND2	7:A:4406:HOH:O	2.44	0.51
1:A:488:GLN:HB2	1:A:496:ILE:HD11	1.92	0.50
1:A:625:HIS:HE1	1:A:704:GLU:OE2	1.94	0.50
1:A:149:ARG:NH2	1:A:225:ASP:OD2	2.45	0.49
1:A:203:ASP:O	1:A:204:THR:HG23	2.12	0.49
1:A:725:ILE:O	5:A:3008:GOL:H31	2.12	0.49
1:A:473:LEU:O	1:A:474:ASP:HB2	2.12	0.49
1:A:464:ASN:HD21	1:A:484:MET:H	1.59	0.48
1:A:691:VAL:HG23	7:A:4147:HOH:O	2.12	0.48
1:A:702:ASP:OD1	5:A:3008:GOL:H12	2.14	0.47
1:A:243:VAL:HG11	1:A:251:ILE:HD11	1.97	0.47
1:A:817:LYS:NZ	7:A:4448:HOH:O	2.46	0.47
1:A:538:HIS:O	1:A:568:VAL:HA	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:366:ASP:HB3	7:A:4054:HOH:O	2.14	0.46
1:A:9:GLU:HB2	1:A:41:VAL:HG11	1.97	0.46
1:A:544:THR:HG22	1:A:577:LEU:HD22	1.98	0.46
1:A:493:GLN:HE21	1:A:497:HIS:CD2	2.25	0.46
1:A:164:GLN:HE22	1:A:462:ASN:HB2	1.80	0.46
1:A:763:GLU:CD	1:A:792:THR:HG22	2.36	0.46
1:A:243:VAL:CG1	1:A:251:ILE:HD11	2.46	0.45
1:A:791:VAL:HA	1:A:795:ARG:O	2.17	0.45
1:A:707:SER:HB3	1:A:848:LEU:HG	1.99	0.44
1:A:26:CYS:CB	1:A:44:CYS:HG	2.30	0.44
1:A:570:PRO:HD2	1:A:594:TYR:CD2	2.52	0.44
1:A:225:ASP:OD1	1:A:227:SER:OG	2.25	0.43
1:A:317:ALA:HB1	1:A:620:LEU:HD21	2.01	0.43
1:A:655:LEU:HD22	1:A:668:VAL:HG11	2.01	0.43
1:A:654:PRO:HD3	7:A:4261:HOH:O	2.19	0.43
1:A:33:TRP:CZ2	1:A:35:PRO:HG3	2.54	0.43
1:A:33:TRP:CE2	1:A:35:PRO:HG3	2.54	0.43
1:A:794:ASN:N	7:A:4190:HOH:O	2.52	0.43
1:A:835:PRO:HA	1:A:836:SER:CB	2.48	0.43
1:A:739:GLN:HB3	1:A:750:ASN:HB3	2.00	0.43
1:A:443:ASP:OD2	3:B:3:AC1:HCB2	2.20	0.42
1:A:793:GLN:O	1:A:794:ASN:HB2	2.20	0.41
1:A:572:ILE:O	1:A:573:CYS:HB2	2.20	0.41
1:A:750:ASN:HD21	4:A:2001[A]:NAG:C8	2.34	0.41
1:A:649:ASP:O	7:A:4619:HOH:O	2.22	0.41
1:A:201:ASN:O	1:A:497:HIS:CE1	2.68	0.40
1:A:526:ARG:O	1:A:538:HIS:CE1	2.73	0.40
1:A:779:VAL:H	5:A:3004:GOL:H32	1.86	0.40
1:A:453:GLY:HA3	1:A:458:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	864/870 (99%)	819 (95%)	42 (5%)	3 (0%)	41 31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	80	SER
1	A	793	GLN
1	A	405	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	761/762 (100%)	752 (99%)	9 (1%)	71 70

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	VAL
1	A	161	PHE
1	A	322	ASP
1	A	347	PHE
1	A	539	TRP
1	A	577	LEU
1	A	793	GLN
1	A	799	ASN
1	A	848	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	A	62	ASN
1	A	84	ASN
1	A	98	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	136	GLN
1	A	148	ASN
1	A	164	GLN
1	A	167	GLN
1	A	183	HIS
1	A	190	HIS
1	A	193	ASN
1	A	239	ASN
1	A	295	HIS
1	A	464	ASN
1	A	493	GLN
1	A	497	HIS
1	A	498	ASN
1	A	625	HIS
1	A	628	ASN
1	A	669	HIS
1	A	714	GLN
1	A	781	ASN
1	A	802	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GLC	B	1	3	12,12,12	0.82	0	17,17,17	1.32	1 (5%)
3	GLC	B	2	3	11,11,12	0.42	0	15,15,17	1.24	2 (13%)
3	AC1	B	3	3	21,22,23	0.90	0	22,32,34	1.39	3 (13%)
2	NAG	X	1	1,2	14,14,15	0.64	0	17,19,21	1.27	2 (11%)
2	NAG	X	2	2	14,14,15	0.62	0	17,19,21	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GLC	B	1	3	-	0/2/22/22	0/1/1/1
3	GLC	B	2	3	-	0/2/19/22	0/1/1/1
3	AC1	B	3	3	-	4/6/43/46	0/2/2/2
2	NAG	X	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	X	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	3	AC1	O6B-C6B-C5B	-4.05	102.82	112.50
2	X	1	NAG	O5-C1-C2	-3.23	106.18	111.29
3	B	2	GLC	C1-O5-C5	2.91	116.14	112.19
3	B	1	GLC	O4-C4-C3	-2.69	104.13	110.35
3	B	3	AC1	O3B-C3B-C2B	2.34	115.75	110.35
2	X	1	NAG	C1-O5-C5	2.23	115.22	112.19
3	B	3	AC1	C1-O5-C5	2.21	117.80	112.78
3	B	2	GLC	C1-C2-C3	2.04	112.17	109.67

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	3	AC1	C7B-C1B-N4A-C4
2	X	2	NAG	C4-C5-C6-O6
3	B	3	AC1	C4A-C5B-C6B-O6B
3	B	3	AC1	C7B-C5B-C6B-O6B
2	X	2	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	3	AC1	C3-C4-N4A-C1B


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	3	AC1	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	3007	-	5,5,5	0.22	0	5,5,5	0.58	0
4	NAG	A	2001[A]	1	14,14,15	0.57	0	17,19,21	2.04	3 (17%)
4	NAG	A	2001[B]	1	14,14,15	0.50	0	17,19,21	2.11	4 (23%)
5	GOL	A	3006	-	5,5,5	0.38	0	5,5,5	0.81	0
5	GOL	A	3004	-	5,5,5	0.42	0	5,5,5	0.61	0
5	GOL	A	3005	-	5,5,5	0.28	0	5,5,5	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	A	4001	-	4,4,4	0.22	0	6,6,6	0.52	0
5	GOL	A	3009	-	5,5,5	0.27	0	5,5,5	0.37	0
4	NAG	A	2005	1	14,14,15	0.59	0	17,19,21	1.76	3 (17%)
5	GOL	A	3003	-	5,5,5	0.36	0	5,5,5	0.43	0
5	GOL	A	3002	-	5,5,5	0.39	0	5,5,5	0.33	0
5	GOL	A	3001	-	5,5,5	0.25	0	5,5,5	0.49	0
5	GOL	A	3008	-	5,5,5	0.46	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	3007	-	-	3/4/4/4	-
4	NAG	A	2001[A]	1	-	3/6/23/26	0/1/1/1
4	NAG	A	2001[B]	1	-	4/6/23/26	0/1/1/1
5	GOL	A	3006	-	-	0/4/4/4	-
5	GOL	A	3004	-	-	1/4/4/4	-
5	GOL	A	3005	-	-	2/4/4/4	-
5	GOL	A	3009	-	-	1/4/4/4	-
4	NAG	A	2005	1	-	0/6/23/26	0/1/1/1
5	GOL	A	3003	-	-	0/4/4/4	-
5	GOL	A	3002	-	-	2/4/4/4	-
5	GOL	A	3001	-	-	2/4/4/4	-
5	GOL	A	3008	-	-	2/4/4/4	-

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2001[A]	NAG	C2-N2-C7	6.77	132.55	122.90
4	A	2001[B]	NAG	C1-O5-C5	6.04	120.37	112.19
4	A	2005	NAG	C1-O5-C5	5.00	118.97	112.19
4	A	2001[B]	NAG	C4-C3-C2	-4.07	105.05	111.02
4	A	2001[B]	NAG	C1-C2-N2	2.78	115.24	110.49
4	A	2005	NAG	C2-N2-C7	2.70	126.75	122.90
4	A	2001[A]	NAG	O5-C5-C6	2.70	111.43	107.20
4	A	2001[B]	NAG	C6-C5-C4	-2.65	106.79	113.00
4	A	2001[A]	NAG	O5-C1-C2	-2.64	107.12	111.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2005	NAG	O5-C5-C4	-2.04	105.86	110.83

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	2001[A]	NAG	C8-C7-N2-C2
4	A	2001[A]	NAG	O7-C7-N2-C2
4	A	2001[B]	NAG	C8-C7-N2-C2
4	A	2001[B]	NAG	O7-C7-N2-C2
5	A	3002	GOL	O1-C1-C2-C3
5	A	3005	GOL	O1-C1-C2-C3
5	A	3007	GOL	C1-C2-C3-O3
5	A	3008	GOL	C1-C2-C3-O3
4	A	2001[B]	NAG	O5-C5-C6-O6
4	A	2001[B]	NAG	C4-C5-C6-O6
5	A	3001	GOL	C1-C2-C3-O3
5	A	3007	GOL	O1-C1-C2-C3
5	A	3008	GOL	O2-C2-C3-O3
5	A	3004	GOL	O1-C1-C2-O2
4	A	2001[A]	NAG	C4-C5-C6-O6
5	A	3002	GOL	O1-C1-C2-O2
5	A	3007	GOL	O2-C2-C3-O3
5	A	3001	GOL	O2-C2-C3-O3
5	A	3005	GOL	O1-C1-C2-O2
5	A	3009	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2001[A]	NAG	2	0
4	A	2001[B]	NAG	1	0
5	A	3004	GOL	11	0
5	A	3008	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	863/870 (99%)	0.08	44 (5%) 28 31	11, 21, 36, 60	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	VAL	7.0
1	A	836	SER	6.5
1	A	48	LYS	5.1
1	A	835	PRO	4.9
1	A	36	GLN	4.6
1	A	73	PRO	4.4
1	A	375	SER	4.4
1	A	870	HIS	4.4
1	A	838	THR	4.4
1	A	793	GLN	4.4
1	A	50	HIS	4.1
1	A	49	ASN	3.9
1	A	37	GLY	3.9
1	A	121	GLY	3.8
1	A	80	SER	3.6
1	A	105	THR	3.5
1	A	376	SER	3.5
1	A	710	ARG	3.4
1	A	51	SER	3.2
1	A	76	PRO	3.1
1	A	616	ASP	3.1
1	A	377	LYS	3.1
1	A	374	SER	3.1
1	A	792	THR	2.7
1	A	433	ASN	2.7
1	A	106	ASN	2.6
1	A	794	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	796	LEU	2.4
1	A	117	GLN	2.4
1	A	74	SER	2.3
1	A	826	ASN	2.3
1	A	135	ARG	2.3
1	A	866	VAL	2.3
1	A	847	ASN	2.2
1	A	615	ALA	2.2
1	A	832	ASN	2.1
1	A	857	ASP	2.1
1	A	38	ALA	2.1
1	A	477	LEU	2.1
1	A	191	ASP	2.1
1	A	869	ALA	2.1
1	A	20	PRO	2.1
1	A	818	ILE	2.1
1	A	204	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	B	1	12/12	0.79	0.36	51,55,59,59	0
2	NAG	X	2	14/15	0.87	0.30	44,49,52,53	0
3	GLC	B	2	11/12	0.88	0.17	36,45,48,49	0
2	NAG	X	1	14/15	0.94	0.10	28,31,35,39	0
3	AC1	B	3	21/22	0.94	0.10	18,26,28,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain X:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q < 0.9
5	GOL	A	3009	6/6	0.68	0.19	67,68,68,69	0
5	GOL	A	3004	6/6	0.78	0.20	41,43,44,45	0
5	GOL	A	3005	6/6	0.82	0.18	40,45,45,45	0
4	NAG	A	2005	14/15	0.82	0.25	42,46,51,52	0
4	NAG	A	2001[B]	14/15	0.84	0.19	25,27,31,31	14
4	NAG	A	2001[A]	14/15	0.84	0.19	20,23,27,27	14
5	GOL	A	3007	6/6	0.84	0.15	30,37,42,47	0
5	GOL	A	3001	6/6	0.84	0.15	35,39,40,46	0
5	GOL	A	3003	6/6	0.90	0.20	32,36,37,39	0
5	GOL	A	3008	6/6	0.91	0.13	43,47,48,51	0
5	GOL	A	3006	6/6	0.93	0.11	38,40,42,43	0
5	GOL	A	3002	6/6	0.94	0.11	27,29,31,31	0
6	SO4	A	4001	5/5	0.97	0.13	38,40,41,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.