

wwPDB X-ray Structure Validation Summary Report (i)

Aug 27, 2023 – 08:13 AM EDT

PDB ID : 3HE3

Title: Crystal Structure of UDP-galactopyranose mutase in complex with UDP

Authors: Partha, S.K.; van Straaten, K.E.; Sanders, D.A.R.

Deposited on : 2009-05-07

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar resolution

Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	397	75%	16%	•	8%			
1	В	397	75%	16%		8%			
1	С	397	75%	15%		9%			
1	D	397	72%	17%	•	9%			
1	Е	397	69%	20%	•	8%			

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain					
	-	225	2%					
1	F	397	76%	15%	٠	9%		
4		207	<u>%</u>					
1	G	397	72%	18%	•	9%		
1	TT	207	%					
1	Н	397	71%	19%	•	9%		
1	т	907	% •			_		
1	1	397	76%	13%	•	8%		
1		207	2%					
1	J	397	72%	17%	• •	9%		



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 31550 atoms, of which 0 are hydrogens and 0 are deuteriums.


In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called UDP-galactopyranose mutase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	366	Total	С	N	О	S	0	0	0
1	Λ	300	2993	1909	520	556	8	0	U	0
1	В	364	Total	\mathbf{C}	N	Ο	S	1	0	0
1	D	304	2971	1896	516	551	8	1	U	0
1	C	362	Total	\mathbf{C}	N	O	S	1	0	0
	C	902	2950	1885	511	546	8	1	U	U
1	D	362	Total	\mathbf{C}	N	O	S	1	0	0
	D	902	2960	1891	514	547	8		Ü	U
1	E	364	Total	\mathbf{C}	N	O	S	0	0	0
	L	301	2975	1899	516	552	8	O	O .	U
1	F	363	Total	\mathbf{C}	N	O	S	1	0	0
1	1	909	2969	1896	516	549	8	1	U	U
1	G	363	Total	\mathbf{C}	N	O	S	0	0	0
1	G G	909	2965	1894	515	548	8	0	O .	U
1	Н	363	Total	\mathbf{C}	Ν	O	S	0	0	0
1	11	909	2969	1896	516	549	8	0	O O	U
1	I	364	Total	\mathbf{C}	N	Ο	S	0	0	0
1	1	504	2975	1899	517	551	8		U	0
1	J	361	Total	С	Ν	О	S	0	0	0
1	9	301	2951	1885	512	546	8		U	

• Molecule 2 is URIDINE-5'-DIPHOSPHATE (three-letter code: UDP) (formula: $C_9H_{14}N_2O_{12}P_2$).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	A	1	Total	С	N	О	Р	0	0	
	Λ	1	25	9	2	12	2	0	U	
2	В	1	Total	С	N	Ο	Р	0	0	
	D	1	25	9	2	12	2	0	U	
2	$^{\rm C}$	1	Total	С	N	Ο	Р	0	0	
	C	1	25	9	2	12	2	0	U	
2	D	1	Total	С	Ν	Ο	Р	0	0	
	D	1	25	9	2	12	2	U	0	
2	E	1	Total	С	N	Ο	Р	0	0	
		Ŧ	25	9	2	12	2	O O	Ŭ	
2	F	1	Total	С	Ν	Ο	Р	0	0	
	1	1	25	9	2	12	2	O	0	
2	G	1	Total	С	Ν	Ο	Р	0	0	
	<u> </u>	1	25	9	2	12	2	O	Ü	
2	Н	1	Total	С	Ν	Ο	Р	0	0	
	11	1	25	9	2	12	2	Ŭ	Ü	
2	I	1	Total	С	N	Ο	Р	0	0	
	1	1	25	9	2	12	2		9	
2	J	1	Total	С	N	Ο	Р	0	0	
	9	1	25	9	2	12	2		U	

 \bullet Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2).$





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
3	A	1	Total	С	N	О	Р	0	0	
3	Λ	1	53	27	9	15	2	U	U	
3	В	1	Total	С	N	Ο	Р	0	0	
	D	1	53	27	9	15	2	O	U	
3	$^{\rm C}$	1	Total	\mathbf{C}	N	Ο	Р	0	0	
		1	53	27	9	15	2	O	U	
3	D	1	Total	\mathbf{C}	N	Ο	Р	0	0	
	D	1	53	27	9	15	2	U	U	
3	E	1	Total	\mathbf{C}	N	Ο	Р	0	0	
	L	1	53	27	9	15	2	O		
3	F	1	Total	С	N	Ο	Р	0	0	
	1	1	53	27	9	15	2	O		
3	G	1	Total	\mathbf{C}	N	Ο	Р	0	0	
	<u> </u>	1	53	27	9	15	2	Ŭ	0	
3	Н	1	Total	С	N	Ο	Р	0	0	
	11	1	53	27	9	15	2	Ŭ	Ŭ .	
3	I	1	Total	\mathbf{C}	N	Ο	Р	0	0	
	*	1	53	27	9	15	2	Ü	U	
3	J	J 1	Total	\mathbf{C}	N	Ο	Р	0	0	
		1	53	27	9	15	2	O	U	

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	135	Total O 135 135	0	0
4	В	123	Total O 123 123	0	0

Continued on next page...



Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	С	94	Total O 94 94	0	0
4	D	95	Total O 95 95	0	0
4	Е	79	Total O 79 79	0	0
4	F	107	Total O 107 107	0	0
4	G	104	Total O 104 104	0	0
4	Н	133	Total O 133 133	0	0
4	I	110	Total O 110 110	0	0
4	J	112	Total O 112 112	0	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UDP-galactopyranose mutase
















4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	134.05Å 176.87Å 222.92Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.88 - 2.40	Depositor
Resolution (A)	39.88 - 2.40	EDS
% Data completeness	98.8 (39.88-2.40)	Depositor
(in resolution range)	98.8 (39.88-2.40)	EDS
R_{merge}	0.18	Depositor
R_{sym}	0.18	Depositor
$< I/\sigma(I) > 1$	1.56 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
D.D.	0.208 , 0.254	Depositor
R, R_{free}	0.202 , 0.245	DCC
R_{free} test set	10272 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 38.6	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	31550	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 1.83% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Во	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.32	0/3077	0.45	0/4183
1	В	0.26	0/3055	0.44	0/4155
1	С	0.27	0/3034	0.44	0/4127
1	D	0.25	0/3044	0.44	0/4139
1	Е	0.33	0/3059	0.45	0/4159
1	F	0.28	0/3053	0.46	0/4151
1	G	0.28	0/3049	0.44	0/4146
1	Н	0.31	0/3053	0.45	0/4151
1	I	0.29	0/3059	0.45	0/4159
1	J	0.39	0/3035	0.56	3/4128 (0.1%)
All	All	0.30	0/30518	0.46	3/41498 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	oms Z Obser		$\operatorname{Ideal}({}^{o})$
1	J	154	GLU	OE1-CD-OE2	-9.80	111.54	123.30
1	J	154	GLU	CA-CB-CG	-8.93	93.76	113.40
1	J	155	GLN	N-CA-CB	8.32	125.58	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2993	0	2862	41	0
1	В	2971	0	2837	37	0
1	С	2950	0	2812	47	0
1	D	2960	0	2835	52	0
1	Ε	2975	0	2846	63	0
1	F	2969	0	2843	34	0
1	G	2965	0	2837	48	0
1	Н	2969	0	2843	51	0
1	I	2975	0	2848	41	0
1	J	2951	0	2822	44	0
2	A	25	0	11	2	0
2	В	25	0	11	0	0
2	С	25	0	11	0	0
2	D	25	0	11	0	0
2	Ε	25	0	11	1	0
2	F	25	0	11	0	0
2	G	25	0	11	0	0
2	Н	25	0	11	0	0
2	I	25	0	11	0	0
2	J	25	0	11	0	0
3	A	53	0	31	1	0
3	В	53	0	31	2	0
3	С	53	0	31	1	0
3	D	53	0	31	0	0
3	Ε	53	0	31	1	0
3	F	53	0	31	1	0
3	G	53	0	31	1	0
3	Н	53	0	31	1	0
3	I	53	0	31	0	0
3	J	53	0	31	1	0
4	A	135	0	0	1	0
4	В	123	0	0	1	0
4	С	94	0	0	2	0
4	D	95	0	0	2	0
4	Ε	79	0	0	1	0
4	F	107	0	0	0	0
4	G	104	0	0	1	0
4	Н	133	0	0	1	0
4	I	110	0	0	1	0
4	J	112	0	0	2	0
All	All	31550	0	28805	433	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 7.



The worst 5 of 433 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{array}{ll} ext{Interatomic} \ ext{distance} \ (ext{Å}) \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (ext{Å}) \end{aligned}$	
1:I:154:GLU:CD	1:I:154:GLU:H	1.51	1.04	
1:J:155:GLN:HG2	1:J:157:ARG:CZ	1.99	0.93	
1:G:58:VAL:HG23	1:G:238:MET:HB3	1.53	0.87	
1:H:162:VAL:O	1:H:166:LYS:HD2	1.76	0.85	
1:G:154:GLU:HG3	1:G:155:GLN:H	1.44	0.83	

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	364/397~(92%)	352 (97%)	12 (3%)	0	100	100
1	В	362/397~(91%)	355 (98%)	7 (2%)	0	100	100
1	C	360/397 (91%)	351 (98%)	8 (2%)	1 (0%)	41	55
1	D	360/397 (91%)	349 (97%)	10 (3%)	1 (0%)	41	55
1	E	362/397~(91%)	351 (97%)	10 (3%)	1 (0%)	41	55
1	F	361/397 (91%)	349 (97%)	12 (3%)	0	100	100
1	G	361/397~(91%)	349 (97%)	12 (3%)	0	100	100
1	Н	361/397 (91%)	353 (98%)	8 (2%)	0	100	100
1	I	362/397~(91%)	347 (96%)	15 (4%)	0	100	100
1	J	359/397~(90%)	345 (96%)	13 (4%)	1 (0%)	41	55
All	All	3612/3970 (91%)	3501 (97%)	107 (3%)	4 (0%)	51	68

All (4) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	Е	154	GLU
1	J	154	GLU
1	D	357	GLN
1	С	154	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	318/346 (92%)	295 (93%)	23 (7%)	1	4 23	
1	В	315/346 (91%)	299 (95%)	16 (5%)	2	4 39	
1	C	311/346 (90%)	296 (95%)	15 (5%)	2	5 41	
1	D	314/346 (91%)	294 (94%)	20 (6%)	1	7 28	
1	E	316/346 (91%)	296 (94%)	20 (6%)	1	8 28	
1	F	315/346 (91%)	301 (96%)	14 (4%)	2	8 45	
1	G	314/346 (91%)	295 (94%)	19 (6%)	1	8 30	
1	Н	315/346 (91%)	296 (94%)	19 (6%)	1	9 31	
1	I	316/346 (91%)	294 (93%)	22 (7%)	1	5 24	
1	J	313/346 (90%)	292 (93%)	21 (7%)	1	6 26	
All	All	3147/3460 (91%)	2958 (94%)	189 (6%)	1	9 31	

5 of 189 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	162	VAL
1	Н	362	VAL
1	G	261	VAL
1	Н	112	LEU
1	I	112	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:



Mol	Chain	Res	Type
1	I	357	GLN
1	J	311	HIS
1	I	390	GLN
1	F	108	GLN
1	I	311	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Вс	ond leng	ths	Bond angles			
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	UDP	A	400	-	24,26,26	0.94	0	37,40,40	1.51	6 (16%)	
2	UDP	D	400	-	24,26,26	0.95	0	37,40,40	1.56	7 (18%)	
2	UDP	В	400	-	24,26,26	0.95	1 (4%)	37,40,40	1.57	6 (16%)	
2	UDP	J	400	-	24,26,26	0.96	0	37,40,40	1.58	6 (16%)	
3	FAD	A	450	-	53,58,58	1.13	4 (7%)	68,89,89	1.30	7 (10%)	
3	FAD	D	450	-	53,58,58	1.12	4 (7%)	68,89,89	1.27	8 (11%)	
3	FAD	J	450	-	53,58,58	1.15	4 (7%)	68,89,89	1.31	10 (14%)	
3	FAD	С	450	-	53,58,58	1.11	4 (7%)	68,89,89	1.31	10 (14%)	



Mol	Trino	Chain	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UDP	F	400	-	24,26,26	0.93	0	37,40,40	1.58	6 (16%)
2	UDP	G	400	-	24,26,26	0.94	1 (4%)	37,40,40	1.55	7 (18%)
2	UDP	Е	400	-	24,26,26	0.93	0	37,40,40	1.50	5 (13%)
2	UDP	С	400	-	24,26,26	0.95	0	37,40,40	1.54	6 (16%)
3	FAD	В	450	-	53,58,58	1.13	4 (7%)	68,89,89	1.30	9 (13%)
3	FAD	F	450	-	53,58,58	1.14	4 (7%)	68,89,89	1.30	8 (11%)
2	UDP	Н	400	-	24,26,26	0.95	0	37,40,40	1.54	6 (16%)
3	FAD	I	450	-	53,58,58	1.13	4 (7%)	68,89,89	1.30	10 (14%)
3	FAD	Н	450	-	53,58,58	1.13	4 (7%)	68,89,89	1.26	9 (13%)
3	FAD	Е	450	-	53,58,58	1.12	4 (7%)	68,89,89	1.25	9 (13%)
3	FAD	G	450	-	53,58,58	1.13	4 (7%)	68,89,89	1.27	8 (11%)
2	UDP	I	400	-	24,26,26	0.94	1 (4%)	37,40,40	1.57	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UDP	A	400	-	-	3/16/32/32	0/2/2/2
2	UDP	D	400	-	-	2/16/32/32	0/2/2/2
2	UDP	В	400	-	-	3/16/32/32	0/2/2/2
2	UDP	J	400	-	-	2/16/32/32	0/2/2/2
3	FAD	A	450	-	-	5/30/50/50	0/6/6/6
3	FAD	D	450	-	-	4/30/50/50	0/6/6/6
3	FAD	J	450	-	-	3/30/50/50	0/6/6/6
3	FAD	С	450	-	-	5/30/50/50	0/6/6/6
2	UDP	F	400	-	-	1/16/32/32	0/2/2/2
2	UDP	G	400	-	-	2/16/32/32	0/2/2/2
2	UDP	Е	400	-	-	2/16/32/32	0/2/2/2
2	UDP	С	400	-	-	1/16/32/32	0/2/2/2
3	FAD	В	450	-	-	2/30/50/50	0/6/6/6
3	FAD	F	450	-	=	1/30/50/50	0/6/6/6
2	UDP	Н	400	-	-	2/16/32/32	0/2/2/2
3	FAD	I	450	-	-	7/30/50/50	0/6/6/6
3	FAD	Н	450	-	-	3/30/50/50	0/6/6/6

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	E	450	-	-	3/30/50/50	0/6/6/6
3	FAD	G	450	-	-	7/30/50/50	0/6/6/6
2	UDP	I	400	-	-	0/16/32/32	0/2/2/2

The worst 5 of 43 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
3	J	450	FAD	C2A-N3A	4.18	1.38	1.32
3	A	450	FAD	C2A-N3A	4.11	1.38	1.32
3	Н	450	FAD	C2A-N3A	4.07	1.38	1.32
3	F	450	FAD	C4X-N5	4.03	1.38	1.30
3	J	450	FAD	C4X-N5	4.00	1.38	1.30

The worst 5 of 150 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
3	В	450	FAD	N3A-C2A-N1A	-5.52	120.05	128.68
3	G	450	FAD	N3A-C2A-N1A	-5.51	120.07	128.68
3	С	450	FAD	N3A-C2A-N1A	-5.47	120.13	128.68
3	Е	450	FAD	N3A-C2A-N1A	-5.47	120.13	128.68
3	F	450	FAD	N3A-C2A-N1A	-5.46	120.15	128.68

There are no chirality outliers.

5 of 58 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	400	UDP	PA-O3A-PB-O3B
2	G	400	UDP	PA-O3A-PB-O3B
2	Н	400	UDP	PA-O3A-PB-O2B
2	Н	400	UDP	PA-O3A-PB-O3B
3	A	450	FAD	O4'-C4'-C5'-O5'

There are no ring outliers.

10 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	400	UDP	2	0
3	A	450	FAD	1	0
3	J	450	FAD	1	0
3	С	450	FAD	1	0


Continued on next page...




Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Е	400	UDP	1	0
3	В	450	FAD	2	0
3	F	450	FAD	1	0
3	Н	450	FAD	1	0
3	Е	450	FAD	1	0
3	G	450	FAD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































































Torsions

Rings

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	366/397~(92%)	-0.44	5 (1%) 75 73	29, 38, 55, 90	1 (0%)
1	В	364/397~(91%)	-0.34	8 (2%) 62 60	30, 39, 59, 102	1 (0%)
1	С	362/397~(91%)	-0.34	3 (0%) 86 84	31, 44, 63, 78	1 (0%)
1	D	362/397~(91%)	-0.11	10 (2%) 53 51	33, 49, 72, 94	1 (0%)
1	E	364/397~(91%)	-0.22	9 (2%) 57 55	20, 49, 70, 107	0
1	F	363/397 (91%)	-0.40	6 (1%) 70 68	31, 41, 57, 82	1 (0%)
1	G	363/397~(91%)	-0.37	4 (1%) 80 79	32, 46, 67, 93	0
1	Н	363/397 (91%)	-0.40	2 (0%) 89 88	20, 41, 61, 91	0
1	I	364/397~(91%)	-0.46	2 (0%) 91 89	30, 41, 60, 75	1 (0%)
1	J	361/397~(90%)	-0.29	6 (1%) 70 68	30, 43, 64, 80	1 (0%)
All	All	3632/3970 (91%)	-0.34	55 (1%) 73 72	20, 42, 66, 107	7 (0%)

The worst 5 of 55 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	154	GLU	5.2
1	В	357	GLN	4.2
1	Е	154	GLU	4.1
1	F	154	GLU	4.1
1	В	155	GLN	4.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	FAD	G	450	53/53	0.96	0.13	38,50,57,61	0
2	UDP	F	400	25/25	0.97	0.11	38,41,46,50	0
3	FAD	A	450	53/53	0.97	0.15	31,37,42,45	0
3	FAD	С	450	53/53	0.97	0.16	36,46,57,61	0
3	FAD	D	450	53/53	0.97	0.17	38,49,62,66	0
3	FAD	Е	450	53/53	0.97	0.15	41,51,59,65	0
2	UDP	Е	400	25/25	0.97	0.16	37,44,53,61	0
3	FAD	I	450	53/53	0.97	0.12	34,42,47,52	0
3	FAD	J	450	53/53	0.97	0.14	37,43,54,58	0
2	UDP	J	400	25/25	0.98	0.15	32,38,48,52	0
2	UDP	С	400	25/25	0.98	0.12	35,41,51,59	0
3	FAD	В	450	53/53	0.98	0.14	32,39,44,52	0
2	UDP	D	400	25/25	0.98	0.15	$41,\!46,\!55,\!57$	0
2	UDP	A	400	25/25	0.98	0.15	34,38,46,52	0
2	UDP	В	400	25/25	0.98	0.12	33,39,49,51	0
3	FAD	F	450	53/53	0.98	0.14	31,40,46,47	0
2	UDP	G	400	25/25	0.98	0.12	38,40,53,59	0
3	FAD	Н	450	53/53	0.98	0.14	33,39,45,49	0
2	UDP	Н	400	25/25	0.98	0.12	35,42,50,55	0
2	UDP	I	400	25/25	0.98	0.13	35,40,51,55	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around UDP F 400:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around FAD A 450: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around FAD C 450: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around FAD E 450:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around UDP E 400:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around FAD I 450:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around FAD J 450:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around UDP J 400:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around FAD B 450:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around UDP D 400:


 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)




























6.5 Other polymers (i)

There are no such residues in this entry.

