



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 15, 2024 – 02:09 AM EST

PDB ID : 3NGU
Title : Structure of Leishmania NDKb complexed with ADP.
Authors : Trindade, D.M.; Sousa, T.A.C.B.; Tonoli, C.C.C.; Santos, C.R.; Arni, R.K.;
Ward, R.J.; Oliveira, A.H.C.; Murakami, M.T.
Deposited on : 2010-06-13
Resolution : 2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	151	
1	B	151	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nucleoside diphosphate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	151	Total 1176	C 748	N 203	O 219	S 6	19	1	0
1	B	151	Total 1173	C 746	N 203	O 219	S 5	17	0	0

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	B	1	Total 27	C 10	N 5	O 10	P 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	25	Total 25	O 25	0	0
3	B	14	Total 14	O 14	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleoside diphosphate kinase



- Molecule 1: Nucleoside diphosphate kinase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	123.21Å 123.21Å 137.98Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.00 – 2.29 32.82 – 2.29	Depositor EDS
% Data completeness (in resolution range)	94.3 (32.00-2.29) 94.4 (32.82-2.29)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.216 , 0.283 0.203 , 0.282	Depositor DCC
R_{free} test set	886 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	48.4	Xtrriage
Anisotropy	0.102	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2442	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	2.01	32/1206 (2.7%)	1.56	14/1629 (0.9%)
1	B	1.97	28/1200 (2.3%)	1.57	12/1621 (0.7%)
All	All	1.99	60/2406 (2.5%)	1.56	26/3250 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (60) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	52	LYS	CD-CE	-12.34	1.20	1.51
1	A	136	ASP	CB-CG	-11.33	1.27	1.51
1	A	53	ASP	CB-CG	-9.81	1.31	1.51
1	B	53	ASP	CB-CG	9.75	1.72	1.51
1	A	134	LYS	CG-CD	-9.56	1.20	1.52
1	B	123	GLU	CB-CG	8.82	1.69	1.52
1	B	74	CYS	CB-SG	-8.77	1.67	1.82
1	B	150	TYR	CD1-CE1	8.51	1.52	1.39
1	B	125	ALA	CA-CB	8.24	1.69	1.52
1	A	4	GLU	CG-CD	8.24	1.64	1.51
1	B	35	VAL	CA-CB	7.93	1.71	1.54
1	A	126	GLU	CG-CD	7.91	1.63	1.51
1	B	22	GLU	CG-CD	7.83	1.63	1.51
1	A	98	SER	CB-OG	-7.74	1.32	1.42
1	A	52	LYS	CA-CB	-7.68	1.37	1.53
1	B	28	GLU	CG-CD	7.62	1.63	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	27	PHE	CE2-CZ	7.47	1.51	1.37
1	A	133	PHE	CE2-CZ	7.47	1.51	1.37
1	A	27	PHE	CE2-CZ	7.39	1.51	1.37
1	A	66	TYR	CD1-CE1	7.06	1.50	1.39
1	B	141	TRP	CB-CG	7.03	1.62	1.50
1	B	23	ILE	CA-CB	6.85	1.70	1.54
1	A	123	GLU	CG-CD	6.73	1.62	1.51
1	B	73	VAL	CB-CG2	-6.51	1.39	1.52
1	B	80	LYS	CB-CG	6.51	1.70	1.52
1	A	11	LYS	CE-NZ	6.48	1.65	1.49
1	B	115	VAL	CB-CG2	-6.43	1.39	1.52
1	A	80	LYS	CG-CD	-6.29	1.31	1.52
1	A	115	VAL	CB-CG2	-6.28	1.39	1.52
1	B	28	GLU	CD-OE2	6.25	1.32	1.25
1	A	46	GLN	CG-CD	6.24	1.65	1.51
1	A	126	GLU	CD-OE1	6.11	1.32	1.25
1	A	33	LYS	CD-CE	6.07	1.66	1.51
1	B	77	TRP	CE3-CZ3	6.00	1.48	1.38
1	A	105	GLY	CA-C	5.96	1.61	1.51
1	A	115	VAL	CA-CB	5.91	1.67	1.54
1	A	90	LEU	C-O	-5.89	1.12	1.23
1	B	45	GLU	CB-CG	5.88	1.63	1.52
1	A	45	GLU	CD-OE1	5.83	1.32	1.25
1	A	113	ARG	CG-CD	5.79	1.66	1.51
1	A	19	LEU	CG-CD1	5.74	1.73	1.51
1	A	45	GLU	CD-OE2	5.74	1.31	1.25
1	A	92	ALA	C-O	5.73	1.34	1.23
1	B	133	PHE	CD2-CE2	5.64	1.50	1.39
1	B	66	TYR	CD1-CE1	-5.58	1.30	1.39
1	B	132	TRP	CB-CG	-5.55	1.40	1.50
1	A	72	ILE	CA-CB	5.49	1.67	1.54
1	A	150	TYR	CE2-CZ	5.47	1.45	1.38
1	B	81	ASN	CB-CG	5.43	1.63	1.51
1	B	116	CYS	CB-SG	-5.39	1.73	1.81
1	B	20	VAL	C-O	5.37	1.33	1.23
1	B	102	THR	CA-CB	5.34	1.67	1.53
1	B	12	PRO	CB-CG	5.33	1.76	1.50
1	A	81	ASN	CB-CG	5.27	1.63	1.51
1	A	66	TYR	CG-CD1	5.23	1.46	1.39
1	A	29	ARG	CZ-NH1	5.16	1.39	1.33
1	B	151	GLU	C-O	5.13	1.33	1.23
1	B	49	GLY	C-O	5.09	1.31	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	13	ASP	CB-CG	5.07	1.62	1.51
1	A	42	PRO	CB-CG	5.05	1.75	1.50

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	136	ASP	CB-CG-OD1	-19.83	100.46	118.30
1	B	53	ASP	CB-CG-OD1	-17.81	102.27	118.30
1	B	53	ASP	CB-CG-OD2	17.67	134.20	118.30
1	A	136	ASP	CB-CG-OD2	16.49	133.14	118.30
1	B	53	ASP	CA-CB-CG	-11.41	88.31	113.40
1	B	5	ARG	NE-CZ-NH2	-10.55	115.02	120.30
1	B	123	GLU	CA-CB-CG	-10.20	90.97	113.40
1	B	54	LEU	CA-CB-CG	7.92	133.52	115.30
1	A	136	ASP	CA-CB-CG	7.65	130.24	113.40
1	A	37	LEU	CA-CB-CG	6.67	130.65	115.30
1	A	5	ARG	NE-CZ-NH1	-6.41	117.10	120.30
1	B	26	ARG	NE-CZ-NH1	6.30	123.45	120.30
1	A	151	GLU	CA-C-O	6.12	132.95	120.10
1	B	134	LYS	CA-CB-CG	6.09	126.79	113.40
1	B	5	ARG	NH1-CZ-NH2	6.01	126.01	119.40
1	A	23	ILE	CG1-CB-CG2	5.94	124.46	111.40
1	B	127	ARG	NE-CZ-NH2	5.91	123.26	120.30
1	A	17	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	80	LYS	CB-CG-CD	5.46	125.80	111.60
1	A	29	ARG	NE-CZ-NH2	-5.41	117.59	120.30
1	A	35	VAL	CG1-CB-CG2	5.34	119.45	110.90
1	A	123	GLU	OE1-CD-OE2	-5.21	117.05	123.30
1	B	150	TYR	CB-CG-CD1	5.08	124.05	121.00
1	B	37	LEU	CA-CB-CG	5.01	126.83	115.30
1	A	103	ILE	CG1-CB-CG2	5.01	122.42	111.40
1	A	37	LEU	CB-CG-CD1	5.00	119.50	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	150	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1176	0	1169	32	0
1	B	1173	0	1164	32	0
2	A	27	0	12	0	0
2	B	27	0	12	2	0
3	A	25	0	0	1	0
3	B	14	0	0	2	0
All	All	2442	0	2357	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:12:PRO:CB	1:B:12:PRO:CG	1.76	1.45
1:A:42:PRO:CG	1:A:42:PRO:CB	1.75	1.37
1:A:90:LEU:HD21	1:A:116[B]:CYS:SG	1.95	1.06
1:A:60:PHE:HB3	1:A:61:PRO:HD3	1.54	0.90
1:B:43:THR:H	1:B:46:GLN:HE21	1.17	0.89
1:A:90:LEU:CD2	1:A:116[B]:CYS:SG	2.63	0.86
1:B:11:LYS:HB3	1:B:12:PRO:HD2	1.62	0.80
2:B:152:ADP:H8	2:B:152:ADP:H5'1	1.52	0.75
1:A:90:LEU:HD21	1:A:116[A]:CYS:HB3	1.71	0.71
1:B:43:THR:N	1:B:46:GLN:HE21	1.92	0.68
1:B:43:THR:H	1:B:46:GLN:NE2	1.89	0.68
1:B:12:PRO:HD3	1:B:72:ILE:HG22	1.76	0.65
1:A:95:PRO:HD3	1:A:111:VAL:HA	1.78	0.65
1:B:37:LEU:HD23	1:B:75:MET:HG2	1.78	0.65
1:A:40:LEU:N	1:A:40:LEU:HD23	2.11	0.65
1:A:125:ALA:O	1:A:129:ILE:HG13	1.96	0.64
1:B:35:VAL:CG2	1:B:76:VAL:HG12	2.29	0.63
1:B:143:SER:O	1:B:146:VAL:HG23	1.99	0.63
1:A:150:TYR:O	1:A:151:GLU:HB2	1.99	0.62
1:A:40:LEU:HD23	1:A:40:LEU:H	1.66	0.61
1:A:135:ALA:O	1:A:136:ASP:CB	2.47	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:PHE:HB3	1:A:61:PRO:CD	2.30	0.60
1:B:86:GLY:O	1:B:90:LEU:HG	2.02	0.60
1:B:90:LEU:HD23	1:B:103:ILE:HD12	1.84	0.58
1:B:104:ARG:NH2	1:B:116:CYS:O	2.32	0.58
1:A:103:ILE:HG21	1:A:116[B]:CYS:SG	2.44	0.58
1:B:76:VAL:HG22	1:B:129:ILE:HG12	1.86	0.56
1:A:94:ASN:HD22	1:A:96:ALA:H	1.54	0.56
1:A:40:LEU:N	1:A:40:LEU:CD2	2.69	0.54
1:A:5:ARG:HA	1:A:77:TRP:O	2.08	0.54
1:B:34:LEU:HD21	1:B:37:LEU:HG	1.88	0.54
1:B:117:HIS:CD2	1:B:117:HIS:C	2.82	0.52
1:A:7:PHE:HE1	1:A:74:CYS:HB3	1.74	0.51
1:A:37:LEU:HG	1:B:37:LEU:HD13	1.93	0.51
1:A:20:VAL:HG22	1:B:24:ILE:HG21	1.93	0.51
1:B:1:MET:CE	1:B:81:ASN:HD21	2.25	0.50
1:B:36:ALA:HB3	1:B:76:VAL:HB	1.93	0.49
1:A:12:PRO:HD3	1:A:72:ILE:HG22	1.94	0.49
1:A:22:GLU:HG2	3:A:158:HOH:O	2.13	0.49
1:B:146:VAL:CG2	3:B:163:HOH:O	2.61	0.49
1:A:11:LYS:HB3	1:A:12:PRO:CD	2.43	0.49
1:A:33:LYS:HG3	1:A:141:TRP:CZ2	2.47	0.49
1:B:43:THR:OG1	1:B:46:GLN:HG3	2.13	0.48
1:A:90:LEU:HD23	1:A:116[B]:CYS:SG	2.52	0.47
1:B:37:LEU:O	1:B:38:LYS:HB3	2.14	0.47
1:B:11:LYS:HB3	1:B:12:PRO:CD	2.40	0.46
1:A:94:ASN:ND2	1:A:96:ALA:H	2.14	0.46
1:B:99:GLN:O	1:B:105:GLY:HA3	2.16	0.45
1:B:94:ASN:ND2	1:B:96:ALA:H	2.15	0.45
2:B:152:ADP:H5'1	2:B:152:ADP:C8	2.43	0.45
1:A:135:ALA:O	1:A:136:ASP:HB2	2.16	0.45
1:A:30:LYS:HD2	1:A:89:LEU:HD11	1.99	0.44
1:B:31:GLY:HA3	1:B:149:ILE:HD13	1.99	0.44
1:B:129:ILE:HG22	1:B:130:ALA:N	2.32	0.44
1:A:99:GLN:HA	1:A:100:PRO:HD3	1.84	0.44
1:B:13:ASP:HB3	1:B:66:TYR:OH	2.16	0.44
1:B:35:VAL:HG23	1:B:76:VAL:HG12	1.97	0.44
1:A:51:TYR:CE2	1:A:63:LEU:HD11	2.53	0.43
1:A:8:ILE:HG23	1:A:8:ILE:HD12	1.79	0.43
1:A:38:LYS:HD3	1:A:133:PHE:CE1	2.54	0.42
1:B:146:VAL:HG21	3:B:163:HOH:O	2.19	0.41
1:B:87:ARG:HD2	1:B:87:ARG:HA	1.87	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:VAL:CG2	1:B:129:ILE:HG12	2.50	0.41
1:A:87:ARG:NH1	1:A:118:GLY:O	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	150/151 (99%)	141 (94%)	6 (4%)	3 (2%)	7	6
1	B	149/151 (99%)	139 (93%)	9 (6%)	1 (1%)	22	26
All	All	299/302 (99%)	280 (94%)	15 (5%)	4 (1%)	12	12

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	52	LYS
1	B	115	VAL
1	A	136	ASP
1	A	115	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/126 (101%)	113 (89%)	14 (11%)	6	7
1	B	126/126 (100%)	110 (87%)	16 (13%)	4	4
All	All	253/252 (100%)	223 (88%)	30 (12%)	5	5

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	PRO
1	A	35	VAL
1	A	39	ILE
1	A	40	LEU
1	A	66	TYR
1	A	84	LYS
1	A	89	LEU
1	A	90	LEU
1	A	94	ASN
1	A	97	ASP
1	A	98	SER
1	A	140	SER
1	A	146	VAL
1	A	151	GLU
1	B	2	SER
1	B	6	THR
1	B	37	LEU
1	B	40	LEU
1	B	53	ASP
1	B	56	SER
1	B	94	ASN
1	B	98	SER
1	B	99	GLN
1	B	115	VAL
1	B	123	GLU
1	B	126	GLU
1	B	127	ARG
1	B	142	THR
1	B	144	HIS
1	B	151	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	ASN
1	B	41	GLN
1	B	46	GLN
1	B	81	ASN
1	B	94	ASN
1	B	144	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	152	-	24,29,29	1.49	4 (16%)	29,45,45	2.18	8 (27%)
2	ADP	B	152	-	24,29,29	1.81	8 (33%)	29,45,45	2.18	10 (34%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	152	-	-	2/12/32/32	0/3/3/3
2	ADP	B	152	-	-	6/12/32/32	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	152	ADP	C5-C4	4.04	1.51	1.40
2	A	152	ADP	C5-C4	3.92	1.51	1.40
2	B	152	ADP	C2'-C1'	3.15	1.58	1.53
2	A	152	ADP	PA-O1A	2.81	1.60	1.50
2	B	152	ADP	C6-C5	2.71	1.53	1.43
2	B	152	ADP	C2-N3	2.62	1.36	1.32
2	B	152	ADP	C2-N1	2.41	1.38	1.33
2	B	152	ADP	PA-O5'	2.23	1.68	1.59
2	A	152	ADP	C8-N7	2.23	1.38	1.34
2	A	152	ADP	O4'-C1'	-2.19	1.38	1.41
2	B	152	ADP	O4'-C1'	2.11	1.44	1.41
2	B	152	ADP	C2'-C3'	2.09	1.59	1.53

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	152	ADP	N3-C2-N1	-5.34	120.34	128.68
2	A	152	ADP	O4'-C4'-C3'	-5.30	94.62	105.11
2	A	152	ADP	C2-N1-C6	5.18	127.61	118.75
2	B	152	ADP	O5'-C5'-C4'	4.92	125.94	108.99
2	B	152	ADP	O4'-C1'-C2'	-4.67	100.10	106.93
2	B	152	ADP	O3'-C3'-C2'	3.72	123.85	111.82
2	B	152	ADP	C3'-C2'-C1'	3.23	105.84	100.98
2	B	152	ADP	N3-C2-N1	-3.11	123.81	128.68
2	B	152	ADP	O3B-PB-O3A	3.04	114.82	104.64
2	B	152	ADP	O2'-C2'-C3'	2.99	121.48	111.82
2	A	152	ADP	C5-C6-N1	-2.70	114.23	120.35
2	B	152	ADP	C4-C5-N7	-2.58	106.71	109.40
2	B	152	ADP	C2-N1-C6	2.57	123.16	118.75
2	A	152	ADP	O3B-PB-O1B	2.41	120.11	110.68
2	B	152	ADP	C5-C6-N1	-2.34	115.05	120.35
2	A	152	ADP	O3'-C3'-C4'	-2.34	104.29	111.05
2	A	152	ADP	N6-C6-N1	2.23	123.21	118.57
2	A	152	ADP	C1'-N9-C4	-2.15	122.87	126.64

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	152	ADP	C5'-O5'-PA-O1A
2	B	152	ADP	PB-O3A-PA-O1A
2	B	152	ADP	PB-O3A-PA-O5'
2	B	152	ADP	C5'-O5'-PA-O2A
2	A	152	ADP	PB-O3A-PA-O1A
2	B	152	ADP	C5'-O5'-PA-O3A
2	A	152	ADP	PB-O3A-PA-O2A
2	B	152	ADP	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	152	ADP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	151/151 (100%)	-0.41	3 (1%) 65 71	42, 54, 68, 85	7 (4%)
1	B	151/151 (100%)	-0.22	2 (1%) 77 81	46, 60, 82, 89	8 (5%)
All	All	302/302 (100%)	-0.31	5 (1%) 70 76	42, 56, 75, 89	15 (4%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	66	TYR	5.3
1	A	130	ALA	2.3
1	A	151	GLU	2.2
1	A	56	SER	2.1
1	B	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ADP	B	152	27/27	0.80	0.21	78,92,106,109	0
2	ADP	A	152	27/27	0.98	0.07	52,58,69,73	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.