



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2023 – 07:13 AM EDT

PDB ID : 4H8J
Title : Structure of GluA2-LBD in complex with MES
Authors : Reiter, A.; Skerra, A.; Trauner, D.; Schiefner, A.
Deposited on : 2012-09-22
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	
1	B	276	
1	C	276	
1	D	276	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	261	2033	1293	339	387	14	0	0	0
1	B	259	2025	1290	337	384	14	0	1	0
1	C	261	2033	1293	339	387	14	0	0	0
1	D	258	2018	1286	336	382	14	0	0	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	GLY	-	expression tag	UNP P19491
A	-10	SER	-	expression tag	UNP P19491
A	-9	ALA	-	expression tag	UNP P19491
A	-8	MET	-	expression tag	UNP P19491
A	-7	GLY	-	expression tag	UNP P19491
A	-6	SER	-	expression tag	UNP P19491
A	-5	GLY	-	expression tag	UNP P19491
A	-4	ASN	-	expression tag	UNP P19491
A	-3	ASP	-	expression tag	UNP P19491
A	-2	THR	-	expression tag	UNP P19491
A	-1	SER	-	expression tag	UNP P19491
A	0	ARG	-	expression tag	UNP P19491
A	1	GLY	-	expression tag	UNP P19491
A	2	ALA	-	expression tag	UNP P19491
A	118	GLY	-	linker	UNP P19491
A	119	THR	-	linker	UNP P19491
B	-11	GLY	-	expression tag	UNP P19491
B	-10	SER	-	expression tag	UNP P19491
B	-9	ALA	-	expression tag	UNP P19491
B	-8	MET	-	expression tag	UNP P19491
B	-7	GLY	-	expression tag	UNP P19491

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	SER	-	expression tag	UNP P19491
B	-5	GLY	-	expression tag	UNP P19491
B	-4	ASN	-	expression tag	UNP P19491
B	-3	ASP	-	expression tag	UNP P19491
B	-2	THR	-	expression tag	UNP P19491
B	-1	SER	-	expression tag	UNP P19491
B	0	ARG	-	expression tag	UNP P19491
B	1	GLY	-	expression tag	UNP P19491
B	2	ALA	-	expression tag	UNP P19491
B	118	GLY	-	linker	UNP P19491
B	119	THR	-	linker	UNP P19491
C	-11	GLY	-	expression tag	UNP P19491
C	-10	SER	-	expression tag	UNP P19491
C	-9	ALA	-	expression tag	UNP P19491
C	-8	MET	-	expression tag	UNP P19491
C	-7	GLY	-	expression tag	UNP P19491
C	-6	SER	-	expression tag	UNP P19491
C	-5	GLY	-	expression tag	UNP P19491
C	-4	ASN	-	expression tag	UNP P19491
C	-3	ASP	-	expression tag	UNP P19491
C	-2	THR	-	expression tag	UNP P19491
C	-1	SER	-	expression tag	UNP P19491
C	0	ARG	-	expression tag	UNP P19491
C	1	GLY	-	expression tag	UNP P19491
C	2	ALA	-	expression tag	UNP P19491
C	118	GLY	-	linker	UNP P19491
C	119	THR	-	linker	UNP P19491
D	-11	GLY	-	expression tag	UNP P19491
D	-10	SER	-	expression tag	UNP P19491
D	-9	ALA	-	expression tag	UNP P19491
D	-8	MET	-	expression tag	UNP P19491
D	-7	GLY	-	expression tag	UNP P19491
D	-6	SER	-	expression tag	UNP P19491
D	-5	GLY	-	expression tag	UNP P19491
D	-4	ASN	-	expression tag	UNP P19491
D	-3	ASP	-	expression tag	UNP P19491
D	-2	THR	-	expression tag	UNP P19491
D	-1	SER	-	expression tag	UNP P19491
D	0	ARG	-	expression tag	UNP P19491
D	1	GLY	-	expression tag	UNP P19491
D	2	ALA	-	expression tag	UNP P19491
D	118	GLY	-	linker	UNP P19491

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	119	THR	-	linker	UNP P19491

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	S			
2	A	1	Total	12	6	1	4	1	0	0
2	C	1	Total	12	6	1	4	1	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	276	Total O 282 282	0	6
5	B	279	Total O 281 281	0	2
5	C	260	Total O 262 262	0	2
5	D	274	Total O 279 279	0	5

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate receptor 2

Chain A: 




- Molecule 1: Glutamate receptor 2

Chain B: 





- Molecule 1: Glutamate receptor 2

Chain C: 



- Molecule 1: Glutamate receptor 2

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.99Å 92.02Å 196.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.93 – 1.80 19.85 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.93-1.80) 100.0 (19.85-1.80)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.35 (at 1.80Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.173 , 0.221 0.174 , 0.222	Depositor DCC
R_{free} test set	5237 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	13.4	Xtrriage
Anisotropy	0.046	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9317	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.34 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.5449e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/2069	0.84	3/2780 (0.1%)
1	B	0.96	0/2064	0.84	1/2775 (0.0%)
1	C	0.93	0/2069	0.83	0/2780
1	D	0.96	0/2054	0.84	2/2762 (0.1%)
All	All	0.95	0/8256	0.84	6/11097 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	203	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	A	139	ASP	C-N-CA	-5.43	108.11	121.70
1	A	96	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	D	180	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	203	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	B	126	ASP	CB-CG-OD1	5.13	122.91	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2033	0	2061	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2025	0	2058	6	0
1	C	2033	0	2061	10	0
1	D	2018	0	2050	7	0
2	A	12	0	13	0	0
2	C	12	0	13	1	0
3	A	15	0	0	0	0
3	B	15	0	0	0	0
3	C	15	0	0	0	0
3	D	15	0	0	0	0
4	A	8	0	12	1	0
4	B	4	0	6	0	0
4	C	4	0	6	0	0
4	D	4	0	6	0	0
5	A	282	0	0	0	0
5	B	281	0	0	3	0
5	C	262	0	0	1	0
5	D	279	0	0	4	0
All	All	9317	0	8286	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:LYS:HE2	5:B:597:HOH:O	1.70	0.91
1:C:218:LYS:HE3	5:C:572:HOH:O	1.77	0.81
1:A:145:GLU:OE1	1:A:148:ARG:NH1	2.22	0.72
1:B:95:VAL:HG21	5:B:528:HOH:O	1.98	0.62
1:C:100:ILE:HD12	1:C:223:ALA:HB1	1.84	0.60
1:C:145:GLU:OE1	1:C:148:ARG:NH1	2.36	0.59
1:B:95:VAL:HG23	5:B:495:HOH:O	2.06	0.56
1:C:65:ASP:OD2	1:C:67:ASP:HB2	2.08	0.54
1:D:45:LYS:HD2	5:D:552:HOH:O	2.07	0.54
1:D:256:TYR:HE2	5:D:538:HOH:O	1.91	0.53
1:B:118:GLY:HA2	1:C:225:PRO:HB3	1.92	0.52
1:D:194:SER:O	1:D:198:GLU:HG3	2.10	0.50
1:D:81:GLY:HA2	5:D:450:HOH:O	2.11	0.50
1:C:65:ASP:O	1:C:69:LYS:HA	2.12	0.49
4:A:305:EDO:H12	1:D:70:ILE:HD12	1.98	0.46
1:D:30:GLU:HG2	5:D:607:HOH:O	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:241:LEU:HD22	1:B:246:LEU:HD23	1.99	0.44
1:C:209:MET:HE2	1:C:209:MET:HB2	1.83	0.43
1:C:203:ARG:HG2	1:C:203:ARG:HH11	1.83	0.43
1:A:166:GLU:HA	1:A:167:PRO:C	2.40	0.42
1:B:100:ILE:HD12	1:B:223:ALA:HB1	2.01	0.41
1:A:50:LYS:HE2	1:A:50:LYS:HB2	1.55	0.41
1:C:117:LYS:HG2	1:C:209:MET:CE	2.51	0.40
1:C:61:TYR:CE1	2:C:301:MES:H81	2.56	0.40
1:D:137:THR:HG22	1:D:191:LEU:HB2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/276 (94%)	258 (100%)	1 (0%)	0	100	100
1	B	258/276 (94%)	256 (99%)	2 (1%)	0	100	100
1	C	259/276 (94%)	256 (99%)	3 (1%)	0	100	100
1	D	256/276 (93%)	255 (100%)	1 (0%)	0	100	100
All	All	1032/1104 (94%)	1025 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/227 (96%)	215 (99%)	3 (1%)	67	59
1	B	218/227 (96%)	217 (100%)	1 (0%)	88	87
1	C	218/227 (96%)	217 (100%)	1 (0%)	88	87
1	D	217/227 (96%)	216 (100%)	1 (0%)	88	87
All	All	871/908 (96%)	865 (99%)	6 (1%)	84	81

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	PRO
1	A	109	LEU
1	A	218	LYS
1	B	109	LEU
1	C	109	LEU
1	D	109	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	303	-	4,4,4	0.34	0	6,6,6	0.84	0
2	MES	C	301	-	12,12,12	2.05	3 (25%)	14,16,16	1.64	3 (21%)
3	SO4	C	302	-	4,4,4	0.75	0	6,6,6	0.85	0
3	SO4	D	301	-	4,4,4	0.45	0	6,6,6	0.70	0
4	EDO	D	304	-	3,3,3	0.48	0	2,2,2	0.78	0
4	EDO	A	306	-	3,3,3	0.50	0	2,2,2	0.85	0
2	MES	A	301	-	12,12,12	1.64	2 (16%)	14,16,16	2.05	5 (35%)
4	EDO	C	305	-	3,3,3	0.44	0	2,2,2	1.13	0
4	EDO	B	304	-	3,3,3	0.42	0	2,2,2	0.32	0
3	SO4	A	302	-	4,4,4	0.44	0	6,6,6	1.09	1 (16%)
3	SO4	C	304	-	4,4,4	0.33	0	6,6,6	0.34	0
3	SO4	A	304	-	4,4,4	0.36	0	6,6,6	0.57	0
3	SO4	D	302	-	4,4,4	0.48	0	6,6,6	0.52	0
3	SO4	B	301	-	4,4,4	0.36	0	6,6,6	1.21	1 (16%)
3	SO4	D	303	-	4,4,4	0.51	0	6,6,6	1.37	1 (16%)
3	SO4	B	303	-	4,4,4	0.46	0	6,6,6	0.39	0
4	EDO	A	305	-	3,3,3	0.36	0	2,2,2	0.92	0
3	SO4	C	303	-	4,4,4	0.19	0	6,6,6	0.49	0
3	SO4	B	302	-	4,4,4	0.44	0	6,6,6	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MES	A	301	-	-	1/6/14/14	0/1/1/1
4	EDO	C	305	-	-	1/1/1/1	-
4	EDO	A	305	-	-	0/1/1/1	-
2	MES	C	301	-	-	2/6/14/14	0/1/1/1
4	EDO	B	304	-	-	0/1/1/1	-
4	EDO	D	304	-	-	0/1/1/1	-
4	EDO	A	306	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	301	MES	C8-S	5.45	1.85	1.77
2	A	301	MES	C8-S	3.91	1.83	1.77
2	A	301	MES	O1S-S	2.75	1.53	1.45
2	C	301	MES	O2S-S	2.55	1.52	1.45
2	C	301	MES	O1S-S	2.31	1.51	1.45

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	MES	O3S-S-C8	4.32	112.75	105.77
2	A	301	MES	C6-C5-N4	-3.35	105.02	110.10
2	C	301	MES	O1-C6-C5	-3.23	104.68	111.80
2	A	301	MES	C6-O1-C2	3.12	120.32	109.89
2	A	301	MES	O3S-S-O1S	-3.00	103.95	111.27
3	B	301	SO4	O4-S-O3	2.45	119.50	109.06
3	A	302	SO4	O4-S-O2	2.33	121.47	109.31
3	D	303	SO4	O4-S-O3	2.31	118.93	109.06
2	C	301	MES	O3S-S-C8	2.23	109.38	105.77
2	C	301	MES	O2S-S-C8	2.19	109.56	106.92
2	A	301	MES	O1-C6-C5	-2.14	107.07	111.80

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	MES	C8-C7-N4-C3
2	C	301	MES	C8-C7-N4-C3
4	C	305	EDO	O1-C1-C2-O2
4	A	306	EDO	O1-C1-C2-O2
2	C	301	MES	C8-C7-N4-C5

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	301	MES	1	0
4	A	305	EDO	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/276 (94%)	-0.32	5 (1%) 66 63	6, 14, 31, 44	0
1	B	259/276 (93%)	-0.33	4 (1%) 73 70	6, 13, 26, 41	0
1	C	261/276 (94%)	-0.27	7 (2%) 54 49	6, 14, 31, 51	0
1	D	258/276 (93%)	-0.35	3 (1%) 79 76	6, 13, 27, 40	0
All	All	1039/1104 (94%)	-0.31	19 (1%) 68 64	6, 13, 29, 51	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	262	GLY	8.1
1	C	66	ALA	5.5
1	A	66	ALA	4.9
1	C	140	SER	4.0
1	C	264	GLY	3.6
1	A	263	SER	3.5
1	C	4	LYS	3.5
1	C	67	ASP	3.4
1	D	4	LYS	3.3
1	B	66	ALA	3.1
1	C	263	SER	2.8
1	A	4	LYS	2.7
1	C	21	LYS	2.6
1	A	21	LYS	2.5
1	A	140	SER	2.5
1	B	67	ASP	2.3
1	B	4	LYS	2.2
1	D	67	ASP	2.2
1	D	259	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	D	303	5/5	0.81	0.30	34,36,40,41	0
4	EDO	C	305	4/4	0.82	0.13	23,24,25,30	0
4	EDO	A	306	4/4	0.83	0.14	23,26,26,28	0
3	SO4	B	302	5/5	0.89	0.30	31,32,36,38	0
3	SO4	B	303	5/5	0.90	0.35	39,40,42,44	0
4	EDO	D	304	4/4	0.93	0.10	18,21,25,26	0
4	EDO	A	305	4/4	0.94	0.11	18,24,25,30	0
3	SO4	B	301	5/5	0.94	0.26	25,31,34,35	0
3	SO4	D	302	5/5	0.94	0.32	35,36,40,40	0
3	SO4	A	304	5/5	0.94	0.32	35,36,40,41	0
2	MES	C	301	12/12	0.95	0.18	21,23,29,29	0
4	EDO	B	304	4/4	0.95	0.10	19,20,21,27	0
3	SO4	A	303	5/5	0.95	0.23	28,31,35,37	0
3	SO4	C	304	5/5	0.95	0.32	30,34,37,37	0
2	MES	A	301	12/12	0.96	0.15	16,21,23,24	0
3	SO4	D	301	5/5	0.96	0.23	25,26,32,32	0
3	SO4	C	302	5/5	0.96	0.13	18,20,22,22	0
3	SO4	C	303	5/5	0.97	0.22	30,31,33,35	0
3	SO4	A	302	5/5	0.97	0.13	20,22,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.