



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 05:04 AM EDT

PDB ID : 4J76
Title : Crystal Structure of a parasite tRNA synthetase, ligand-free
Authors : Koh, C.Y.; Kim, J.E.; Verlinde, C.L.M.J.; Hol, W.G.J.
Deposited on : 2013-02-12
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2096 (2.36-2.32)
Clashscore	141614	2193 (2.36-2.32)
Ramachandran outliers	138981	2159 (2.36-2.32)
Sidechain outliers	138945	2160 (2.36-2.32)
RSRZ outliers	127900	2067 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	 4% 87% 9%
1	B	409	 9% 82% 5% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophanyl-tRNA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	372	3061	1977	510	561	13	0	3	0
1	B	355	2928	1900	486	530	12	0	2	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP Q8IDW3
A	-3	PRO	-	expression tag	UNP Q8IDW3
A	-2	GLY	-	expression tag	UNP Q8IDW3
A	-1	SER	-	expression tag	UNP Q8IDW3
A	0	MET	-	expression tag	UNP Q8IDW3
B	-4	GLY	-	expression tag	UNP Q8IDW3
B	-3	PRO	-	expression tag	UNP Q8IDW3
B	-2	GLY	-	expression tag	UNP Q8IDW3
B	-1	SER	-	expression tag	UNP Q8IDW3
B	0	MET	-	expression tag	UNP Q8IDW3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	206	Total O 206 206	0	0
3	B	118	Total O 118 118	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tryptophanyl-tRNA synthetase



- Molecule 1: Tryptophanyl-tRNA synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	117.30Å 190.67Å 51.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.34 49.95 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-2.34) 99.5 (49.95-2.34)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.95 (at 2.34Å)	Xtrriage
Refinement program	REFMAC refmac_5.7.0032	Depositor
R, R_{free}	0.179 , 0.209 0.183 , 0.211	Depositor DCC
R_{free} test set	2515 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtrriage
Anisotropy	0.135	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6337	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3144	0.54	0/4237
1	B	0.38	0/3006	0.54	0/4053
All	All	0.38	0/6150	0.54	0/8290

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3061	0	3050	10	0
1	B	2928	0	2923	10	0
2	A	18	0	24	0	0
2	B	6	0	8	0	0
3	A	206	0	0	3	0
3	B	118	0	0	0	0
All	All	6337	0	6005	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:617:TYR:O	1:A:623:LYS:HE3	2.01	0.59
1:A:342:LEU:HD11	1:A:367:VAL:HG22	1.84	0.59
1:B:342:LEU:HD11	1:B:367:VAL:HG22	1.84	0.58
1:B:598:GLU:O	1:B:602:LYS:HG3	2.04	0.58
1:B:557:SER:HA	1:B:592:LEU:HD22	1.86	0.57
1:A:557:SER:HA	1:A:592:LEU:HD22	1.88	0.56
1:A:266:ARG:NH1	3:A:957:HOH:O	2.39	0.54
1:B:245:ILE:HG21	1:B:247:TYR:CZ	2.42	0.54
1:B:321:LEU:HD22	1:B:519:LEU:HD21	1.90	0.53
1:B:617:TYR:O	1:B:623:LYS:HE3	2.10	0.51
1:A:301:LYS:NZ	3:A:930:HOH:O	2.43	0.50
1:A:492:LYS:O	1:A:493:MET:C	2.52	0.49
1:A:319:GLY:HA3	1:A:493:MET:HE2	1.98	0.45
1:A:310:GLY:HA3	1:A:347:LYS:HE2	2.00	0.44
1:B:578:TYR:HB2	1:B:583:MET:CE	2.48	0.43
1:B:552:LEU:HD11	1:B:579:LYS:HB3	2.01	0.43
1:B:452:GLN:O	1:B:479:HIS:HA	2.19	0.42
1:B:578:TYR:HB2	1:B:583:MET:HE2	2.02	0.42
1:A:266:ARG:HG3	3:A:957:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	371/409 (91%)	367 (99%)	4 (1%)	0	100	100
1	B	351/409 (86%)	348 (99%)	3 (1%)	0	100	100
All	All	722/818 (88%)	715 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	346/377 (92%)	345 (100%)	1 (0%)	92	96
1	B	331/377 (88%)	328 (99%)	3 (1%)	78	87
All	All	677/754 (90%)	673 (99%)	4 (1%)	86	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	331	GLN
1	B	321	LEU
1	B	331	GLN
1	B	490	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	490	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	701	-	5,5,5	0.26	0	5,5,5	0.21	0
2	GOL	A	703	-	5,5,5	0.32	0	5,5,5	0.27	0
2	GOL	B	701	-	5,5,5	0.32	0	5,5,5	0.22	0
2	GOL	A	702	-	5,5,5	0.22	0	5,5,5	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	701	-	-	4/4/4/4	-
2	GOL	A	703	-	-	2/4/4/4	-
2	GOL	B	701	-	-	2/4/4/4	-
2	GOL	A	702	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	701	GOL	O1-C1-C2-C3
2	A	701	GOL	C1-C2-C3-O3
2	A	702	GOL	O1-C1-C2-O2
2	A	702	GOL	O1-C1-C2-C3
2	A	703	GOL	O1-C1-C2-C3
2	B	701	GOL	C1-C2-C3-O3
2	A	701	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	703	GOL	O1-C1-C2-O2
2	B	701	GOL	O2-C2-C3-O3
2	A	701	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	372/409 (90%)	0.06	15 (4%) 38 49	17, 33, 80, 144	0
1	B	355/409 (86%)	0.44	38 (10%) 6 9	17, 44, 114, 150	0
All	All	727/818 (88%)	0.25	53 (7%) 15 22	17, 39, 100, 150	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	244	GLY	11.1
1	B	580	LYS	8.1
1	B	583	MET	6.6
1	B	576	GLU	5.4
1	B	581	GLY	5.1
1	B	572	ASN	4.5
1	B	578	TYR	4.1
1	A	244	GLY	4.1
1	A	492	LYS	4.1
1	B	587	GLU	4.0
1	B	482	PHE	3.9
1	B	584	LEU	3.7
1	B	571	LEU	3.7
1	B	552	LEU	3.7
1	A	482	PHE	3.6
1	B	579	LYS	3.6
1	B	314	LEU	3.5
1	B	489	VAL	3.5
1	B	569	ASN	3.4
1	B	574	ILE	3.4
1	B	573	GLU	3.3
1	B	577	LYS	3.3
1	B	570	LYS	3.2
1	A	245	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	314	LEU	3.1
1	B	582	GLU	3.1
1	B	551	ASN	3.1
1	B	313	SER	3.0
1	A	576	GLU	2.8
1	B	575	GLY	2.7
1	B	245	ILE	2.6
1	B	532	LYS	2.6
1	A	490	ASN	2.6
1	A	312	SER	2.5
1	B	533	TYR	2.5
1	A	493	MET	2.5
1	B	632	MET	2.5
1	B	512	HIS	2.5
1	B	534	ALA	2.5
1	B	265	LYS	2.4
1	B	527	LYS	2.3
1	A	489	VAL	2.2
1	B	274	LYS	2.2
1	B	591	ILE	2.2
1	A	543	GLN	2.2
1	B	588	ILE	2.1
1	B	594	ASP	2.1
1	B	566	GLU	2.1
1	A	572	ASN	2.1
1	A	547	GLU	2.1
1	A	580	LYS	2.0
1	A	315	SER	2.0
1	B	530	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	701	6/6	0.80	0.22	67,69,70,71	0
2	GOL	A	703	6/6	0.89	0.16	69,70,72,72	0
2	GOL	A	701	6/6	0.91	0.16	48,49,51,52	0
2	GOL	A	702	6/6	0.92	0.18	46,56,60,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.