



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 19, 2024 – 04:08 PM EST

PDB ID : 4K6T
Title : Crystal structure of Ad37 fiber knob in complex with trivalent sialic acid inhibitor ME0385
Authors : Stehle, T.; Bauer, J.
Deposited on : 2013-04-16
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	194	
1	B	194	
1	C	194	
1	E	194	
1	F	194	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	194	 90% 5% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ACT	B	407	-	-	X	-

2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 9858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fiber protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	1466	943	231	286	6	0	5	0
1	B	186	1460	941	231	283	5	0	4	0
1	C	182	1459	940	230	284	5	0	8	0
1	E	185	1458	939	231	282	6	0	4	0
1	F	186	1448	927	232	284	5	0	3	0
1	G	185	1469	946	231	287	5	0	6	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	172	GLY	-	expression tag	UNP Q64823
A	173	ALA	-	expression tag	UNP Q64823
A	174	MET	-	expression tag	UNP Q64823
A	175	GLY	-	expression tag	UNP Q64823
A	176	SER	-	expression tag	UNP Q64823
B	172	GLY	-	expression tag	UNP Q64823
B	173	ALA	-	expression tag	UNP Q64823
B	174	MET	-	expression tag	UNP Q64823
B	175	GLY	-	expression tag	UNP Q64823
B	176	SER	-	expression tag	UNP Q64823
C	172	GLY	-	expression tag	UNP Q64823
C	173	ALA	-	expression tag	UNP Q64823
C	174	MET	-	expression tag	UNP Q64823
C	175	GLY	-	expression tag	UNP Q64823
C	176	SER	-	expression tag	UNP Q64823
E	172	GLY	-	expression tag	UNP Q64823
E	173	ALA	-	expression tag	UNP Q64823

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	174	MET	-	expression tag	UNP Q64823
E	175	GLY	-	expression tag	UNP Q64823
E	176	SER	-	expression tag	UNP Q64823
F	172	GLY	-	expression tag	UNP Q64823
F	173	ALA	-	expression tag	UNP Q64823
F	174	MET	-	expression tag	UNP Q64823
F	175	GLY	-	expression tag	UNP Q64823
F	176	SER	-	expression tag	UNP Q64823
G	172	GLY	-	expression tag	UNP Q64823
G	173	ALA	-	expression tag	UNP Q64823
G	174	MET	-	expression tag	UNP Q64823
G	175	GLY	-	expression tag	UNP Q64823
G	176	SER	-	expression tag	UNP Q64823

- Molecule 2 is N-acetyl-alpha-neuraminic acid (three-letter code: SIA) (formula: $C_{11}H_{19}NO_9$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
2	A	1	Total	C	N	O	0	0
			20	11	1	8		
2	B	1	Total	C	N	O	0	0
			20	11	1	8		
2	C	1	Total	C	N	O	0	0
			20	11	1	8		
2	E	1	Total	C	N	O	0	0
			20	11	1	8		
2	F	1	Total	C	N	O	0	0
			20	11	1	8		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	G	1	Total	C	N	O	0	0
			20	11	1	8		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Zn	0	0
			3	3		
3	B	3	Total	Zn	0	0
			3	3		
3	C	1	Total	Zn	0	0
			1	1		
3	E	2	Total	Zn	0	0
			2	2		
3	F	2	Total	Zn	0	0
			2	2		
3	G	3	Total	Zn	0	0
			3	3		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	G	1	Total C O 4 2 2	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	C	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0

- Molecule 6 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0

- Molecule 7 is 2,2',2''-[nitrilotris(methanediyl-1H-1,2,3-triazole-4,1-diyl)]triethanol (three-letter code: 1P0) (formula: C₁₅H₂₄N₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			28	15	10	3		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	Mg	0	0
			1	1		
8	C	2	Total	Mg	0	0
			2	2		
8	E	1	Total	Mg	0	0
			1	1		
8	F	2	Total	Mg	0	0
			2	2		

- Molecule 9 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	E	1	Total	C	O	0	0
			6	3	3		
9	G	1	Total	C	O	0	0
			6	3	3		
9	G	1	Total	C	O	0	0
			6	3	3		

- Molecule 10 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	G	1	Total Ca 1 1	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	143	Total O 143 143	0	0
11	B	146	Total O 146 146	0	0
11	C	131	Total O 131 131	0	0
11	E	129	Total O 129 129	0	0
11	F	127	Total O 127 127	0	0
11	G	146	Total O 146 146	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fiber protein




- Molecule 1: Fiber protein




- Molecule 1: Fiber protein




- Molecule 1: Fiber protein



- Molecule 1: Fiber protein



- Molecule 1: Fiber protein

Chain G:  90% 5% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.94Å 67.18Å 121.43Å 90.00° 96.81° 90.00°	Depositor
Resolution (Å)	37.26 – 2.00 37.26 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (37.26-2.00) 99.4 (37.26-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0025	Depositor
R, R_{free}	0.160 , 0.200 0.170 , 0.204	Depositor DCC
R_{free} test set	1615 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.6	Xtrriage
Anisotropy	0.306	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9858	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0119e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, 1P0, CL, ACT, EDO, GOL, SIA, CA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/1515	0.69	0/2064
1	B	0.61	0/1506	0.69	0/2052
1	C	0.56	0/1514	0.67	0/2062
1	E	0.57	0/1504	0.70	0/2046
1	F	0.54	1/1490 (0.1%)	0.66	1/2030 (0.0%)
1	G	0.59	0/1518	0.70	1/2069 (0.0%)
All	All	0.57	1/9047 (0.0%)	0.69	2/12323 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	343	TRP	CD2-CE2	5.37	1.47	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	304	ARG	NE-CZ-NH2	-5.49	117.56	120.30
1	G	318	ASP	CB-CG-OD1	5.00	122.80	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1466	0	1431	9	0
1	B	1460	0	1423	13	0
1	C	1459	0	1446	7	0
1	E	1458	0	1431	13	0
1	F	1448	0	1407	6	0
1	G	1469	0	1440	5	0
2	A	20	0	17	0	0
2	B	20	0	17	0	0
2	C	20	0	17	0	0
2	E	20	0	17	0	0
2	F	20	0	17	0	0
2	G	20	0	17	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
3	C	1	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
3	G	3	0	0	0	0
4	A	8	0	6	0	0
4	B	8	0	6	2	0
4	E	12	0	9	0	0
4	F	4	0	3	0	0
4	G	4	0	3	0	0
5	A	8	0	12	0	0
5	B	12	0	18	3	0
5	C	4	0	6	0	0
5	E	16	0	24	3	0
5	F	12	0	18	2	0
6	A	1	0	0	0	0
7	B	28	0	21	2	0
8	B	1	0	0	0	0
8	C	2	0	0	0	0
8	E	1	0	0	0	0
8	F	2	0	0	0	0
9	E	6	0	8	0	0
9	G	12	0	16	0	0
10	G	1	0	0	0	0
11	A	143	0	0	0	0
11	B	146	0	0	0	0
11	C	131	0	0	0	0
11	E	129	0	0	1	0
11	F	127	0	0	0	0
11	G	146	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	9858	0	8830	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:402:1P0:H19	7:B:402:1P0:NAR	1.75	0.99
7:B:402:1P0:NAR	7:B:402:1P0:CAN	2.36	0.89
1:E:285:GLY:H	5:E:407:EDO:H22	1.38	0.88
1:B:231:HIS:HE1	4:B:407:ACT:O	1.67	0.77
1:B:296:PRO:HA	1:B:297:SER:CB	2.22	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/194 (97%)	182 (96%)	7 (4%)	0	100	100
1	B	188/194 (97%)	179 (95%)	8 (4%)	1 (0%)	29	23
1	C	188/194 (97%)	180 (96%)	8 (4%)	0	100	100
1	E	187/194 (96%)	179 (96%)	7 (4%)	1 (0%)	29	23
1	F	187/194 (96%)	179 (96%)	8 (4%)	0	100	100
1	G	189/194 (97%)	182 (96%)	7 (4%)	0	100	100
All	All	1128/1164 (97%)	1081 (96%)	45 (4%)	2 (0%)	47	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	297	SER
1	E	299	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/171 (95%)	162 (99%)	1 (1%)	86	90
1	B	160/171 (94%)	158 (99%)	2 (1%)	69	74
1	C	166/171 (97%)	162 (98%)	4 (2%)	49	51
1	E	162/171 (95%)	159 (98%)	3 (2%)	57	61
1	F	161/171 (94%)	158 (98%)	3 (2%)	57	61
1	G	164/171 (96%)	161 (98%)	3 (2%)	59	63
All	All	976/1026 (95%)	960 (98%)	16 (2%)	65	67

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	248[A]	LEU
1	G	187	TRP
1	E	278	THR
1	F	353	GLU
1	E	187	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	298	ASN
1	C	253	ASN
1	G	319	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 54 ligands modelled in this entry, 22 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SIA	A	401	7	20,20,21	0.89	0	24,28,31	1.22	3 (12%)
5	EDO	F	407	-	3,3,3	0.60	0	2,2,2	0.28	0
2	SIA	F	403	-	20,20,21	0.88	0	24,28,31	1.47	5 (20%)
4	ACT	A	406	3	3,3,3	0.67	0	3,3,3	1.12	0
4	ACT	E	405	-	3,3,3	0.64	0	3,3,3	1.08	0
4	ACT	E	404	-	3,3,3	0.76	0	3,3,3	0.69	0
4	ACT	F	404	3	3,3,3	0.92	0	3,3,3	0.29	0
5	EDO	E	407	-	3,3,3	0.46	0	2,2,2	0.60	0
9	GOL	G	406	-	5,5,5	0.34	0	5,5,5	0.35	0
9	GOL	G	407	-	5,5,5	0.39	0	5,5,5	0.83	0
2	SIA	E	403	-	20,20,21	0.79	0	24,28,31	1.28	3 (12%)
7	1P0	B	402	2	30,30,30	2.76	10 (33%)	27,39,39	3.18	13 (48%)
5	EDO	F	406	-	3,3,3	0.41	0	2,2,2	0.36	0
2	SIA	G	404	-	20,20,21	0.73	0	24,28,31	1.22	3 (12%)
5	EDO	E	408	-	3,3,3	0.24	0	2,2,2	0.44	0
5	EDO	B	409	-	3,3,3	0.44	0	2,2,2	0.32	0
4	ACT	E	406	-	3,3,3	0.91	0	3,3,3	0.46	0
2	SIA	C	401	7	20,20,21	0.78	0	24,28,31	1.11	1 (4%)
5	EDO	F	405	-	3,3,3	0.58	0	2,2,2	0.47	0
2	SIA	B	401	7	20,20,21	0.70	0	24,28,31	1.16	2 (8%)
4	ACT	B	407	3	3,3,3	0.92	0	3,3,3	0.57	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	E	410	-	3,3,3	0.44	0	2,2,2	0.37	0
5	EDO	B	408	-	3,3,3	0.57	0	2,2,2	0.15	0
9	GOL	E	411	-	5,5,5	0.49	0	5,5,5	0.40	0
5	EDO	A	408	-	3,3,3	0.56	0	2,2,2	0.25	0
4	ACT	B	406	3	3,3,3	0.63	0	3,3,3	1.20	0
4	ACT	G	405	10	3,3,3	0.69	0	3,3,3	0.84	0
4	ACT	A	405	3	3,3,3	0.91	0	3,3,3	0.37	0
5	EDO	A	407	-	3,3,3	0.58	0	2,2,2	0.22	0
5	EDO	B	410	-	3,3,3	0.43	0	2,2,2	0.45	0
5	EDO	E	409	-	3,3,3	0.52	0	2,2,2	0.19	0
5	EDO	C	403	-	3,3,3	0.45	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIA	A	401	7	-	0/18/34/38	0/1/1/1
5	EDO	F	407	-	-	0/1/1/1	-
2	SIA	F	403	-	-	2/18/34/38	0/1/1/1
5	EDO	E	407	-	-	1/1/1/1	-
9	GOL	G	406	-	-	3/4/4/4	-
9	GOL	G	407	-	-	0/4/4/4	-
2	SIA	E	403	-	-	1/18/34/38	0/1/1/1
7	1P0	B	402	2	-	12/21/21/21	0/3/3/3
5	EDO	F	406	-	-	1/1/1/1	-
2	SIA	G	404	-	-	0/18/34/38	0/1/1/1
5	EDO	E	408	-	-	1/1/1/1	-
5	EDO	B	409	-	-	1/1/1/1	-
2	SIA	C	401	7	-	0/18/34/38	0/1/1/1
5	EDO	F	405	-	-	0/1/1/1	-
2	SIA	B	401	7	-	4/18/34/38	0/1/1/1
5	EDO	E	410	-	-	1/1/1/1	-
5	EDO	B	408	-	-	0/1/1/1	-
9	GOL	E	411	-	-	2/4/4/4	-
5	EDO	A	408	-	-	1/1/1/1	-
5	EDO	A	407	-	-	0/1/1/1	-
5	EDO	B	410	-	-	1/1/1/1	-
5	EDO	E	409	-	-	1/1/1/1	-
5	EDO	C	403	-	-	0/1/1/1	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	402	1P0	NAR-NAU	-7.95	1.20	1.34
7	B	402	1P0	NAQ-NAT	-6.71	1.22	1.34
7	B	402	1P0	NAU-NBB	-5.12	1.24	1.34
7	B	402	1P0	NAS-NAZ	-4.22	1.26	1.34
7	B	402	1P0	CAJ-NAZ	3.66	1.54	1.47

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	B	402	1P0	CAW-CAN-NAY	-8.62	94.62	112.78
7	B	402	1P0	CAX-CAO-NAY	-6.27	99.57	112.78
7	B	402	1P0	NAR-NAU-NBB	5.49	111.45	107.31
7	B	402	1P0	NAQ-NAT-NBA	5.24	111.27	107.31
7	B	402	1P0	CAF-CAX-NAR	-4.04	105.34	111.34

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	402	1P0	OAA-CAG-CAJ-NAZ
7	B	402	1P0	CAG-CAJ-NAZ-CAD
7	B	402	1P0	CAG-CAJ-NAZ-NAS
7	B	402	1P0	NAY-CAM-CAV-NAP
7	B	402	1P0	NAY-CAO-CAX-CAF


There are no ring outliers.


8 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	F	407	EDO	1	0
5	E	407	EDO	1	0
7	B	402	1P0	2	0
5	E	408	EDO	2	0
5	B	409	EDO	2	0
5	F	405	EDO	1	0
4	B	407	ACT	2	0
5	B	410	EDO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	186/194 (95%)	-0.36	2 (1%) 80 79	15, 23, 39, 68	0
1	B	186/194 (95%)	-0.11	7 (3%) 40 39	17, 24, 39, 72	0
1	C	182/194 (93%)	0.03	7 (3%) 40 39	17, 28, 50, 75	0
1	E	185/194 (95%)	-0.16	4 (2%) 62 60	16, 24, 43, 80	0
1	F	186/194 (95%)	-0.02	9 (4%) 30 29	17, 28, 46, 75	0
1	G	185/194 (95%)	-0.38	0 100 100	14, 22, 36, 46	0
All	All	1110/1164 (95%)	-0.17	29 (2%) 56 54	14, 25, 43, 80	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	181	TYR	4.6
1	F	277	SER	4.4
1	C	277	SER	4.3
1	F	180	LYS	3.6
1	E	332	GLY	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
9	GOL	E	411	6/6	0.64	0.25	41,54,57,60	0
2	SIA	G	404	20/21	0.73	0.27	37,57,75,77	0
5	EDO	F	405	4/4	0.77	0.15	31,36,47,50	0
2	SIA	E	403	20/21	0.82	0.23	39,58,72,74	0
7	1P0	B	402	28/28	0.82	0.21	19,44,56,68	0
5	EDO	E	410	4/4	0.82	0.16	54,56,58,59	0
2	SIA	F	403	20/21	0.83	0.27	33,56,69,76	0
5	EDO	B	408	4/4	0.85	0.12	46,55,59,62	0
4	ACT	G	405	4/4	0.86	0.28	48,52,53,54	0
2	SIA	A	401	20/21	0.86	0.14	23,30,42,44	0
2	SIA	B	401	20/21	0.87	0.17	24,34,45,55	0
5	EDO	E	407	4/4	0.87	0.14	34,36,39,47	0
4	ACT	E	405	4/4	0.88	0.20	27,30,38,46	0
5	EDO	A	407	4/4	0.89	0.15	23,28,32,37	0
5	EDO	F	406	4/4	0.89	0.13	33,42,44,45	0
5	EDO	E	409	4/4	0.89	0.20	48,49,54,57	0
8	MG	F	409	1/1	0.89	0.09	36,36,36,36	0
3	ZN	E	402	1/1	0.89	0.08	87,87,87,87	0
3	ZN	F	402	1/1	0.90	0.12	73,73,73,73	0
5	EDO	C	403	4/4	0.90	0.09	34,36,50,53	0
5	EDO	F	407	4/4	0.91	0.19	28,29,40,50	0
5	EDO	A	408	4/4	0.91	0.13	31,32,42,51	0
3	ZN	G	403	1/1	0.91	0.08	80,80,80,80	0
5	EDO	B	409	4/4	0.91	0.13	26,34,36,40	0
2	SIA	C	401	20/21	0.92	0.13	26,33,43,44	0
3	ZN	B	405	1/1	0.92	0.10	81,81,81,81	0
6	CL	A	409	1/1	0.92	0.10	48,48,48,48	0
9	GOL	G	406	6/6	0.92	0.21	32,44,53,57	0
9	GOL	G	407	6/6	0.92	0.12	35,43,53,57	0
5	EDO	B	410	4/4	0.93	0.12	31,32,36,39	0
8	MG	C	405	1/1	0.93	0.07	37,37,37,37	0
8	MG	F	408	1/1	0.94	0.07	41,41,41,41	0
4	ACT	E	404	4/4	0.94	0.20	27,33,33,36	0
3	ZN	A	404	1/1	0.94	0.07	68,68,68,68	0
8	MG	C	404	1/1	0.94	0.09	40,40,40,40	0
4	ACT	B	407	4/4	0.94	0.10	24,31,34,40	0
4	ACT	E	406	4/4	0.95	0.21	24,36,44,53	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ACT	B	406	4/4	0.96	0.08	23,24,29,35	0
3	ZN	C	402	1/1	0.96	0.04	37,37,37,37	0
8	MG	B	411	1/1	0.96	0.04	28,28,28,28	0
4	ACT	F	404	4/4	0.96	0.10	25,25,32,34	0
3	ZN	B	404	1/1	0.96	0.17	71,71,71,71	0
8	MG	E	412	1/1	0.97	0.09	39,39,39,39	0
4	ACT	A	405	4/4	0.97	0.10	15,22,31,40	0
10	CA	G	408	1/1	0.97	0.07	45,45,45,45	0
5	EDO	E	408	4/4	0.98	0.14	31,41,41,41	0
4	ACT	A	406	4/4	0.98	0.14	18,21,22,27	0
3	ZN	G	402	1/1	0.99	0.04	35,35,35,35	0
3	ZN	F	401	1/1	0.99	0.05	33,33,33,33	0
3	ZN	E	401	1/1	0.99	0.06	21,21,21,21	1
3	ZN	G	401	1/1	0.99	0.04	28,28,28,28	0
3	ZN	B	403	1/1	1.00	0.03	40,40,40,40	0
3	ZN	A	403	1/1	1.00	0.05	23,23,23,23	0
3	ZN	A	402	1/1	1.00	0.03	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SIA G 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA E 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around 1P0 B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA F 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SIA A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SIA B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.